

Environmental Sensitivity Index Maps

A Guide to Coastal Resources at Risk to Spilled Oil

Chesapeake Bay: Volume 5 Lower Potomac River




Office of Response and Restoration

September 2016

NOAA
Environmental Sensitivity
Index Maps

A Guide to Coastal Resources at Risk to Spilled Oil




**Chesapeake Bay and the Outer Coasts of
Maryland and Virginia**



Chesapeake Bay: Volumes 1 and 2
(Chesapeake North:
Upper Bay and Choptank River)

Chesapeake Bay: Volume 3
(Chesapeake West:
Upper Potomac River)

Chesapeake Bay: Volumes 4 and 5
(Central Bay:
Patuxent River and Lower Potomac River)



W
E
N
S

1:100,000 Grid Tiles
1:50,000 Grid Tiles

10 5 0 10
KM
MILES

CONTENTS

Introduction	iv
Shoreline Habitat Mapping	iv
Sensitive Biological Resources	vi
Marine Mammals	viii
Birds	xi
Herpetofauna	xv
Fish	xvii
Invertebrates	xxii
Benthic	xxv
Habitats	xxvii
Invasive Species	xxix
Terrestrial Mammals	xxix
Wildlife Rehabilitation	xxix
Human Use Resources	xxx
Geographic Information System	xxxiii
Acknowledgements	xxxiii
Appropriate Use of Atlas and Data	xxxiv
Species List	xxxv
Shoreline Descriptions	xl
Exposed, Rocky Shores: ESI = 1A	xl
Exposed, Solid Man-Made Structures: ESI = 1B	xli
Exposed Scarps and Steep Slopes (Clay): ESI = 2B	xlii
Fine- to Medium-Grained Sand Beaches: ESI = 3A	xliii
Scarps and Steep Slopes (Sand): ESI = 3B	xliv
Coarse-Grained Sand Beaches: ESI = 4	xlv
Mixed Sand and Gravel Beaches: ESI = 5	xlii
Gravel Beaches: ESI = 6A	xlvii
Riprap: ESI = 6B	xlviii
Boulder Rubble: ESI = 6D	xlix
Exposed Tidal Flats: ESI = 7	l
Sheltered Scarps (Bedrock/Mud/Clay): ESI = 8A	li
Sheltered, Solid Man-Made Structures: ESI = 8B	lii
Sheltered Riprap: ESI = 8C	liii
Sheltered, Rocky, Rubble Shores: ESI = 8D	liv
Sheltered Tidal Flats: ESI = 9A	lv
Vegetated Low Banks: ESI = 9B	lvi
Salt- and Brackish-Water Marshes: ESI = 10A	lvii
Freshwater Marshes: ESI = 10B	lviii
Swamps: ESI = 10C	lix
Scrub and Shrub Wetlands: ESI = 10D	lx

Environmental Sensitivity Index Maps (Maps 1-11 contained in Volume 1;
Maps 12-16, 18-21 contained in Volume 2; Maps 17, 22-23, 30-33, 42-45 contained in Volume 3;
Maps 25-29, 37-41 contained in Volume 4; Maps 24, 34-36, 46-48 contained in Volume 6)

Map 49: Human Use Resources	1127
Map 49A: Biological Resources.....	1131
Map 49B: Biological Resources.....	1137
Map 49C: Biological Resources	1143
Map 49D: Biological Resources.....	1149
Map 50: Human Use Resources	1155
Map 50A: Biological Resources.....	1159
Map 50B: Biological Resources.....	1165
Map 50C: Biological Resources	1171
Map 50D: Biological Resources.....	1177
Map 51: Human Use Resources	1185
Map 51A: Biological Resources.....	1189
Map 51B: Biological Resources.....	1195
Map 51C: Biological Resources	1201
Map 51D: Biological Resources.....	1207
Map 52: Human Use Resources	1213
Map 52A: Biological Resources.....	1217
Map 52B: Biological Resources.....	1223
Map 52C: Biological Resources	1229

Maps 53-54 contained in Volume 3; Maps 55-56 contained in Volume 6

Map 57: Human Use Resources	1235
Map 57A: Biological Resources.....	1239
Map 57B: Biological Resources.....	1247
Map 57C: Biological Resources	1255
Map 57D: Biological Resources	1263
Map 58: Human Use Resources	1269
Map 58A: Biological Resources.....	1273
Map 58B: Biological Resources.....	1281
Map 58C: Biological Resources	1289
Map 58D: Biological Resources	1297
Map 59: Human Use Resources	1303
Map 59A: Biological Resources.....	1307
Map 59B: Biological Resources.....	1315
Map 60: Human UseResources	1321
Map 60A: Biological Resources.....	1325
Map 60B: Biological Resources.....	1331
Map 60C: Biological Resources	1337
Map 61: Human Use Resources	1343
Map 61A: Biological Resources.....	1347
Map 61B: Biological Resources.....	1353
Map 61C: Biological Resources	1359
Map 62 contained in Volume 3; Maps 63-65 contained in Volume 7	
Map 66: Human Use Resources	1365
Map 66A: Biological Resources.....	1369
Map 66B: Biological Resources.....	1375
Map 66C: Biological Resources	1381
Map 66D: Biological Resources	1389
Map 67: Human Use Resources	1397
Map 67A: Biological Resources.....	1401
Map 67B: Biological Resources.....	1407
Map 67C: Biological Resources	1413
Map 67D: Biological Resources	1419

Maps 68-70 contained in Volume 3;
 Maps 71-72, 78-79, 86-87 contained in Volume 7;
 Maps 73-77, 80-85 contained in Volume 8;
 Maps 88-97 contained in Volume 9

ENVIRONMENTAL SENSITIVITY INDEX: CHESAPEAKE BAY AND THE OUTER COASTS OF MARYLAND AND VIRGINIA

INTRODUCTION

This Environmental Sensitivity Index (ESI) atlas was developed for the land and marine and coastal areas of Chesapeake Bay and the outer coasts of Maryland and Virginia. The atlas covers the entire coastal area of Maryland and Virginia from the Pennsylvania and Maryland border to the north in the Susquehanna River, from the Delaware and Maryland border to the east at Fenwick Island, to Washington District of Columbia to the west at the Potomac River, and to the Virginia and North Carolina border to the south in Back Bay. The Maryland and Virginia ESI extends 12 nautical miles seaward. The inland extent of the study area includes lands and waters ranging from five to ten nautical miles from the shoreline of all tidal waters. The study area covers all Atlantic Ocean coastal bays, including (from north to south): Assawoman Bay, Isle of Wight Bay, Sinepuxent Bay, Newport Bay, Chincoteague Bay, Toms Cove, Metompkin Bay, Hog Island Bay, Cobb Bay, South Bay, Fishermans Inlet, and Back Bay. In addition, the study area encompasses all of Chesapeake Bay and major tidal tributary rivers, including (from north to south along the western Bay shoreline): Susquehanna River (south of the Pennsylvania and Maryland border), Back River, Patapsco River, Magothy River, Severn River, Rhode River, West River, Herring Bay, Patuxent River, Potomac River (north to Washington D.C.), Rappahannock River (west to Fredericksburg), York River, Mattaponi River (west to White Oak Landing), Pamunkey River (west to Hampstead), James River (west to Richmond), Appomattox River (west to the I-95 corridor), Nansemond River (west to Hwy 58), Elizabeth River (east to Hwy 168), North Landing River (south to the North Carolina border), Lynnhaven River, Wolfsnare Creek, and Linkhorn Bay. Also included in the study area are major tidal tributary rivers along the eastern Bay shoreline, including (from north to south): Elk River (and Back Creek to the Maryland and Delaware border), Sassafras River, Chester River, Eastern Bay, Wye River, Tred Avon River, Choptank River (east to Providence Landing), Fishing Bay, Nanticoke River (east to the Maryland and Delaware border), Manokin River (east to Kings Creek), Back Creek (east to Hwy 13), Big Annemessex River (east to Hwy 413), Pocomoke River (east to Pocomoke City), Beasley Bay, Pompco Creek, Deep Creek, Onancock Creek, Pungoteague Creek, Nandua Creek, Curratuck Creek, Craddock Creek, Occohannock Creek, Nassawaddox Creek, Hungars Creek, Cherrystone Inlet, and Longs Pond. The ESI represents a compilation of information about three main categories: shoreline habitats, sensitive biological resources, and human-use resources. Though the data will be useful for many shoreline applications, the goal of the ESI data is to present a concise summary of resources that may be particularly vulnerable to spilled oil. The intent of the data should caveat other uses. As an example, the ESI is not intended to present a catalog or comprehensive listing of species present in an area, rather the focus is on species particularly sensitive to oiling and life stages where vulnerability may increase.

SHORELINE HABITAT MAPPING

The shoreline and classifications were fully updated using the following sources and methods. The Chesapeake Bay and the outer coasts of Maryland and Virginia shoreline was derived from the integration of the Virginia Institute of Marine Science (VIMS) Center for Coastal Resources Management (CCRM) Tidal Marsh Inventory (TMI, 2010-2013); the National Oceanic and Atmospheric Administration (NOAA) Continually Updated Shoreline Product (CUSP, 1990-2015) and National Shoreline composite (2003-2015); the U.S. Geological Survey (USGS) high resolution National Hydrography Dataset (NHD, 2002-2014); the U.S. Fish and Wildlife Service (USFWS) National Wetlands Inventory (NWI) dataset (1974-2010); and manual digitization at 1:4,000 from BING Bird's Eye and Aerial imagery (2003-2014), ESRI Basemap World Imagery (2011-2015), and Google Earth aerial imagery (2013-2015). The most recent shoreline was utilized where available.

The intertidal shoreline habitats were classified based on the USFWS NWI wetland polygons and low-altitude oblique and nadir aerial imagery from: BING Bird's Eye, BING Aerial, ESRI Basemap, and Google Earth.

The ESI shoreline classification and ranking scale has been used to assess vulnerability of shoreline to spilled oil since the mid-1970s. Rankings range from 1 – least vulnerable, to 10 – most vulnerable, with a variety of qualifiers unique to the geographic region. The scale incorporates the following considerations:

- 1) Shoreline type (substrate, grain size, tidal elevation, origin)
- 2) Exposure to wave and tidal energy
- 3) Biological productivity and sensitivity
- 4) Ease of cleanup

Prediction of the behavior and persistence of oil in intertidal habitats is based on an understanding of the dynamics of the coastal environments, not just the substrate type and grain size. The intensity of energy expended upon a shoreline by wave action, tidal currents, and river currents directly affects the persistence of stranded oil. The need for shoreline cleanup activities is determined, in part, by the slowness of natural processes in removal of oil stranded on the shoreline. The potential for biological injury and ease of cleanup of spilled oil are also important factors in the ESI shoreline ranking. Thus, shorelines exposed to high levels of physical energy, such as wave action and tidal currents, and low biological activity rank low on the scale, whereas sheltered shorelines with associated high biological activity have the highest ranking. The shoreline types delineated for Maryland and Virginia, presented in order of increasing sensitivity to spilled oil, are listed below.

- | | |
|--|--|
| 1A) Exposed, Rocky Shores | 7) Exposed Tidal Flats |
| 1B) Exposed, Solid Man-Made Structures | 8A) Sheltered Scarps (Bedrock/Mud/Clay) |
| 2B) Exposed Scarps and Steep Slopes (Clay) | 8B) Sheltered, Solid Man-Made Structures |
| 3A) Fine- to Medium- Grained Sand Beaches | 8C) Sheltered Riprap |
| 3B) Scarps and Steep Slopes (Sand) | 8D) Sheltered, Rocky, Rubble Shores |
| 4) Coarse-Grained Sand Beaches | 9A) Sheltered Tidal Flats |
| 5) Mixed Sand and Gravel Beaches | 9B) Vegetated Low Banks |
| 6A) Gravel Beaches | 10A) Salt- and Brackish-Water Marshes |
| 6B) Riprap | 10B) Freshwater Marshes |
| 6D) Boulder Rubble | 10C) Swamps |
| | 10D) Scrub and Shrub Wetlands |

For each of these shoreline types, a photo and description of the physical attributes, predicted oil behavior, and response considerations are included at the end of the introductory pages.

SENSITIVE BIOLOGICAL RESOURCES

Biological information presented in this atlas was collected, compiled, and reviewed with the assistance of biologists and resource managers from the following agencies:

- Center for Conservation Biology (CCB)
- Maryland Department of Natural Resources (MD DNR)
- National Park Service (NPS), Assateague Island National Seashore
- NPS Northeast Coastal and Barrier Network (NCBN)
- NPS Northeast Region
- National Oceanic and Atmospheric Administration (NOAA)
- North Carolina Division of Marine Fisheries – Habitat Protection Section, Albemarle-Pamlico National Estuary Program
- Saltmarsh Habitat and Avian Research Program (SHARP)
- The Nature Conservancy (TNC)
- Virginia Aquarium and Marine Science Center (VAMSC)
- Virginia Commonwealth University (VCU)
- Virginia Department of Conservation and Recreation (VA DCR)
- Virginia Department of Game and Inland Fisheries (VDGIF)
- Virginia Institute of Marine Science (VIMS), College of William and Mary
- Virginia Marine Resources Commission (MRC)
- U.S. Fish and Wildlife Service (USFWS), Division of Migratory Bird Management (DMBM)
- USFWS National Wildlife Refuge (NWR) System

The above organizations provided the majority of the biological information included in the atlas. Other participating organizations will be featured in the sources table and cited in the metadata accompanying the digital product.

The biological resources shown in this atlas were extracted from the ESI GIS data compiled for this region. The extracted features were mapped at scale of 1:50,000 and appear on the maps referenced by a combination of number and letter. For example, Map 1B will show the biological features in conjunction with the ESI shoreline. The biology maps on these maps is “layered” in the PDF maps. This allows the user to turn off the biological features to more clearly see the underlying shoreline and habitat polygons.

The data published date appearing at the bottom of the maps and on the cover page reflect when the data collection and compilation was completed. This atlas represents those data and was published September 2016.

KEY BIOLOGICAL FEATURES ON ESI MAPS

- 1) Occurrences of animal and plant species that are at risk to spilled oil or may be impacted during a spill response are represented in the database by polygons, points, and lines.
- 2) To avoid clutter, the front of the map features occurrences that cover less than 10 kilometers of the map extent. A Map ID is associated with each of these polygonal, linear or point features.
- 3) Each map includes a tabular report summarizing the species found in the area. Features that are shown on the map are referenced by their Map ID. Features that cover more than 10 kilometers are presented in the report as Widespread in Mapped Area. Species occurrences that appear in the database as General Distribution are listed in a third category, Also Present in Mapped Area. To fully understand the diversity of species present, ALL sections of the map report should be reviewed.
- 4) Associated with each species in the table is the state (S) and federal (F) protected status as threatened (T) or endangered (E), as well as concentration, seasonality, and life-history information. Federal listings were provided by USFWS. State listings were provided by MD DNR and VDGIF.
- 5) The table includes a Mapping Qualifier with each species record (see table of mapping qualifiers and guidelines below). The mapping qualifier should help users understand particular vulnerabilities associated with the map data.
- 6) Feature level source information is included in the GIS database used to create these maps. The GIS data also provides the extent polygons for all mapped features; it can be queried, filtered, and used with other GIS datasets.
- 7) Species have been divided into groups and subgroups based on their behavior, morphology, taxonomic classification, and spill vulnerability and sensitivity. The icons below reflect this grouping scheme.
- 8) Colors depicting monthly seasonality roughly reflect Winter/Spring/Summer/Fall, but are primarily intended to ease readability.

MARINE MAMMAL

	Dolphin
	Manatee
	Pinniped
	Whale

INVERTEBRATE

	Bivalve
	Cephalopod
	Crab
	Gastropod

BIRD

	Alcid/Pelagic
	Diving
	Gull/Tern
	Passerine

TERRESTRIAL MAMMAL

	Bat
	Small Mammal
	Ungulate

FISH

	Fish
	Nursery

HABITAT

	Upland/Wetland/Plant
---	----------------------

HERPETOFAUNA

	Amphibian
	Turtle

Mapping Qualifiers and Guidelines

Element	Qualifier	Guidelines
All	Concentration Area	Areas where concentrations are considerably higher than other records of the same species in the area of interest.
All	General Distribution	Used for broad, general distributions of species that are often mapped to landscape- or habitat-scale features.
All	Vulnerable Occurrence	Intended for records of rare species with discrete occurrences, where the conservation value of the species should be highlighted for spill response.
Birds, Herpetofauna, Marine Mammals, Fish, Invertebrates	Migration	Used when an area is a known staging area of high importance to the species for birds; and/or areas are potential or known migration corridors in the marine environment for other elements.
Birds, Herpetofauna	Nesting	Applicable to all nesting birds and herps. Should represent known nesting areas rather than all potential nesting habitat.
Birds	Rafting	Similar to 'Concentration Area' qualifier, but specific to large on-water concentrations.
Birds	Wintering	Designates known areas of importance to wintering birds.
Benthic	High Ecological Value	For use in areas where benthic organisms provide high ecological services, high quality habitat, or known areas of high biodiversity.
Fish and Invertebrates	Harvest Area	May be used as a qualifier for distributions in special cases, where the general distribution was not mapped and/or widespread and the distribution of the harvested resources is used to depict important areas.
Fish and Invertebrates	Nursery Area	Refers to specific areas of known importance to early life history stages (e.g., larvae, juveniles) of a species.
Fish and Invertebrates	Spawning Area	Areas where animals are spawning. Spawning is loosely defined as the release of gametes or eggs from the adult.

MARINE MAMMALS

Marine mammals depicted in this atlas include whales, dolphins, seals, and manatees. Species that are federally and state listed, and those that are vulnerable to oiling and response activities, are specifically emphasized. Marine mammal occurrence and seasonality information displayed in this atlas are based on data from the Ocean Biogeographic Information System (OBIS) and Global Biodiversity Information Facility (GBIF), NOAA Fisheries Marine Mammal Stock Assessment Reports (SARS), and correspondence with local resource experts from Virginia Aquarium and Marine Science Center and other agencies (see expert contact list). Both hardcopy and digital data sources are listed below and included in the metadata.

Four of the marine mammal species included in this ESI atlas are currently listed as endangered at both the federal and state level. These include the North Atlantic right whale (*Eubalaena glacialis*, FE, VA/MD SE), fin whale (*Balaenoptera physalus*, FE, VA/MD SE), humpback whale (*Megaptera novaeangliae*, FE, VA/MD SE) and the Florida manatee (*Trichechus manatus latirostris*, FE, VA SE), a sub-species of the West Indian manatee. Two of these species currently have federal proposals under review in 2016 that may change their federal statuses within the boundaries of this ESI, by either removing them from the ESA

listing altogether (humpback whale) or reclassifying them from endangered to threatened (Florida manatee). These proposals were under review at the time this atlas was published (hence, both are still considered endangered); however, their listings may change so it is important to review their listings prior to any planning or response activities.

North Atlantic right whales may be present in both Maryland and Virginia marine waters from October to May. The waters off the mid-Atlantic coast are an important migration corridor between their wintering and calving areas, located off the southeastern United States, to their summer feeding and nursery grounds in New England waters. Right whales are known to primarily occur in coastal and shelf waters, although movements into deeper waters have been observed.

Fin whales can be found off the U.S. Atlantic coast from Cape Hatteras, NC northward and are the dominant large cetacean in this region. Although typically observed over continental shelf waters eastward of the 12 nautical mile boundary of the ESI, they may be present in shallower waters closer to the coastline from November through April.

Humpback whales may be present in Maryland and Virginia coastal waters throughout the year, but are primarily observed during the winter months as they migrate between their winter calving grounds in the West Indies and their summer feeding grounds in the Gulf of Maine and areas further north. Not all humpbacks migrate every winter, and significant numbers can be found in mid- and high-latitude regions along the Atlantic coast during this time. During their migration, humpback whales tend to stay near the surface of the ocean. They are known to enter Chesapeake Bay, which may be a supplemental feeding ground for this species, especially for juveniles; however, most sightings occur near the mouth of the bay and the surrounding marine waters.

The Florida manatee was mapped from information provided by peer-reviewed literature and local resource experts. This species is an irregular visitor to the Chesapeake Bay region and has, on rare occasion, been observed in the Maryland portion of the Chesapeake Bay, as far north as the Susquehanna River. They are a warm-water species (68°F or above) that feed on seagrasses and other aquatic plants, and can be found in marine, brackish and freshwater habitats. They usually travel along shorelines and within channels (i.e., Intracoastal Waterway) and may be present in shallow nearshore waters during the summer and fall months when water temperatures are above 68°F. Most of the reported sightings in Chesapeake Bay have been south of the Potomac River, with the majority of them occurring in the waters proximal to Norfolk, VA.

Other marine mammals were mapped using information provided by local marine mammal experts. Literature and thirty years of observation data from the SARs reports, OBIS, and GBIF were also reviewed. Bottlenose dolphins are the only species likely to occur regularly in inshore and offshore areas and may be present year round. Other species of cetaceans and pinnipeds that may occur in Chesapeake Bay and/or the coastal or continental shelf waters of Maryland and Virginia that were mapped in this atlas include gray seals, harbor seals, harbor porpoises, and short-beaked common dolphins.

In addition to those species mapped, other pelagic cetacean and pinniped species can occasionally move into the area without predictable occurrence. Species have been reported via stranding data or other observations, but are not included due to their relative rarity of occurrence within the mapped 'Area of Interest' (AOI) for this atlas.

Expert contacts for Chesapeake Bay and the outer coasts of Maryland and Virginia marine mammals*:

Name	Agency	City	Phone	Species
Susan Barco	VAMSC	Virginia Beach, VA	757/385-6476	Cetaceans, Pinnipeds and Sirenia
Mark Swingle	VAMSC	Virginia Beach, VA	757/437-6022	Cetaceans, Pinnipeds and Sirenia
Jennifer Dittmar	National Aquarium, Baltimore	Baltimore, MD	410/986-2377	Cetaceans and Pinnipeds
Amanda Weschler	MD DNR	Oxford, MD	410/226-5193	Cetaceans and Pinnipeds
Kevin Holcomb	USFWS	Chincoteague, VA	757/336-6122	Chincoteague and Wallops Island NWRs
Alex Wilke	TNC	Nassa-wadox, VA	757/442-3049	Eastern Shore species
Ruth Boettcher	VDGIF	Machi-pongo, VA	757/709-0766	Eastern Shore species

*Note: this list is not meant to represent all marine mammal experts for the region.

Major Data Sources Used: Marine Mammals

Barco, S.G., W.A. McLellan, J.M. Allen, R.A. Asmutis-Silia, R. Mallon-Day, E.M. Meagher, D.A. Pabst, J. Robbins, R.E. Seton, W.M. Swingle, M.T. Weinrich, and P.J. Clapham. 2002. Population identity of humpback whales (*Megaptera novaeangliae*) in the waters of the US mid-Atlantic states. Journal of Cetacean Research Management 4(2): 135-141.

Biodiversity Research Institute. 2015. Wildlife Densities and Habitat Use Across Temporal and Spatial Scales on the Mid-Atlantic Continental Shelf. Online vector digital data accessed on January 26, 2016 at: <http://www.briloon.org/mabs/data>.

Biodiversity Research Institute. 2015. Baseline Wildlife Studies in Atlantic Waters Offshore of Maryland. Online vector digital data accessed on January 26, 2016 at: <http://www.briloon.org/mabs/data>.

Cummings, E. W., D.A. Pabst, J.E. Blum, S.G. Barco, S.J. Davis, V.G. Thayer, N. Adimey, and W.A. McLellan. 2014. Spatial and Temporal Patterns of Habitat Use and Mortality of the Florida Manatee (*Trichechus manatus latirostris*) in the Mid-Atlantic States of North Carolina and Virginia from 1991 to 2012. Aquatic Mammals 40(2): 126-138.

Engelhardt, A., J. Aschettino, T.A. Jefferson, M. Richlen, and D. Engelhardt. 2015. Occurrence, Distribution, and Density of Marine Mammals Near Naval Station Norfolk & Virginia Beach, VA: Annual Progress Report. Final Report. Prepared for U.S. Fleet Forces Command. Submitted to Naval Facilities Engineering Command Atlantic, Norfolk, Virginia, under Contract No. N62470-10-3011, Task Orders 031 and 043, issued to HDR Inc., Virginia Beach, Virginia. 7 August 2015.

Engelhardt, A., M. Richlen, T.A. Jefferson, and D. Engelhardt. 2014. Occurrence, Distribution, and Density of Marine Mammals Near Naval Station Norfolk & Virginia Beach, VA: Annual Progress Report. Prepared for U.S. Fleet Forces Command. Submitted to Naval Facilities Engineering Command (NAVFAC) Atlantic, Norfolk, Virginia, under Contract No. N62470-10-3011, Task Orders 031 and 043, issued to HDR Inc., Norfolk, Virginia. 22 July 2014.

Global Biodiversity Information Facility (GBIF). 2015. Observation data for Cetacea, Phocidae and Sirenia in Maryland and Virginia from 1986-2015. Spreadsheet (.csv). Online data accessed on January 27, 2016 from: <http://www.gbif.org/>.

Halpin, P.N., A.J. Read, E. Fujioka, B.D. Best, B. Donnelly, L.J. Hazen, C. Kot, K. Urian, E. LaBrecque, A. Dimatteo, J. Cleary, C. Good, L.B. Crowder, and K.D. Hyrenbach. 2009. OBIS-SEAMAP: The world data center for marine mammal, sea bird, and sea turtle distributions. Oceanography. 22(2): 104-115. Spreadsheet (.csv). Online data accessed on December 17, 2015 from <http://seamap.env.duke.edu/>.

Roberts, J. J., B.D. Best, L. Mannucci, E. Fujioka, P.N. Halpin, D.L. Palka, L.P. Garrison, K.D. Mullin, T.V.N. Cole, C.B. Khan, W.A. McLellan, D.A. Pabst, and G.G. Lockhart. 2016. Habitat-based cetacean density models for the US Atlantic and Gulf of Mexico. Scientific reports, 6 (#22615). <http://seamap.env.duke.edu/models/Duke-EC-GOM-2015/>.

Waring G.T., E. Josephson, K. Maze-Foley, and P.E. Rosel, editors. 2015. U.S. Atlantic and Gulf of Mexico Marine Mammal Stock Assessments -- 2014. NOAA Tech Memo NMFS NE 231, 361 p. doi: 10.7289/V5TQ5ZH0. Accessed at <http://www.nmfs.noaa.gov/pr/sars/region.htm>.

BIRDS

Birds displayed in this atlas include: alcids, diving birds, gulls, terns, passerines, pelagic birds, raptors, shorebirds, wading birds, and waterfowl. Species that are federally and state listed, and coastal nesting, roosting, and migratory staging locations are specifically emphasized. Bird occurrence information displayed in this atlas is based on information gathered at workshops and via phone/email correspondence with local resource experts from VDGIF, MD DNR, NPS, USFWS, CCB, and TNC. Additional hardcopy and digital sources are listed below and included in the metadata.

Shorebirds, diving birds, gulls and terns – Survey data on locations of breeding, migrating, and/or wintering shorebirds, diving birds, gulls, and terns were provided by various agencies via shapefiles, spreadsheets, primary literature, and expert local knowledge, and was supplemented with information from eBird. In general, data from the various data sources were compiled and either mapped to habitat, park or refuge, or specific locations, per expert recommendations. Additional source information is provided in the accompanying data tables and metadata.

Colonial waterbird nesting areas – For Virginia, VDGIF provided nesting data for colonial waterbirds from 2013 that originated from various partner organizations including: CCB, VDGIF, TNC, VA DCR, USFWS, USGS, and College of William and Mary. Each nesting colony location was buffered by 200 meters (m), and where buffered nest locations overlapped, buffers were dissolved to form single polygons, and counts from the dissolved colonies were summed. Number of nesting pairs for each species for each colony and aggregated colony were reported in the ESI.

For Maryland, MD DNR provided nesting data for colonial waterbirds from surveys conducted annually during 2004-2013. Each nesting colony location was buffered by 200 m, and where buffered nest locations overlapped, buffers were dissolved to form single polygons, and counts from the dissolved colonies were summed. Number of nesting pairs for each species for each colony and aggregated colony were reported in the ESI.

Additionally, CCB provided nesting heron and egret survey data for the entire Chesapeake Bay drainage area (including MD), also from 2013. Where the heron/egret colonies overlapped with the VA colonial waterbird colonies, the heron/egret counts were added to the existing colony. Elsewhere, the heron and egret survey locations were mapped as points with counts of pairs for each species for each point location. Overall, species mapped using colonial waterbird datasets in VA include: black-crowned night-heron, black skimmer (MD SE), brown pelican, Caspian tern, cattle egret, common tern, double-crested cormorant, Forster's tern, glossy ibis, great black-backed gull, great blue heron, great egret, green heron, gull-billed tern (VA ST, MD SE), herring gull, laughing gull, least tern (MD ST), little blue heron, royal tern, sandwich tern, snowy egret, tricolored heron, white ibis, and yellow-crowned night-heron.

Wintering and nesting waterfowl – Tens of thousands of waterfowl migrate through Chesapeake Bay and Atlantic coastal bays with numbers of southbound transients peaking October through December and northbound transients peaking February through April. High numbers of scoters, red-throated loon, northern gannet, and other seabirds frequently concentrate at the mouth of Chesapeake Bay, Tangiers Sound, and the mouth of the Choptank River from November through April. Chesapeake Bay marsh islands and coastal bays provide habitat for the highest concentrations of nesting American black duck in the region. Waterfowl data for Virginia and Maryland were provided by VDGIF and USFWS and included mid-winter waterfowl surveys (2004-2015) and expert knowledge from VDGIF and MD DNR.

In Maryland, diving ducks and geese along the Chesapeake Bay were mapped into Areas of Critical State Concern for waterfowl (defined under Maryland's State Planning enabling legislation, article 88C). In coastal bays these species were mapped with a 250-m offshore buffer based on MD DNR surveys (2013-2014). Dabbling ducks were mapped based on USFWS mid-winter survey blocks and MD DNR coastal bay surveys with a 250-m onshore/offshore buffer. In Virginia, species were mapped according to areas defined by VDGIF when compiling aerial survey data for USFWS mid-winter surveys (2004-2015) and expert knowledge. Counts were binned into Present, (<10), 10s, 100s, 1,000s, or 10,000s.

Nesting waterfowl occurrences in Maryland were derived from MD DNR expert knowledge and data from the Second Atlas of the Breeding Birds of Maryland and the District of Columbia compiled by U.S. Geological Survey; they are mapped as a 250-m onshore buffer. In Virginia, American black duck nesting distributions were mapped based on VDGIF expert knowledge; mallard and Canada goose were mapped using species' presence during mid-winter surveys in or near marsh habitats; and wood duck were mapped using species' presence during mid-winter surveys in swamps and freshwater marshes. Nesting concentrations for waterfowl are represented as Present, Low, Moderate, High, or Very High. Onshore buffers that intersected with larger areas of freshwater or salt-and brackish-water marsh included those areas as part of the mapped distribution. Numbers from wintering surveys are likely underestimates, since surveys do not necessarily occur during each species' months of peak abundance.

Atlantic Ocean and Chesapeake Bay seabirds – Distribution of pelagic birds, gulls, terns, diving birds, alcids, phalaropes, and waterfowl in open waters of Chesapeake Bay and tributaries and the Atlantic Ocean (nearshore and offshore) were generated from discussions with resource experts, reports, and survey data. Species in open waters of Chesapeake Bay and the Atlantic Ocean were mapped from compiled aerial and shipboard survey data from USFWS that included: USFWS Atlantic Coast Wintering Sea Duck Surveys (ACWSD; preliminary survey in 2008, full surveys 2009-11), USFWS Atlantic Marine Assessment Program for Protected Species (AMAPPS) seabird surveys (preliminary survey in 2010, full-coast surveys in summer 2011, spring and fall 2012, winter 2014), NOAA EcoMon surveys, and Department of Energy (DOE)/Biological Research Institute (BRI) surveys (2012-2014). Distributions in Chesapeake Bay tributaries were mapped from USFWS midwinter surveys (2004-2015), BRI reports (2015), and eBird records from 2004-2016. Based on frequency of occurrence, abundance in these areas, and expert advice, data were grouped to create divisions throughout Chesapeake Bay and its tributaries, at the mouth of Chesapeake Bay, and as offshore polygons made as either a 0-5 nautical mile (nm) offshore buffer or a 0-12 nm offshore buffer. Counts were binned into Present (<10), 10s, 100s, 1,000s, or 10,000s.

Raptors – Bald eagle nest survey data (2015) were provided by CCB for Virginia. Nest locations were mapped as points in the ESI. Recent bald eagle nest survey data was not available for MD, as the state no longer surveys for the species. Historical data (1988-2010) of MD nest locations were mapped as points with the last observation date noted in the ESI concentration field. According to local experts (D. Brinker, L. Davidson), bald eagle nests are common in the state and the historical dataset is incomplete and does not fully cover where the species currently nests. VDGIF provided a Bald Eagle Concentration and Roosts (BECAR) dataset that features concentration areas along shorelines that have been identified through surveys to support a high density of non-breeding bald eagles in summer and/or winter. Roost sites are sites that have been documented as repeatedly used as night-time resting places for multiple bald eagles, and are displayed as both polygons (based on the VDGIF BECAR data set) and points (based on the CCB data set). Peregrine falcon and osprey nest sites were mapped based on expert knowledge and/or nesting survey data. Northern harrier were mapped based on expert knowledge.

Secretive marsh birds – Black rail (VA SE) detection data for Virginia were provided by CCB for 2007-2008 and 2014. Data from both datasets largely overlapped, so the datasets were combined for the ESI. Points were buffered by 200 m. Patch analysis data for clapper rail, saltmarsh sharp-tailed sparrow, seaside sparrow, and willet were provided by SHARP. The patch analysis used habitat data from NWI to delineate marsh patches, and field surveys of the target species to model species densities in each patch. Species densities were converted to number of birds per square kilometer and mapped to the appropriate marsh patches. Additionally, there were four SHARP survey sites on the western side of the Chesapeake Bay that had detected clapper rails in numbers too low for patch analysis. These survey sites were mapped as 50-m buffer polygons. NPS NCBN provided survey data of the following marsh birds on Assateague Island: clapper rail, king rail, saltmarsh sharp-tailed sparrow, and seaside sparrow. Each survey location with birds detected was mapped with a 200-m buffer. Where overlap occurred between SHARP data and NCBN data, SHARP data were used because they contained more detailed spatial and concentration information.

Rare birds - A few rare bird species were mapped using data supplied by MD DNR from their Ecologically Significant Areas database. Data from this program came in the form of large generalized polygons that often include the presence of more than one species. Bird species mapped from this data set include: American bittern, black rail (MD SE, VA SE), black skimmer (MD SE), common moorhen, gull-billed tern (MD SE, VA ST), least bittern, peregrine falcon (VA ST), and royal tern (MD SE).

Note that locations of nesting, wintering, and/or migratory sites, species composition within polygons, and particularly concentration values, are based on a compilation of observations made over multi-year periods and are not meant to accurately reflect ‘current’ conditions in the case of an event. Survey limitations and adjustments in protocols over the years, changes in shoreline geomorphology (particularly on small/ephemeral islands), weather, and other ecological factors contribute to the condition of nesting colonies and migratory or other concentrations at any given time. Also, note that concentrations vary throughout the multi-month nesting, migratory, and wintering periods listed in the seasonality table. Please contact local resource experts in the event of a spill or if data are to be used for any reason other than spill planning or response.

Expert contacts for Chesapeake Bay and the outer coasts of Maryland and Virginia birds*:

Name	Agency	City	Phone	Species
Ruth Boettcher	VDGIF	Machipongo, VA	757/709-0766	Birds of Virginia Eastern Shore
Bryan Watts	CCB	Williamsburg, VA	757/221-2247	Birds of Virginia and Chesapeake Bay
Alex Wilke	TNC	Nassawadox, VA	757/442-3049	Birds of Virginia Eastern Shore
Kevin Holcomb	USFWS	Chincoteague, VA	757/336-6122	Chincoteague and Wallops Island NWRs
Pam Denmon	USFWS	Cape Charles, VA	757/331-2760	Eastern Shore of Virginia and Fisherman Island NWRs
Lauren Billodeaux Mowbray	USFWS	Virginia Beach, VA	757/301-7329	Back Bay NWR and Eastern Virginia Rivers NWRC
Doug Brewer	USFWS	Virginia Beach, VA	757/301-7329	Back Bay NWR
Geralyn Mireles	USFWS	Virginia Beach, VA	757/301-7329	Back Bay NWR
Jack Kumer	NPS	Berlin, MD	410/629-6070	Assateague Island NS
Bill Hulslander	NPS	Berlin, MD	410/629-6061	Assateague Island NS

Name	Agency	City	Phone	Species
Dave Brinker	MD DNR	Annapolis, MD	410/744-8939	Maryland colonial-waterbirds
Lynn Davidson	MD DNR	Annapolis, MD	410/260-8563	Digital data provider
Gary Costanzo	VDGIF	Charles City, VA	804/367-1000	Waterfowl
Ben Lewis	VDGIF	Charles City, VA	804/367-1000	Waterfowl
Bill Harvey	MD DNR	Cambridge, MD	410/221-8838	Waterfowl
Don Webster	MD DNR	Cambridge, MD	410/221-8838	Waterfowl
Larry Hindman	MD DNR (Retired)	Cambridge, MD	-	Waterfowl
Paul Padding	USFWS	Laurel, MD	301/497-5851	Waterfowl
Tim Jones	USFWS	Laurel, MD	301/497-5674	Seabirds
Emily Silverman	USFWS	Laurel, MD	301/497-5801	Seabirds
Caleb Spiegel	USFWS	Hadley, MA	413/253-8490	Seabirds
Matt Whitbeck	USFWS	Cambridge, MD	410/221-2034	Chesapeake Marshlands NWR Complex

*Note: this list is not meant to represent all bird experts for the region.

Major Data Sources Used: Birds

Biodiversity Research Institute. 2014. Mid-Atlantic Baseline Studies: Boat surveys 2002-2014, vector digital data.

Bradshaw, D. 2008. George Washington Birthplace National Monument Avian Inventory. Technical Report NPS/NER/NRTR-2008.112 National Park Service. Philadelphia, PA, 66 pp.

Breeding Bird Atlas Explorer (online resource). 2016. U.S. Geological Survey Patuxent Wildlife Research Center. Data compiled from: Maryland and the District of Columbia Breeding Bird Atlas 2002-2006. Maryland Ornithological Society. Available at: <http://www.pwrc.usgs.gov/bba> (Accessed: November 2015).

Department of Energy. 2014. Seabird Surveys 2007-2014, vector digital data.

eBird. 2016. eBird: An online database of bird distribution and abundance [web application]. eBird, Ithaca, New York. Available at: <http://www.ebird.org>. (Accessed: March 2016).

Iliff, M.J., R.F. Ringler, and J.L. Stasz. 1996. Field List of the Birds of Maryland, Third Edition. Maryland Ornithological Society, Baltimore, MD, 53 pp.

MD DNR. 2015. Coastal Bay Waterfowl Surveys 2013-2015, vector digital data.

MD DNR. 2015. Colonial Waterbird Quarter mile Buffer, Chesapeake Bay AOI, vector digital data.

MD DNR. 2015. Waterfowl Areas, vector digital data.

NOAA. Atlantic Marine Assessment Program and Protected Species surveys 2011-2014, vector digital data.

Reese, J.G. and Weterings, R. 2013. Transient Waterfowl in Chesapeake Bay. Unpublished.

Rottenborn, S.C. and E.S. Brinkley, editors. 2007. Virginia's Birdlife: An Annotated Checklist, Fourth Edition. Virginia Society of Ornithology, Ervington, VA. 330 pp.

Saltmarsh Habitat and Avian Research Program. 2015. Patch analysis of marsh bird distribution, vector digital data.

The Nature Conservancy. 2015. Shorebird Flats ESI 2015, vector digital data.

TNC, VDGIF, USFWS. 2013. American Oystercatcher (AMOY) 2013 statewide survey pair locations, vector digital data.

- TNC, VDGIF, USFWS. 2016. American oystercatcher (AMOY) winter roost sites, vector digital data.
- USFWS Back Bay National Wildlife Refuge. Back Bay National Wildlife Refuge and False Cape State Park Aerial Surveys 2009-2014, spreadsheet.
- USFWS. 2014. Back Bay NWR peak bird numbers, 2013 and 2014, spreadsheets.
- USFWS DMB. 2014. Atlantic coast seabird survey data, vector digital data.
- USFWS Division of Migratory Birds DMB. 2012. Atlantic coast wintering sea duck survey data, vector digital data.
- USFWS Division of Migratory Birds DMB. 2014. Mid-winter waterfowl surveys 2004-2015, vector digital data and spreadsheet.
- USFWS. 2015. James River National Wildlife Refuge Final Comprehensive Conservation Plan. U.S. Fish & Wildlife Service, Warsaw, VA, 11 pp.
- USFWS. 2008. Mackay Island National Wildlife Refuge Comprehensive Conservation Plan. U.S. Fish & Wildlife Service, Virginia Beach, VA, 214 pp.
- USFWS. 2012. Presquile National Wildlife Refuge Comprehensive Conservation Plan. U.S. Fish & Wildlife Service, Warsaw, VA, 306 pp.
- USFWS DMB. 2011. Southern black scoter survey, vector digital data.
- VDGIF 2015. Midwinter waterfowl surveys 2004-2015, vector digital data and spreadsheet.
- VDGIF. 2015. Virginia Fish and Wildlife Information System (VaFWIS) – Wildlife and Environmental Review Map Service (WERMS). Bald Eagle Concentration Areas and Roosts – BECAR, vector digital data.
- Watts, B.D. and B.J. Paxton (CCB). 2015. Chesapeake Bay Drainage Great Blue Heron and Great Egret Survey 2013, vector digital data.
- Watts, B.D. and B.R. Truitt. 2015. Spring migration of red knots along the Virginia barrier islands. The Journal of Wildlife Management 79: 288-295.
- Watts, B.D. and B.J. Paxton (CCB). 2014. Status and distribution of colonial waterbirds in coastal Virginia: 2013 breeding season, spreadsheet and document.
- Watts, B.D. and M.A. Byrd (CCB). 2015. Virginia Eagle Nests 20151223. College of William & Mary and Virginia Commonwealth University. Williamsburg, VA, vector digital data.
- Watts, B.D. and E. Mojica (CCB). 2015. Virginia peregrine falcon monitoring and management program: Year 2015 report.
- Williams K.A., I.J. Stenhouse, E.E. Connelly, and S.M. Johnson. 2015. Baseline Wildlife Studies Offshore of Maryland. Biodiversity Research Institute. Portland, Maine. Science Communications Series BRI 2015-20, 8 pp
- Wilson. M.D. (CCB). 2016. Black rail VA data 2007-2014, vector digital data.

HERPETOFAUNA

Herpetofauna depicted in this atlas include four species of sea turtles, diamondback terrapin, barking treefrog (VA ST), chicken turtle (VA SE), and snapping turtle.

Nesting and in-water distribution data for diamondback terrapins were provided by VIMS (in-water data set), USGS (nesting data set), and USFWS (pers. comm.). The VIMS data set was based on 2012 and 2013 spring-summer surveys at 165 sites and was conducted to study the relationship between terrestrial-aquatic connectivity and the distribution of an estuarine turtle. The results of the survey included predicted probability of occupancy of diamondback terrapins in the Chesapeake Bay, and the 'high' and 'low' probabilities of occupancy that were determined during the study were adapted for inclusion and

represented as likely terrapin distribution polygons in the ESI. The USGS nesting data set was based on the results of the Patuxent Wildlife Research Center investigation of over 3,000 kilometers of Chesapeake Bay shoreline in 2002 for terrapins, tracks, and active and disturbed nests.

Green (FT/MD ST, VA ST), Kemp's ridley (FE/MD SE, VA SE), leatherback (FE/MD SE, VA SE) and loggerhead (FT/MD ST, VA ST) sea turtles were mapped in this atlas. Many barrier island, and some mainland, outer sand beaches are important nesting habitat for loggerhead sea turtle, and potential (but very rare) nesting habitat for Kemp's ridley and green sea turtles. Nesting typically occurs in Virginia and Maryland from May to August with hatching from June-October (loggerhead sea turtle). Data that represent documented nesting in Virginia from 1970-2015 were provided by VDGIF. The concentration values were based on the range of nest counts observed on a particular stretch of beach over time (e.g., 0-8 nests, <10 nests). Beach buffers were used rather than point locations due to the potential for beaches adjacent to known nest sites to also provide suitable nesting habitats. Green sea turtle and Kemp's ridley sea turtle were documented as having 'false crawls' only and no nest counts. Additional nesting and general distribution and seasonality information were provided during interviews with NPS, USFWS, and the VA Aquarium.

Distributions for the barking treefrog (VA ST) and chicken turtle (VA SE) were mapped using data provided as part of VDGIF's Wildlife Environmental Review Map Service (WERMS) database. Records of both species were pulled from survey dates between the years 1996 and 2013, with 2013 being the most recent surveys for herpetofauna in the database.

Expert contacts for Chesapeake Bay and the outer coasts of Maryland and Virginia herpetofauna*:

Name	Agency	City	Phone	Species
Ruth Boettcher	VDGIF	Machipongo, VA	757/709-0766	Sea turtles
Jack Kumer	NPS	Berlin, MD	410/629-6070	Assateague Island NS
Lauren Billodeaux Mowbray	USFWS	Virginia Beach, VA	757/301-7329	Back Bay NWR
Susan Barco	VA Aquarium	Virginia Beach, VA	757/385-6476	Sea turtles
Robert Isdell III	VIMS	Gloucester Pt., VA	804/684-7179	Diamond-back terrapins
Jay Kapalczynski	VDGIF	Henrico, VA	804/367-6796	WERMS Database

*Note: this list is not meant to represent all herpetofauna experts for the region.

Major Data Sources Used: Herpetofauna

Isdell, R.E., R.M. Chambers, D.M. Bilkovic, and M. Leu. 2015. Effects of terrestrial-aquatic connectivity on an estuarine turtle, *Diversity and Distributions* (2015) 1-11.

National Marine Fisheries Service. 2015. Virginia and Maryland Sea Turtle Conservation Plan. National Marine Fisheries Service, Virginia Beach, 58 pp.

USFWS. 2012. Presquile National Wildlife Refuge. Comprehensive Conservation Plan, USFWS, Warsaw, VA, 306 pp.

U.S. Geological Survey. 2002. Terrapin survey 2002, vector digital data.

VDGIF. 2015. Virginia sea turtle nest data, 1970-2015, tabular data.

VDGIF. 2015. Wildlife environmental review map service, species observation database. Vector digital data.

Virginia Institute of Marine Science. 2015. Terp occupancy, vector digital data.

FISH

Fish species depicted in this atlas include select marine, estuarine, and freshwater species. Species of conservation interest, commercial or recreational importance, or ecological importance are emphasized. Fish polygons were created based on fisheries-independent survey information, digital data, and expert opinion provided primarily by resource experts at MD DNR, VDGIF, VIMS, VMR, ASMFC, USFWS, and NOAA.

Atlantic and Shortnose sturgeon - Populations of shortnose sturgeon (FE, VA/MD SE) and Atlantic sturgeon (FE, VA SE) are found in Chesapeake Bay and adult and sub-adult Atlantic sturgeon distributions also encompass the coastal waters of Virginia and Maryland. Data on shortnose sturgeon populations in Chesapeake Bay are limited to incidental catches during a tagging program for Atlantic sturgeon by USFWS, and to a few recent studies of movements in the Potomac river. Generally, shortnose sturgeon are concentrated in the upper portions of the Chesapeake Bay, roughly from the mouth of the Chester River and north, but they are not uncommon to the mouth of the Potomac, and have been found to stray as far south as the James River in Virginia. No spawning of shortnose sturgeon has been documented in Chesapeake Bay, but there is some evidence of potential for spawning in the Potomac River and possibly the Susquehanna River. Atlantic sturgeon inhabit all areas in the Chesapeake Bay at various life stages. For the purposes of the ESI, life stages of Atlantic sturgeon were defined using definitions published by the Atlantic States Marine Fisheries Commission (ASMFC). In the ESI data tables, larvae refer to fish less than 4 weeks old, juveniles refer to young-of-year, and adults include both sub-adults (age 2 to 15) and adults. Eggs, larvae, and juveniles are restricted to known or potential spawning rivers, including the James, Nanticoke, Pocomoke, Rappahannock Rivers, and the York River system, which includes the Mattaponi and Pamunkey Rivers.

Special concentration information was also included for Atlantic and shortnose sturgeon spawning and migration areas. For Atlantic sturgeon, the concentration field was used to differentiate between spring and fall spawning migration runs in the James River, where adult sturgeon have been documented making spawning forays during both spring and fall months. However, spawning itself has only been conclusively documented during the fall. Areas within the Mattaponi, Pocomoke, Nanticoke, Rappahannock, and James Rivers (spring run) were marked as 'Potential Spawning' due to the fact that while milting and migrating adult fish have been documented, fertilized eggs have not been documented in these systems. Fall spawning has only been documented for the Pamunkey and James Rivers. Finally, two areas were marked as concentration areas for Atlantic sturgeon, one in the Nanticoke River, and one in the James River. Both areas are known to attract high numbers of adults and sub-adults and are believed to be staging areas for spawning runs upstream.

Anadromous Fish – Virginia - A number of other anadromous species' spawning runs and nursery areas were mapped based on multiple data sources. Alewife, blueback herring, American shad, and hickory shad distributions were mapped based on expert knowledge provided by Eric Brittle and Alan Weaver (VDGIF), The Nature Conservancy's Chesapeake Fish Passage Prioritization Project data, and the Habitat Requirements for Chesapeake Bay Living Resources. White and yellow perch spawning and nursery areas were initially mapped using the Habitat Requirements for Chesapeake Bay Living Resources and were modified based on comments from VDGIF, USFWS, and VIMS biologists. Striped bass spawning and nursery areas were mapped primarily based on expert knowledge from VIMS and VDGIF biologists.

Anadromous Fish – Maryland - Alewife, blueback herring, American shad, and hickory shad distributions were mapped based on expert knowledge provided by Jim Thompson and Margaret McGinty (VDGIF), The Nature Conservancy's Chesapeake Fish Passage Prioritization Project data, and MD DNR data on river herring spawning and nursery habitat. White and yellow perch spawning and nursery areas were initially mapped using the Habitat Requirements for Chesapeake Bay Living Resources and were modified based on comments from MD DNR and USFWS biologists. Striped bass spawning and nursery areas were mapped using MD DNR data on spawning reaches and rivers, and were expanded based on expert knowledge of MD DNR staff. Anadromous fish presence and spawning runs in the Coastal Bays of Maryland were mapped based on expert knowledge provided by Steve Doctor and Angel Willey.

It should be noted that alewife, blueback herring, and shad species spawn outside of the areas identified as 'spawning areas' in the ESI. This is especially true of the two herring species which utilize freshwater streams and lakes to a large degree for spawning. Also, note that eel life histories do not match the standard ESI terminology, so for the purpose of this ESI, 'juveniles' correspond to immigrating glass eels and 'adults' refer to resident yellow eels and emigrating silver eels. Lastly, striped bass juveniles and sub-adults can be found year-round in Chesapeake Bay. Mature females typically join the coastal migration at year three or four, and while the majority of males do as well, some are known to become permanent residents in the Bay. Areas in the ESI identified as 'spawning' or 'nursery' areas, specifically large areas in the Potomac, York, James, and Rappahannock Rivers, represent the most important early life stage areas; however, striped bass likely utilize other smaller tributaries and water bodies for spawning as well.

Finally, there was special consideration given to a number of rivers on the western shore of Chesapeake Bay in Maryland. These are rivers where anadromous fish still make migratory runs and attempt to spawn. However, the deteriorated condition of these rivers prevents the successful maturation of eggs and larvae. Thus, while these rivers still attract significant numbers of spawning adult fish, they are functionally sterile. Rivers on the western shore that were identified by MD DNR staff as described above include: Bush River, Severn River, South River, and Magothy River. Anadromous fish species that attempt to utilize these rivers for spawning have been recorded in the ESI attribute table 'Concentration' field with the designation 'Historical Spawning'.

Freshwater fish – Maryland - Freshwater finfish distributions were initially mapped to the river systems based on sampling data (2000-2014) from the Maryland Biological Stream Survey (MBSS). This survey is conducted on a yearly basis during the summer months in streams throughout Maryland using a stratified-random sampling method to determine the yearly sampling locations, except for a limited series of sentinel or reference sites that are sampled every year. Although the MBSS does not sample in the main stems of Maryland's rivers, as it is specifically a survey conducted in smaller tributaries that feed the major rivers, MBSS survey data were used to assign initial fish distributions to the fresh water reaches of Maryland's river systems (Back, Bohemia, Bush, Chester, Choptank, Elk, Gunpowder, Middle, Nanticoke, North East, Patuxent, Potomac, Sassafras, and Susquehanna) included in this ESI. MBSS data were utilized due to a lack of sampling data available for the fresh water portions of these rivers. Fish species that were captured in multiple years (>3) and in multiple sampling locations (>5) within the watersheds were included in the initial freshwater fish distribution lists and assigned to river polygons proximal to the sites where they were captured. In the Potomac River, freshwater fish species captured in the VIMS Juvenile Fish and Blue Crab Trawl Survey, which encompasses sampling effort up to the approximate fresh/salt water mixing zone in the Potomac (and James, Rappahannock and York Rivers in VA), but not captured by the MBSS, were also included in the initial distributions to supplement the MBSS data (see also methods below under marine and estuarine fish). Seasonalities for spawning and the presence of various life stages were assigned using information contained in the Field Guide to Fishes of the Chesapeake Bay, when available, due to a lack of year round sampling data.

Through consultations and in-person meetings with resource managers/scientists at MD DNR who were responsible for the sampling programs and management of Maryland's freshwater fisheries, fish species included in the initial distribution assessment (as described above) were systematically reviewed for their spatial distributions and life history characteristics (seasonalities) and either accepted, modified, or removed based on the comprehensive knowledge and expert opinions of the MD DNR reviewers.

Freshwater fish – Virginia - In Virginia, freshwater species were mapped in the river systems, bays, and sounds using data from the VIMS Juvenile Fish and Blue Crab Trawl survey, VDGIF Wildlife Environmental Review Map Service (WERMS) Species Observations Database, USFWS Back Bay National Wildlife Refuge fishery management plans and survey data, and expert opinion from VIMS scientists and VGDIF staff.

Freshwater species distributions were mapped in the main rivers in Virginia (James, Rappahannock and York Rivers) primarily using data from the VIMS Juvenile Fish and Blue Crab Trawl dataset, which was summarized by VIMS for the sampling years 2005-2014, to give total number of fish captured by species per sampling station. Utilizing this summary, the frequency of each species caught during each month

was calculated for the designated sampling regions, which were based on polygonal regions designated by VIMS for the trawl survey. Initial distributions were based on species that were captured in the trawl survey at a 10% frequency or higher (based on number of trawls) in a given month for a given sampling region. Species that met the 10% threshold were marked as present for that month and region.

Some rivers and bays in Virginia had only limited or no sampling data, including Currituck Sound, Back Bay, and North Landing River. Initial fish distributions for these areas were developed using available publications, including management plans, and limited survey data (when available). Fresh water regions lacking all data were initially populated for species using data from adjacent sampling regions (i.e., VIMS Juvenile Fish and Blue Crab Trawl) and the WERMS Species Observations Database. Additional distributions and life history seasonalities, such as larvae, eggs, and spawning months, were typically gathered from the Field Guide to Fishes of the Chesapeake Bay (Murdy and Musick, 2013) and expert knowledge provided by VIMS scientists or VDGIF staff.

Through a series of consultations and in-person meetings with resource managers/scientists at VIMS and VDGIF, who are responsible for the sampling programs and/or management of Virginia's freshwater fisheries, fish species included in the initial distribution assessment (as described above) were systematically reviewed for their spatial distributions and life histories (seasonalities) and either accepted, modified or removed based on the knowledge and expert opinions of the reviewers.

Additional fresh water fish species known to occur in the river systems of Maryland and Virginia, which were not contained within the initial data request and assessment, were also included in this ESI based on the expert opinion of the reviewers from the respective states.

Nearshore and Estuarine Fish Species – Nearshore marine and estuarine distributions of fish species were mapped primarily using fishery-independent data from the Northeast Area Monitoring and Assessment Program (NEAMAP), the Chesapeake Bay Multispecies Monitoring and Assessment Program (ChesMMAP), MD DNR Coastal Bays Fisheries Investigation, and VIMS Juvenile Fish and Blue Crab Trawl Survey. These data sets were supplemented with information from Funderburk et al. (1991), Murdy and Musick (2013), and then systematically reviewed by resource managers and scientists from MD DNR, VDGIF, USFWS and VIMS.

Data from ChesMMAP and the VIMS Juvenile Fish and Blue Crab Trawl Survey were used to map fish species distributions within Chesapeake Bay and its tributaries. ChesMMAP data, which was initially requested for 40 commercially and/or recreationally important species, is collected during the months of March, May, July, September and November. The VIMS Juvenile Fish and Blue Crab Trawl Survey is conducted in every month throughout the year. The VIMS Trawl survey was also used to map species into the three main Virginia tributaries of the Chesapeake Bay (James River, Rappahannock River and the York River) and the Potomac River in Maryland, up to, and including, the fresh water boundary regions (see fresh water fish species above). Both datasets were used as a reference for fish distributions in Maryland's river systems, where there was a lack of sampling data available. Each requested dataset was summarized by the data provider for the sampling years 2005-2014 to give total number of occurrences per sampling station by species. From these summaries, the frequency of capture of each species caught during each month for each sampling region was calculated (based on how many times a species was captured in a trawl divided by the total number of trawls conducted in the sampling region). The standardized polygonal sampling regions provided by the respective surveys for each of these datasets were used to create the geographic distributions of species. Species that were found to occur with a 10% frequency or higher in a given month for a given sampling region were marked as present for that month and sampling region. Due to the limited sampling times in the ChesMMAP data, the final monthly presence and absence data was determined through in-person consultations with VIMS and MD DNR scientists.

For marine species in the Atlantic Ocean from the coast out to 12 nm for both Maryland and Virginia, fish species distributions and seasonalities were initially created using data from the NEAMAP trawl survey, which began sampling in the fall of 2007. NEAMAP sampling takes place in the coastal waters of Maryland and Virginia in the spring (April and occasionally May) and fall (October) each year. Data used for the ESI included trawls from the years 2007-2014, and the frequencies of species caught were used to

determine monthly presence/absence for the sampling months. Presence within a sampling month was determined by the same 10% frequency cut-off outlined for the ChesMMAP and VIMS Trawl data above. Due to the limited sampling times in the NEAMAP data the final monthly presence and absence data were determined through face-to-face consultations with VIMS and MD DNR scientists.

Maryland's Coastal Bays Fisheries Investigation (CBFI) Survey data, summarized by MD DNR, were used to initially map species distributions in the coastal bays of Maryland. Due to a lack of survey data in Virginia's coastal bays, CBFI data were also used as a reference for initial species distributions in the coastal bays of Virginia. Species distributions for the Back Bay region were mapped based on reports and data supplied by USFWS and expert opinion from VDGIF scientists.

Additional species distributions and life history traits, such as larvae, eggs, and spawning months, were gathered from the Field Guide to Fishes of the Chesapeake Bay (Murdy and Musick, 2013), ELMR Program data (Stone et al., 1994) and expert opinion from MD DNR and VIMS. All marine and estuarine species distributions and presence/absence data included in this ESI were systematically reviewed and approved through consultations with MD DNR, VDGIF, USFWS and VIMS scientists.

Sharks - The southern region of Chesapeake Bay is an important nursery area for several shark species. Nursery areas and general distributions of sharks were mapped based on expert knowledge from scientists at MD DNR and VIMS, who conduct both adult longline and juvenile shark (COASTSPAN) surveys within the area covered by this atlas. Note that the life-history stages for fish in the ESI data structure do not match the correct terminology for sharks. As a result, areas with months for shark 'larvae' represent months for which that polygon serves as habitat or a nursery area for young-of-year sharks (i.e., primary nursery areas).

Estuarine and marine fish species generally spawn where and when environmental conditions are suitable and spatially defined spawning areas have not been determined through surveys for the majority of the marine, estuarine, and fresh water species included in this atlas. Spawning information for these species' life history is contained in the 'general distribution' polygons to capture seasons in which spawning, eggs, and larvae are likely to be present. Exceptions include anadromous species which have known spawning locations.

This atlas does not contain all of the fish species that are present in the Chesapeake Bay, in the coastal bays of Maryland and Virginia, or in the Atlantic Ocean. Some species, like killifish spp. (in Chesapeake Bay), butterfish, mummichog, and needlefish, were not included due to their ubiquitous nature in the bay and/or marine environment; while others species were not included due to lack of data or on the expert opinion of the scientists and data providers.

Expert contacts for Virginia fish*:

Name	Agency	City	Phone	Species
Eric Brittle	VDGIF	Chesapeake, VA	757/334-5075	Anadromous and Freshwater Fish
Alan Weaver	VDGIF	Richmond, SC	804/367-6795	Anadromous and Freshwater Fish
Bob Greenlee	VDGIF	Charles City, VA	804/829-6715	Atlantic and shortnose sturgeon
Katie May Laumann	VMRC	Newport News, VA	757)-247-2263	Atlantic sturgeon
Albert Spells	USFWS	Charles City, VA	804/829-5627	Atlantic and shortnose sturgeon
Jason Kahn	NOAA	Silver Springs, MD	301/427-8464	Atlantic sturgeon
Troy Tuckey	VIMS	Gloucester Point, VA	804/684-7328	Estuarine fish and striped bass
Mary C. Fabrizio	VIMS	Gloucester Point, VA	804/684-7308	Estuarine fish

Name	Agency	City	Phone	Species
Chris Bonzek	VIMS	Gloucester Point, VA	804/684-7291	Marine finfish
Debra Gauthier	VIMS	Gloucester Point, VA	804/684-7891	Marine finfish

*Note: this list is not meant to represent all fish experts for the region.

Expert contacts for Maryland fish*:

Name	Agency	City	Phone	Species
Steve Doctor	MD DNR	Ocean City, MD	410/213-1531	Marine and estuarine fish
Angel Willey	MD DNR	Matapeake, MD	410/643-6801 x 2101	Marine and estuarine fish
Jim Thompson	MD DNR	Annapolis, MD	410/260-8269	Anadromous fish
Beth Versak	MD DNR	Annapolis, MD	410/260-8304	Striped bass
Paul Piavis	MD DNR	Annapolis, MD	410/643-6801	Estuarine and anadromous fish
Margaret McGinty	MD DNR	Annapolis, MD	410/260-8297	Estuarine and anadromous fish
Chuck Stence	MD DNR	Stevensville, MD	410/643-6788	Atlantic sturgeon
Steve Minkinnen	USFWS	Annapolis, MD	410/573-4506	Atlantic and shortnose sturgeon
Mike Mangold	USFWS	Annapolis, MD	410/573-4509	Atlantic and shortnose sturgeon
Matt Ashton	MD DNR	Annapolis, MD	410/260-8604	Freshwater fish
Scott Stranko	MD DNR	Annapolis, MD	410/260-8603	Freshwater fish

*Note: this list is not meant to represent all fish experts for the region.

Major Data Sources Used: Fish

Funderburk, S.L., S.J. Jordan, J.A. Mihursky, and D. Riley, editors. 1991. Habitat requirements for Chesapeake Bay living resources (Revised edition). Chesapeake Bay Program, Annapolis, MD.

MD DNR. 2014. Coastal Bays Fisheries Investigation Trawl and Beach Seine Survey data, 2005-2014, spreadsheet.

MD DNR. 2014. Maryland Biological Stream Survey (MBSS) data, 2000-2014, spreadsheet.

MD DNR. 2015. Maryland river herring spawning and nursery habitats, vector digital data.

MD DNR. 2012. Maryland striped bass spawning areas, vector digital data.

Murdy, E.O. and J.A. Musick. 2013. Field guide to fishes of the Chesapeake Bay. The John Hopkins University Press, Baltimore, MD, 345 pp.

Stone, S.L., T.A. Lowery, J.D. Field, C.D. Williams, D.M. Nelson, S.H. Jury, M.E. Monaco and L. Andreasen. 1994. Distribution and abundance of fishes and invertebrates in Mid-Atlantic estuaries. ELMR Rep. No. 12. NOAA/NOS Strategic Environmental Assessments Division, Silver Spring, MD. 280 p.

The Nature Conservancy. 2015. Chesapeake Bay Fish Passage Prioritization Project data, vector digital data.

- USFWS. 2011. Maryland sturgeon reward program captures – Atlantic and shortnose sturgeon, spreadsheet
- VDGIF. 2015. Wildlife Environmental Review Map Service: Species Observation Database (SppObs), vector digital data.
- VIMS. 2014. VIMS Multispecies Research Group, Chesapeake Bay Multispecies Monitoring and Assessment Program (ChesMMAP) data, 2005-2014, spreadsheet.
- VIMS. 2014. VIMS Multispecies Research Group, Northeast Area Monitoring and Assessment Program (NEAMAP) data, 2005-2014, spreadsheet.
- VIMS. 2014. VIMS Recruitment Program Juvenile Fish and Blue Crab Trawl Survey data, 2005-2014, spreadsheet.

INVERTEBRATES

Invertebrates depicted in this atlas include selected marine and estuarine species of commercial, ecological, and/or conservation value. Blue crabs, clams, oysters, and shrimp support valuable commercial and/or recreational shellfisheries with Chesapeake Bay and the coastal waters of Virginia and Maryland.

Marine and Estuarine Invertebrate Species - Sources used to depict distribution and seasonality of invertebrate species include: personal interviews with resource experts from MD DNR, VDGIF, VA MRC and VIMS; fishery-independent sampling data provided by MD DNR, NEAMAP, ChesMMAP and VIMS; species observation datasets for Virginia; reports and hardcopy maps for soft clam and razor clam distributions in Maryland; and vector digital data covering oyster reef locations provided by VIMS and MD DNR and shellfish lease areas provided by VA MRC.

Blue crabs - Blue crab were initially mapped in Chesapeake Bay and the Atlantic Ocean using fishery-independent data for the time period 2005–2014 from NEAMAP, ChesMMAP, and VIMS Juvenile Fish and Blue Crab Trawl survey. Spatial data from Habitat Requirements for Chesapeake Bay Living Resources (Funderburk et al., 1991) was used to depict summer high density areas. Winter high density areas were mapped using independent fishery data from the Blue Crab Winter Dredge survey (2009–2010) provided by VIMS. Spatial distributions in the coastal bays of Maryland and Virginia were mapped using expert knowledge from biologists at MD DNR and VIMS. Blue crab presence/absence data were then systematically reviewed and modified through consultations with MD DNR and VIMS scientists to the final distributions depicted on this ESI.

Chesapeake Bay Shellfish - Oysters and other commercially and/or recreationally important shellfish species, including hard clams, soft clams and stout razor clams, were mapped in Chesapeake Bay using a combination of independent fishery survey data, shellfish lease areas, reports and expert knowledge from MD DNR, VA MRC and VIMS. In Maryland, shellfish distributions were mapped using a combination of MD DNR Oyster Dredge Survey data (2005–2014), Maryland’s Historical Oyster Bottom vector digital data, the Assessment of Chesapeake Bay Commercial Softshell Clams (Homer et al., 2011), Habitat Requirements for Chesapeake Bay Living Resources (Funderburk et al., 1991), and expert opinion from MD DNR biologists. In Virginia, shellfish spatial distributions were mapped using VA MRC 2015 active shellfish lease data, Habitat Requirements for Chesapeake Bay Living Resources (Funderburk et al., 1991), and expert opinion from VA MRC and VIMS scientists.

Coastal Bays Shellfish - Oysters, stout razor clam and softshell clam distributions in the coastal bays of Maryland were mapped based on expert opinion of MD DNR. Hard clam distributions were mapped using data from MD DNR Coastal Hard Clam survey from 2004–2014. In Virginia’s coastal bays, oyster general distributions were based on shellfish lease digital data and expert opinion provided by VA MRC, and oyster reefs were mapped using digital data provided by VIMS (Ross et al., 2009), which had a resolution of approximately 1 m². The oyster reef polygons in this dataset were buffered by 25 m and

then merged together to conform to the minimum resolution requirements of the ESI maps and to take into account the growth and shifting nature of oyster reefs. Oyster reefs mapped in the ESI are not the only reefs that could occur in coastal areas, but represent current and potential oyster habitat.

Other Invertebrates - Distributions of other marine and estuarine invertebrates, including brown shrimp, white shrimp, mantis shrimp, and squid species were mapped in Chesapeake Bay and the Atlantic Ocean using data from NEAMAP, ChesMMAP, and VIMS Juvenile Fish and Blue Crab Trawl surveys from 2005–2014. Presence and absence of species was calculated using the NEAMAP, CHesMMAP and VIMS Juvenile Fish and Blue Crab survey data sets and followed the same methods outlined under the fish section (above). Additional invertebrate species with spatial distributions in the coastal bays of Maryland and Virginia and in marine waters (out to 12 nm), including life history traits, were based on expert knowledge provided by MD DNR, VA DGIF and VIMS. All marine and estuarine invertebrate spatial distributions and presence/absence data depicted in this ESI were systematically reviewed and finalized through consultations with MD DNR, VA MRC and VIMS scientists prior to inclusion.

Horseshoe crabs - Horseshoe crab spawning beaches were mapped from three different sources. In the Coastal Bays of Maryland, spawning beaches were identified using expert knowledge provided by Steve Doctor. Tom's Cove in the Chincoteague National Wildlife Refuge holds arguably the most important spawning habitat within the Chesapeake Bay and the Outer Coasts of Maryland and Virginia ESI. Spawning beaches in Tom's Cove were mapped from the Tom's Cove, Virginia Horseshoe Crab Tag Recovery Study Interim Report. Lastly, a number of small spawning sites within Chesapeake Bay itself were mapped using public sightings data provided by MD DNR. It should be noted that horseshoe crabs utilize whatever appropriate habitat is available inside Chesapeake Bay, and that these areas can shift over time. No comprehensive surveys for spawning beaches have been conducted, so the public sightings data are unlikely to capture all the ecologically significant areas.

Freshwater mussels - Freshwater mussels in Virginia and Maryland were mapped using VDGIF's Freshwater Mussel database, MD DNR's Ecologically Significant Areas database and the Maryland Biological Stream Survey data, respectively. Mussels mapped in Virginia represent rare species that are of conservation concern and are of significant ecological importance. Species observations mapped from the VDGIF database were restricted to the dates 1995–2013, and only include records that fell within 1000 m of ESI water bodies that are continuous to the ocean. Species mapped in Maryland have similar importance, but also include two endangered species: the dwarf wedge mussel (FE, MD SE) and the triangle floater (MD SE). Species mapped from MD DNR's Ecologically Significant Areas database are restricted based on date of last survey and include the years 1996 to 2016.

Insects - A number of insects of conservation concern were mapped in Maryland using MD DNR's Ecologically Significant Areas database. Records were selected based upon date of last survey and include data from 1996 to 2016. Insect species mapped from this database include three beetles: a darkling beetle, *Schoenicus puberulus* (MD SE); a helops darkling beetle, *Helops cistelooides* (MD SE); and the white tiger beetle, *Habroscelimorpha dorsalis media* (formerly *Cicindela dorsalis media*) (MD SE); and two species of butterfly: Palamedes swallowtail, *Papilio palamedes* (MD SE), and Selys' sunfly, *Helocordulia selysi* (MD ST). USFWS provided shapefiles depicting occurrences of FT/SE insects; common names were not included in the data tables for these species.

Tiger beetles - Northeastern beach tiger beetle (*Cicindela dorsalis dorsalis*, FT, MD SE, VA ST) occurrences in Virginia were mapped using a hard copy map from USFWS. On the eastern shore of Chesapeake Bay in Virginia, two sites account for more than 50% of the total population along that shore. These sites, Parker's Marsh and Savage Neck, have been marked as 'Concentration Areas' in the ESI database. The western shore harbors a smaller total population of northeastern beach tiger beetles, mostly due to recent hurricane events in 2003 and 2006, that seriously altered beach habitat and resulted in a large drop in population size. As of the last survey in 2008, approximately 10,000 tiger beetles were found on the western shore, as opposed to more than 38,000 beetles found on the eastern shore in 2005. Northeastern beach tiger beetles were also mapped in various parts of the study area according to expert knowledge.

Assateague National Seashore and USFWS National Wildlife Refuges - Resource experts provided distribution information for little white tiger beetle (*Elliptoptera lepida*, MD SE), white tiger beetle (see above),

northeastern beach tiger beetle (see above), and monarch butterfly (*Danaus plexippus*) on federal lands for which they had information.

The species depicted in this atlas do not represent all of the invertebrate species present in this area of interest, including species like ribbed mussels which are ubiquitous to salt marsh habitats. Other invertebrate species, like blue mussels and some razor clam species which can be found in the coastal bays of Maryland and Virginia as juveniles, are also ecologically important as a food source for migrating birds and other invertebrate/fish species, but because of their ephemeral nature and lack of stable adult populations, were not included in this atlas.

Expert contacts for Chesapeake Bay and the outer coasts of Maryland and Virginia invertebrates*:

Name	Agency	City	Phone	Species/ Program
Steve Doctor	MD DNR	Ocean City, MD	410/213-1531	Horseshoe crabs and marine invertebrates
Marek Topolski	MD DNR	Annapolis, MD	410/260-8263	Chesapeake Bay horseshoe crabs
Andy Moser	USFWS	Annapolis, MD	410/573-4537	Tiger Beetles
Brian Watson	VDGIF	Forest, VA	434/525-7522	Freshwater mussels
Matt Ashton	MD DNR	Annapolis, MD	410/260-8604	Freshwater mussels
Lynn Davidson	MD DNR	Annapolis, MD	410/260-8563	Digital data provider
Mark Luckenback	VIMS	Gloucester Point, VA	804/684-7108	Marine and estuarine shellfish
Jim Wesson	VA MRC	Newport News, VA	757/247-2121	Marine and estuarine shellfish
Roger Mann	VA MRC	Gloucester Point, VA	804/ 684-7360	Marine and estuarine shellfish
Angel Willey	MD DNR	Matapeake, MD	410/643-6801 X 2101	Marine invertebrates
Mitch Tarnowski	MD DNR	Annapolis, MD	410/260-8258	Marine and estuarine shellfish
Ben Stagg	VA MRC	Newport News, VA	757-247-2225	Oysters
Leslie Gerlich	USFWS	Annapolis, MD	410/573-4587	Federal T/E species

*Note: this list is not meant to represent all invertebrate experts for the region.

Major Data Sources Used: Invertebrates

Funderburk, S.L., S.J. Jordan, J.A. Mihursky, and D. Riley, editors. 1991. Habitat requirements for Chesapeake Bay living resources (Revised edition). Chesapeake Bay Program, Annapolis, MD.

Hallerman, E. 2009. Tom's Cove, Virginia horseshoe crab tag recovery study: interim report. A report to permitting authorities: Assateague Island National Seashore, Chincoteague National Wildlife Refuge, Virginia Marine Resources Commission. Virginia Polytechnic Institute and State University, Blacksburg, VA.

Homer, M.L., C.F. Dungan, and M.L. Tarnowski. 2011. Assessment of Chesapeake Bay Commercial softshell clams, *Mya arenaria* and *Tagelus plebeius*, with emphasis on abundance and disease status. Maryland Department of Natural Resources, Award NA07NMF4570326.

MD DNR. 2016. Ecologically significant areas of Maryland, vector digital data.

- MD DNR. 2014. Maryland Biological Stream Survey (MBSS) data, *Elliptio complanata* occurrences, 1995-2014, spreadsheet.
- MD DNR. 2014. Maryland Department of Natural Resources Coastal Hard Clam Survey data, 2004-2014, spreadsheet.
- MD DNR. 2014. Maryland Department of Natural Resources Oyster Dredge Survey data, 2005-2014, spreadsheet.
- MD DNR. 1997. Maryland Department of Natural Resources Historical Oyster Bottom. Vector digital data.
- MD DNR. 2015. Maryland horseshoe crab spawning public siting data, 2007-2015, spreadsheet.
- Ross, P.G. and M.W. Luckenbach. 2009. Population assessment of Eastern Oysters (*Crassostrea virginica*) in the Seaside Coastal Bays, vector digital data.
- USFWS. 2011. Northeastern beach tiger beetle occupied habitat, Chesapeake Bay. Gloucester, VA.
- VDGIF. 2013. VDGIF Freshwater Mussel Database, 1995-2013, vector digital data.
- VA MRC. 2015. Virginia Shellfish Lease Areas. Vector digital data.
- VIMS. 2010. Blue Crab Winter Dredge Survey data, 2009-2010. Map images and spatial analyses.
- VIMS. 2014. VIMS Multispecies Research Group, Chesapeake Bay Multispecies Monitoring and Assessment Program (ChesMMAP) data, 2005-2014, spreadsheet.
- VIMS. 2014. VIMS Multispecies Research Group, Northeast Area Monitoring and Assessment Program (NEAMAP) data, 2005-2014. Spreadsheet. Tarnowski, Mitch (MD DNR). Shellfish distributions and life histories in Chesapeake Bay and the coastal bays of Maryland. Personal communication.
- VIMS. 2014. VIMS Recruitment Program Juvenile Fish and Blue Crab Trawl Survey data, 2005-2014, spreadsheet.

BENTHIC HABITATS

Benthic habitats mapped in the ESI atlas include submerged aquatic vegetation (SAV) in the Chesapeake Bay and Back Bay. SAV data for the Chesapeake Bay from 2010-2014 aerial surveys were provided by VIMS. VIMS categorized SAV density for each year for all survey locations using a scale from 1, very sparse (<10% cover), to 4, dense (70-100% cover). For the ESI, VIMS provided five years (2010-2014) of SAV data to account for interannual variation in bed size and density. The five years of data were layered and the resulting polygons were categorized as either "high" or "low" density based on whether a majority of the years (three or more years) had high density categories (3 or 4) or low density categories (1 or 2). Areas with a density category of 0 for all five years were not mapped. Areas with one or two 0 density years were assigned to the ESI "high" or "low" categories based on the density categories of the majority of remaining years. All areas with at least one year of SAV presence, regardless of density value, that was not already covered by the "high" and "low" polygons were mapped with a concentration of "ephemeral". "High", "low", and "ephemeral" polygons were dissolved to make contiguous polygons. SAV in Back Bay was mapped where present using the SAV map of North Carolina and Southeastern Virginia originated by the North Carolina Division of Marine Fisheries and provided by Back Bay NWR. Submerged aquatic vegetation mapped in the ESI is comprised of a variety of species, and the species assemblages are often determined by salinity. The following table lists species assemblages known to occur in the Chesapeake Bay and the months when they are present. Density of all SAV communities in Chesapeake Bay peak in the summer months, and some species are present year-round (Moore et al., 2000). The SAV survey data provided were not species-specific, so all SAV mapped in the Chesapeake Bay were mapped as present year-round. Freshwater SAV communities defoliate and are not present in winter (Moore et al., 2000), so SAV in Back Bay, an oligohaline bay, was mapped only during months present for freshwater SAV.

SAV species assemblages and months present:

SAV community type	Species associations	Months present
Zostera Community	<i>Zostera marina</i> * <i>Ruppia maritima</i>	January - December
Ruppia Community	<i>Ruppia maritima</i> * <i>Potamogeton perfoliatus</i> <i>Potamogeton pectinatus</i> <i>Zannichellia palustris</i>	January - December
Potamogeton Community	<i>Potamogeton pectinatus</i> * <i>Potamogeton perfoliatus</i> * <i>Potamogeton crispus</i> <i>Elodea canadensis</i>	May - November
Freshwater Mixed Community	<i>Myriophyllum spicatum</i> * <i>Hydrilla verticillata</i> * <i>Vallisneria americana</i> * <i>Ceratophyllum demersum</i> <i>Heteranthera dubia</i> <i>Elodea canadensis</i> <i>Najas</i> spp. <i>Potamogeton crispus</i> <i>Potamogeton pusillus</i>	May - November

*Dominant species

Expert contacts for Chesapeake Bay and the outer coasts of Maryland and Virginia benthic habitats*:

Name	Agency	City	Phone	Species/ Program
Bob Orth	VIMS	Gloucester Point, VA	804/684-7392	SAV
David Wilcox	VIMS	Gloucester Point, VA	804/684-7088	SAV
Lauren Billodeaux Mowbray	USFWS	Virginia Beach, VA	757/301-7329 x3108	Back Bay NWR SAV
Geralyn Mireles	USFWS	Virginia Beach, VA	757/301-7329 x153	Back Bay NWR SAV

*Note: this list is not meant to represent all benthic habitat experts for the region.

Major Data Sources Used: Benthic Habitats

Moore, K.A., D.J. Wilcox, and R.J. Orth. 2000. Analysis of the abundance of submersed aquatic vegetation communities in the Chesapeake Bay. *Estuaries* 23: 115-127.

North Carolina Division of Marine Fisheries, Habitat Protection Section, Albemarle-Pamlico Sound National Estuary Program Submerged Aquatic Vegetation (SAV) Mapping Partnership. 2008. SAV Map of North Carolina and Southeastern Virginia 2007-2008, vector digital data.

VIMS. 2015. SAV distribution, 2010-2014, vector digital data.

HABITATS

Plant Species and Sensitive Ecological Communities – The MD DNR provided the majority of data for rare, threatened, and endangered plant species from their Ecologically Significant Areas (ESAs) of Maryland database. Data from the ESAs came in the form of generalized polygons within which multiple threatened or endangered species can often be found. Species include two federally listed species: seabeach amaranth (FT, VA ST, MD SE) and sensitive joint-vetch (FT, VA ST, MD SE) as well as numerous Maryland state listed species. To simplify display of these data, species names were generalized to reflect the Maryland state status with one of two designations, either ‘threatened plant’ or ‘endangered plant’. When more than one threatened or endangered species is found in the same polygon, this is recorded in the concentration field of the attribute table. For example, if two distinct endangered plants occur in the same polygon, this would result in a single record attributed with ‘2 species present’ in the concentration field. If there is no attribute in the concentration field, than only a single species is present.

State threatened plant genera mapped in Maryland:

<i>Antennaria</i>	<i>Linum</i>	<i>Sclerolepis</i>
<i>Carex</i>	<i>Lygodium</i>	<i>Scutellaria</i>
<i>Chamaedaphne</i>	<i>Lysimachia</i>	<i>Smilax</i>
<i>Chelone</i>	<i>Paspalum</i>	<i>Solidago</i>
<i>Diplazium</i>	<i>Phemeranthus</i>	<i>Sphenopholis</i>
<i>Eleocharis</i>	<i>Polygala</i>	<i>Stenanthium</i>
<i>Eriocaulon</i>	<i>Rhynchospora</i>	<i>Thelypteris</i>
<i>Eupatorium</i>	<i>Sagittaria</i>	
<i>Hypericum</i>	<i>Sanguisorba</i>	

State endangered plant genera mapped in Maryland:

<i>Aeschynomene</i>	<i>Equisetum</i>	<i>Leersia</i>	<i>Parnassia</i>	<i>Sesuvium</i>
<i>Agalinis</i>	<i>Eriocaulon</i>	<i>Lemna</i>	<i>Paspalum</i>	<i>Sida</i>
<i>Amaranthus</i>	<i>Eurybia</i>	<i>Liatris</i>	<i>Pedicularis</i>	<i>Sorghastrum</i>
<i>Aristida</i>	<i>Gaylussacia</i>	<i>Limnobium</i>	<i>Persea</i>	<i>Sporobolus</i>
<i>Asclepias</i>	<i>Glyceria</i>	<i>Limosella</i>	<i>Pluchea</i>	<i>Stachys</i>
<i>Bidens</i>	<i>Gymnopogon</i>	<i>Lipocarpha</i>	<i>Polanisia</i>	<i>Stellaria</i>
<i>Cardamine</i>	<i>Hasteola</i>	<i>Lithospermum</i>	<i>Polygonum</i>	<i>Symphyotrichum</i>
<i>Carex</i>	<i>Helianthemum</i>	<i>Litsea</i>	<i>Potamogeton</i>	<i>Tephrosia</i>
<i>Ceratophyllum</i>	<i>Hexastylis</i>	<i>Lobelia</i>	<i>Prenanthes</i>	<i>Trachelospermum</i>
<i>Chamaecrista</i>	<i>Honckenya</i>	<i>Ludwigia</i>	<i>Prunus</i>	<i>Triglochin</i>
<i>Cicuta</i>	<i>Hottonia</i>	<i>Lycopodiella</i>	<i>Pycnanthemum</i>	<i>Triosteum</i>
<i>Coelorachis</i>	<i>Hudsonia</i>	<i>Lysimachia</i>	<i>Ranunculus</i>	<i>Utricularia</i>
<i>Deschampsia</i>	<i>Hypericum</i>	<i>Lythrum</i>	<i>Rhynchospora</i>	<i>Valeriana</i>
<i>Desmodium</i>	<i>Iris</i>	<i>Matelea</i>	<i>Rumex</i>	<i>Veratrum</i>
<i>Dichanthelium</i>	<i>Juncus</i>	<i>Mecardonia</i>	<i>Sabatia</i>	<i>Xyris</i>
<i>Dryopteris</i>	<i>Krigia</i>	<i>Melothria</i>	<i>Sacciolepis</i>	<i>Zizaniopsis</i>
<i>Elatine</i>	<i>Kyllinga</i>	<i>Morella</i>	<i>Salix</i>	<i>Stellaria</i>
<i>Eleocharis</i>	<i>Lachnanthes</i>	<i>Nymphoides</i>	<i>Scleria</i>	<i>Symphyotrichum</i>
<i>Elephantopus</i>	<i>Lathyrus</i>	<i>Panicum</i>	<i>Scutellaria</i>	

Data for sensitive joint-vetch and seabeach amaranth in National Wildlife Refuges were provided through personal communication with USFWS staff and in NWR reports. Occurrences of seabeach amaranth within the Assateague Island National Seashore were provided by NPS staff. Additional occurrences of rare plant species within Chincoteague and Wallops Island National Wildlife Refuges were provided by USFWS staff via personal communication.

Expert contacts for Chesapeake Bay and the outer coasts of Maryland and Virginia rare plants*:

Name	Agency	City	Phone	Species/ Program
Jack Kumer	NPS	Berlin, MD	410/629-6070	Assateague Island NS
Lauren Billodeaux Mowbray	USFWS	Virginia Beach, VA	757/301-7329	Back Bay NWR
Kevin Holcomb	USFWS	Chincoteague, VA	757/336-6122	Chincoteague and Wallops Island NWRs
Lynn Davidson	MD DNR	Annapolis, MD	410/260-8563	Digital data provider

*Note: this list is not meant to represent all rare plant experts for the region.

Major Data Sources Used: Habitats

MD DNR. 2016. Ecologically Significant Areas of Maryland, vector digital data.

USFWS. 2015. Eastern Virginia Rivers National Wildlife Refuge Complex – sensitive joint-vetch map.

INVASIVE SPECIES

The spread of invasive or non-native species can degrade habitat, increase the potential for crop damage and diseases in humans, livestock and natural resources, reduce biodiversity through competition and limit recreational opportunities. Invasive species often opportunistically spread after disturbance events alter the natural landscape. Oil spill response and clean up often alters the landscape in a manner conducive to the spread of invasive species as crews often mobilize from all over the U.S. in response to large scale spill events. Boats, trailers, waders and clean up equipment can spread invasive species from waterbody to waterbody unless properly cleaned after use. Invasive species that were mapped are shown on the HUMAN-USE RESOURCE maps.

TERRESTRIAL MAMMALS

Terrestrial mammals depicted in this atlas include bats, squirrels, shrews, horses, and otters. Species that are federally and state listed, and those that are vulnerable to oiling and response activities, are specifically emphasized. Threatened and endangered mammals include the Delmarva fox squirrel (VA SE, MD SE), and the Rafinesque's eastern big-eared bat (VA SE). Terrestrial mammal occurrence and seasonality information displayed in this atlas are based on data supplied by the USFWS, NPS, and VDGIF. Note that distributions for Delmarva fox squirrel's (VA SE, MD SE) are areas where they are likely to occur in their preferred habitat. Fox squirrels are primarily found in forests with mixed hardwoods and pines.

Expert contacts for Chesapeake Bay and the outer coasts of Maryland and Virginia terrestrial mammals*:

Name	Agency	City	Phone	Species
Jack Kumer	NPS	Berlin, MD	410/629-6070	Assateague Island NS
Lauren Billodeaux Mowbray	USFWS	Virginia Beach, VA	757/301-7329	Back Bay NWR
Jay Kapalczynski	VDGIF	Henrico, VA	804/367-6796	WERMS Database

*Note: this list is not meant to represent all terrestrial mammal experts for the region.

Major Data Sources Used: Terrestrial Mammals

USFWS. 2008. Delmarva peninsula fox squirrel. Arlington, VA.

VDGIF. 2015. Wildlife environmental review map service, species observation database, vector digital data.

WILDLIFE REHABILITATION

The following contact provides veterinary care and/or retrieval of wildlife adversely affected by an event:
Tri-State Bird Rescue & Research. 170 Possum Hollow Road, Newark, DE 19711. (302)-737-9543.

HUMAN-USE RESOURCES

The human-use resources shown in this atlas were extracted from the ESI GIS data compiled for this region. The extracted features were mapped at scale of 1:100,000 and appear on the maps referenced by a number. For example, Map 1 will show the human-use features in conjunction with the ESI shoreline.

Management areas such as wildlife refuges and state parks are mapped as polygons. Where the feature is a known point location (e.g., marinas, airports, water intakes), the specific location is displayed.

Map IDs can be found in the accompanying data tables for point and polygon features mapped. The Map ID may provide more information (i.e., name, contact) for that particular resource. The types of human use resources mapped in this atlas are depicted below.

Note that some data provided by MD DNR, as part of their Ecologically Significant Areas (ESA) database, were included in the Essential Habitats resource. Species mapped this way include some of the most sensitive species in the ESA database. Many of these species are federally threatened or endangered. MD DNR considers these species so sensitive that they were unwilling to disclose species' names or what ESI element they represent. Without knowing the element these species fall under, they could not be mapped with the biological data.

	Abandoned Vessel		Marina
	Airport		Military Installation
	Anchorage		Mooring Site
	Aquaculture		National Estuarine Research Reserve
	Artificial Reef		National Park
	Beach		Nature Conservancy
	Boat Ramp		Park
	Commercial Fishing		Port
	Critical Habitat		Recreational Fishing
	EPA Facility		Repeated Measurement Site
	Ferry		State Protected Area
	Heliport		Tribal Land
	Historical Site		Water Intake
	Lock and Dam		Wildlife Refuge
	Management Area		

Abandoned Vessel: Data on locations of abandoned and derelict vessels comes from NOAA's Office of Coast Survey Automated Wrecks and Obstructions Information System (AWOIS).

Airport: Locations of airports, airfields, landing strips, etc. were downloaded from the National Transportation Atlas Databases maintained by the Federal Aviation Administration.

Anchorage: Locations where commercial vessels can be anchored offshore were provided by the NOAA Office for Coastal Management.

Aquaculture: Data were provided by the Center for Coastal Resources Management, Virginia Institute of Marine Science, College of William and Mary.

Army Corps of Engineers: U.S. Army Corps of Engineers (USACE) districts and divisions data were provided from the geospatial platform maintained by USACE.

Artificial Reef: Artificial reef data were provided by VA Department of Game and Inland Fisheries, MD DNR, The Nature Conservancy, and NOAA's Office of Coastal Management.

Beach: Locations of recreational beaches used for activities such as swimming, sun-bathing, fishing, etc. were mapped from expert knowledge.

Boat Ramp: Boat ramp location data were provided by the MD DNR, MD Geographic Information Office, VA Department of Conservation and Recreation, VA Department of Game and Inland Fisheries, and the Center for Coastal Resource Management at Virginia Institute of Marine Science, College of William and Mary.

Coast Guard: U.S. Coast Guard (USCG) stations were provided by the USCG and the U.S. Army Corps of Engineers. USCG polygonal district boundaries were provided by the USCG.

Commercial Fishing: Commercial pound net sites were provided by Maryland Department of Natural Resources.

County Boundary: Boundaries were downloaded from the U.S. Census Bureau MAF/TIGER (Topologically Integrated Geographic Encoding and Referencing) geographic database.

Critical Habitat: Designated Critical Habitat was mapped for the Maryland darter (*Etheostoma sellare*). Data were provided by the United States Fish and Wildlife Service.

Environmental Protection Agency Facility: Represents facilities required to file a Risk Management Plan (RMP) due to the presence of extremely hazardous substances that may result in a chemical accident. The data comes from the USEPA.

Environmental Protection Agency Region: Jurisdictional region boundaries were provided by the Environmental Protection Agency.

Essential Habitat: Essential habitat for Maryland's vulnerable, rare, threatened, and/or endangered species were obtained from MD Department of Natural Resources.

Federal Emergency Management Agency Region: Jurisdictional region boundaries were provided by the Federal Emergency Management Agency.

Ferry: U.S. Department of Transportation's Intermodal Passenger Connectivity Database provided ferry terminal data.

Fishery Area: Virginia's Baylor Grounds, public oyster grounds, were provided by the Center for Coastal Resources Management, Virginia Institute of Marine Science, College of William and Mary, in addition to the black drum and oyster management areas. Public Shellfish Fishing Areas in Maryland were mapped by MD DNR.

Heliport: Locations of heliports were downloaded from the National Transportation Atlas Databases maintained by the Federal Aviation Administration.

Historical Site: Historical sites were depicted from sites in the National Park Service National Register of Historic Places (2007) and MD State Historic Preservation Office (The Maryland Historical Trust).

Lock and Dam: Marine lock and dam system data were downloaded from the National Transportation Atlas Database.

Management Area: Management areas in this ESI atlas include: wildlife management areas, nature preserves, and conservation areas and were provided by the VA Department of Conservation and Recreation, VA Department of Game and Inland Fisheries, and MD DNR.

Marina: Marina location data were provided by MD DNR.

Military Installation: Military installation data were provided by the U.S. Census Bureau's MAF/TIGER (Topologically Integrated Geographic Encoding and Referencing) geographic database.

Mooring Site: NOAA's Electronic Navigational Charts (NOAA ENCs) provided mooring locations.

National Estuarine Research Reserve (NERR): NERR boundary data were provided by the Chesapeake Bay Virginia and Maryland NERR.

National Guard: Army National Guard Units were provided by Environmental Systems Research Institute, ArcGIS Online.

National Park: National Park boundaries were provided by the National Park Service (NPS).

Nature Conservancy: Boundaries of The Nature Conservancy (TNC) properties were obtained from TNC Lands database.

NOAA Facility: Locations of NOAA facilities were digitized as points from address matching and verified with Google Earth imagery.

Park: State park boundaries were provided by MA DNR and VA Department of Conservation and Recreation.

Pipeline: Natural gas inter/intrastate lines were provided by the U.S. Energy Information Association.

Port: Major port location data were compiled from ACE National Transportation Atlas databases.

Rail Route: Locations of railways were downloaded from the National Transportation Atlas Databases, maintained by the Department of Transportation.

Recreational Fishing: Recreational fishing locations were provided by Maryland DNR and Virginia Department of Game and Inland Fisheries.

Repeated Measurement Site: Repeated measurement sites obtained from NOAA include: national data buoy locations (DB), Mussel Watch monitoring sites (MWS), National Water Level Observation Network (NWLON), and Physical Oceanographic Real-Time System (PORTS). Water quality monitoring stations (WQ) were obtained from the Chesapeake Bay Program. The Long Term Ecological Research (LTER) Network provided the locations for their LTER Stations.

Road (or Bridge): Bridges were provided by NOAA Continually Updated Shoreline Product (CUSP) and also NOAA National Geodetic Survey's (NGS) Composite/National Shoreline Product.

Shipping Lane: Shipping lanes in U.S. waters were provided by NOAA's Coastal and Marine Spatial Planning office.

State (or State Border): State Borders were compiled from a variety of sources and verified by Research Planning, Inc.

State Protected Area: Areas that receive varied levels of state protection data were provided by VA Department of Game and Inland Fisheries, MD DNR, and NOAA's U.S. Marine Protected Areas Boundaries Inventory.

State Waters: State jurisdictional waters (low water mark extending 3 nm offshore) were derived from the Marine Cadastre, Coastal Zone Management Act offshore boundary.

Tribal Land: Tribal lands were downloaded from the U.S. Census Bureau's Census MAF/TIGER database.

Tunnel: Locations of tunnels were digitized as lines from information provided by Virginia Department of Transportation and Maryland Transportation Authority and verified with Google Earth imagery by Research Planning, Inc.

Washover: A washover, or washover fan, is a relatively flat surface on the top of a barrier spit complex that slopes gently landward. It is usually created when water, forced landward by breaking waves, flows across the top of the barrier spit during high spring tides or storms. This process creates a flattened-off surface along which sand is transported across the top of the spit into the standing water (lagoon) or marsh landward of the spit. The resulting deposit usually has a fan-like shape. Washover locations are represented by points that were digitized from ESRI World Imagery by Research Planning, Inc. at a scale of 1:8,000.

Waste Disposal: Locations represent ocean disposal sites of dredged material removed from the bottom of waterbodies in order to maintain navigation channels and berthing areas. Data were provided by NOAA's Office of Coastal Management.

Water Intake: Location of the fish hatchery water intake was provided by USFWS.

Wildlife Refuge: Twenty-one National Wildlife Refuges (NWR) fall within the ESI atlas. Locations of NWRs were provided by the U.S. Fish and Wildlife Service.

GEOGRAPHIC INFORMATION SYSTEM

All maps were produced using the ESI Geographic Information System (GIS) data compiled for this region. These data are stored as spatial data layers and associated databases. The format for the data varies depending on the type of information or features for which the data are being stored.

Under separate cover is a metadata document that details the data dictionary, processing techniques, data lineage, and other descriptive information for the digital datasets and maps that were used to create this atlas. Below is a brief synopsis of the information contained in the digital version. Refer to the metadata file for a full explanation of the data and its structure.

MAJOR ROADS

The major roads polyline layer represents major thoroughfares within the United States (ArcGIS Content Team (ESRI) and Tele Atlas North America Inc., U.S. Major Roads, ed. 10, published June 30, 2010, ESRI® Data & Maps series, Redlands, CA, USA). These roads are shown on the maps, but are not part of the underlying ESI GIS data.

SHORELINE CLASSIFICATIONS

The ESI shoreline habitat classification is stored as lines and polygons with associated attributes. In many cases, a shoreline may have two or three different classifications or colored lines on the shoreline. These multiple classifications are represented in the database by ESI#1/ESI#2, where ESI#1 is the landward-most classification and ESI#2 is the seaward-most classification. In addition to the line features, marshes (ESI=10A, ESI=10B), swamps (ESI=10C), and scrub-shrub wetlands (ESI=10D) are also stored as polygons.

SENSITIVE BIOLOGICAL RESOURCES

Biological resources are stored as points, lines and polygons. Associated with each feature is a unique identification number that is linked to a series of data tables that further identify the resources. The main biological resource table consists of a list of species identification numbers for each site, the concentration of each species at each site, a mapping qualifier, and identification codes for seasonality and source information. This data table is linked to other tables that describe the seasonality and life-history time periods for each species (at month resolution) for the specified map feature. Other data tables linked to the first table include: the species identification table, which includes common and scientific names; the species status table, which gives information for state and/or federal threatened or endangered listings; and the source database, which provides source metadata at the feature-species level (specific sources are listed for each species occurring at each mapped feature in the biology feature classes).

HUMAN-USE RESOURCES

Human-use features are represented as polygons, points or lines. Management areas such as wildlife refuges, national parks and state parks are mapped as polygons. Known locations such as marinas, high use beaches, airports and water intakes are displayed as points when security risks allow. Bridges and railroads are mapped as line features.

ACKNOWLEDGMENTS

This project was supported by the NOAA Office of Response and Restoration, Hazardous Materials Response Division, under the direction of Jill Petersen, NOAA's ESI Program Manager. The development of this atlas was part of a larger effort to update much of the Atlantic coast after the destruction caused by Hurricane Sandy in October 2012. Funding was provided by the Disaster Relief Appropriations Act of 2013. Frank Csulak (NOAA ERD/Scientific Support Coordinator) assisted greatly in all aspects of the project's completion. Cover photo provided by Joseph Fehrer of the Nature Conservancy.

The biological and human-use data included on the maps were provided by numerous individuals and agencies, including: CCB, MD DNR, NPS, NCBN, NOAA, SHARP, TNC, VAMSC, VCU, VADCR, VDGIF, VIMS, MRC, USFWS and others. Staff at these organizations contributed a vast amount of information to this effort, including first-hand expertise, publications, maps, and digital data.

At Research Planning, Inc. in Columbia, South Carolina, numerous scientific, GIS, and graphic staff were involved with different phases of the project. Mark White, GIS Director, and Christine Boring, Biology Dept. Manager, were co-Project Managers. The ESI shoreline was classified by Jennifer Horsman and Dr. Tim McClinton. The biological and human-use data were collected and compiled onto base maps by Christine Boring, Lauren Szathmary, Lincoln Smith, Michael Bauman, Mike Thompson, and Emily Watson. Lee Diveley, Katy Beckham, Jeff Dahlin, Emily Watson, Chris Locke, and Bryan Thom processed and produced the GIS data. Joe Holmes produced the final documents.

APPROPRIATE USE OF ATLAS AND DATA

This atlas and the associated database were developed to provide summary information on sensitive natural and human-use resources for the purposes of oil and chemical spill planning and response. Although the atlas and database should be very useful for other environmental and natural resource planning purposes, it should not be used in place of data held by any contributing agencies. Likewise, information contained in the atlas and database cannot be used in place of consultations with natural and cultural resource agencies, or in place of field surveys. Also, this atlas should not be used for navigation.

SPECIES LIST

Common Name	Scientific Name	Common Name	Scientific Name
BENTHIC			
SAV			
Submerged aquatic vegetation	-		
BIRDS			
ALCID			
Alcids	-		
DIVING			
American white pelican	<i>Pelecanus erythrorynchos</i>		
Brown pelican	<i>Pelecanus occidentalis</i>		
Common loon	<i>Gavia immer</i>		
Diving birds	-		
Double-crested cormorant	<i>Phalacrocorax auritus</i>		
Grebes	-		
Horned grebe	<i>Podiceps auritus</i>		
Pied-billed grebe	<i>Podilymbus podiceps</i>		
Red-necked grebe	<i>Podiceps grisegena</i>		
Red-throated loon	<i>Gavia stellata</i>		
GULL_TERN			
<u>Black skimmer</u>	<i>Rynchops niger</i>		
Bonaparte's gull	<i>Larus philadelphia</i>		
Caspian tern	<i>Hydroprogne caspia</i>		
Common tern	<i>Sterna hirundo</i>		
Forster's tern	<i>Sterna forsteri</i>		
Great black-backed gull	<i>Larus marinus</i>		
<u>Gull-billed tern</u>	<i>Gelochelidon nilotica</i>		
Gulls	-		
Herring gull	<i>Larus argentatus</i>		
Laughing gull	<i>Larus atricilla</i>		
<u>Least tern</u>	<i>Sternula antillarum</i>		
Ring-billed gull	<i>Larus delawarensis</i>		
<u>Roseate tern</u>	<i>Sterna dougallii</i>		
<u>Royal tern</u>	<i>Thalasseus maximus</i>		
Sandwich tern	<i>Thalasseus sandvicensis</i>		
Terns	-		
PASSERINE			
Coastal plain swamp sparrow	<i>Melospiza georgiana nigrescens</i>		
Le Conte's sparrow	<i>Ammodramus leconteii</i>		
BIRDS, cont.			
PASSERINE, cont.			
Marsh passerine	-		
Marsh wren			<i>Cistothorus palustris</i>
Nelson's sharp-tailed sparrow			<i>Ammodramus nelsoni</i>
Northern parula			<i>Setophaga americana</i>
Prothonotary warbler			<i>Protonotaria citrea</i>
Purple martin			<i>Progne subis</i>
Saltmarsh sharp-tailed sparrow			<i>Ammodramus caudacutus</i>
Savannah sparrow			<i>Passerculus sandwichensis</i>
Seaside sparrow			<i>Ammodramus maritimus</i>
<u>Sedge wren</u>			<i>Cistothorus platensis</i>
Swamp sparrow			<i>Melospiza georgiana</i>
PELAGIC			
Northern gannet			<i>Morus bassanus</i>
Shearwaters			-
Storm-petrels			<i>Oceanodroma spp.</i>
RAPTOR			
Bald eagle			<i>Haliaeetus leucocephalus</i>
Northern harrier			<i>Circus cyaneus</i>
Osprey			<i>Pandion haliaetus</i>
<u>Peregrine falcon</u>			<i>Falco peregrinus</i>
Red-tailed hawk			<i>Buteo jamaicensis</i>
SHOREBIRD			
American oystercatcher			<i>Haematopus palliatus</i>
Black-bellied plover			<i>Pluvialis squatarola</i>
Common snipe			<i>Gallinago gallinago</i>
Dowitchers			<i>Limnodromus spp.</i>
Dunlin			<i>Calidris alpina</i>
Greater yellowlegs			<i>Tringa melanoleuca</i>
Killdeer			<i>Charadrius vociferus</i>
Least sandpiper			<i>Calidris minutilla</i>
Lesser yellowlegs			<i>Tringa flavipes</i>
Phalaropes			<i>Phalaropus spp.</i>
<u>Piping plover</u>			<i>Charadrius melanodus</i>
<u>Red knot</u>			<i>Calidris canutus</i>
Ruddy turnstone			<i>Arenaria interpres</i>
Sanderling			<i>Calidris alba</i>

SPECIES LIST

Common Name	Scientific Name	Common Name	Scientific Name	
BIRDS, cont.			BIRDS, cont.	
SHOREBIRDS, cont.			WATERFOWL	
Semipalmated plover	<i>Charadrius semipalmatus</i>	American black duck	<i>Anas rubripes</i>	
Semipalmated sandpiper	<i>Calidris pusilla</i>	American coot	<i>Fulica americana</i>	
Shorebirds	-	American wigeon	<i>Anas americana</i>	
Short-billed dowitcher	<i>Limnodromus griseus</i>	Blue-winged teal	<i>Anas discors</i>	
Solitary sandpiper	<i>Tringa solitaria</i>	Brant	<i>Branta bernicla</i>	
Spotted sandpiper	<i>Actitis macularia</i>	Bufflehead	<i>Bucephala albeola</i>	
Stilt sandpiper	<i>Calidris himantopus</i>	Canada goose	<i>Branta canadensis</i>	
Western sandpiper	<i>Calidris mauri</i>	Canvasback	<i>Aythya valisineria</i>	
Whimbrel	<i>Numenius phaeopus</i>	Common goldeneye	<i>Bucephala clangula</i>	
Willet	<i>Tringa semipalmata</i>	Common moorhen	<i>Gallinula chloropus</i>	
<u>Wilson's plover</u>	<u><i>Charadrius wilsonia</i></u>	Dabbling ducks	-	
Yellowlegs	<i>Tringa spp.</i>	Diving ducks	-	
WADING			Gadwall	
American bittern	<i>Botaurus lentiginosus</i>	Geese	-	
American woodcock	<i>Scolopax minor</i>	Greater scaup	<i>Aythya marila</i>	
<u>Black rail</u>	<u><i>Laterallus jamaicensis</i></u>	Green-winged teal	<i>Anas crecca</i>	
Black-crowned night-heron	<i>Nycticorax nycticorax</i>	Hooded merganser	<i>Lophodytes cucullatus</i>	
Cattle egret	<i>Bubulcus ibis</i>	Lesser scaup	<i>Aythya affinis</i>	
Clapper rail	<i>Rallus longirostris</i>	Long-tailed duck	<i>Clangula hyemalis</i>	
Glossy ibis	<i>Plegadis falcinellus</i>	Mallard	<i>Anas platyrhynchos</i>	
Great blue heron	<i>Ardea herodias</i>	Mergansers	-	
Great egret	<i>Ardea alba</i>	Northern pintail	<i>Anas acuta</i>	
Green heron	<i>Butorides virescens</i>	Northern shoveler	<i>Anas clypeata</i>	
King rail	<i>Rallus elegans</i>	Red-breasted merganser	<i>Mergus serrator</i>	
Least bittern	<i>Ixobrychus exilis</i>	Redhead	<i>Aythya americana</i>	
Little blue heron	<i>Egretta caerulea</i>	Ring-necked duck	<i>Aythya collaris</i>	
Rails	-	Ruddy duck	<i>Oxyura jamaicensis</i>	
Secretive marsh birds	-	Scaup	<i>Aythya spp.</i>	
Snowy egret	<i>Egretta thula</i>	Scoters	<i>Melanitta spp.</i>	
Sora	<i>Porzana carolina</i>	Snow goose	<i>Chen caerulescens</i>	
Tricolored heron	<i>Egretta tricolor</i>	Surf scoter	<i>Melanitta perspicillata</i>	
Virginia rail	<i>Rallus limicola</i>	Tundra swan	<i>Cygnus columbianus columbianus</i>	
Wading birds		Waterfowl	-	
White ibis	<i>Eudocimus albus</i>	Wood duck	<i>Aix sponsa</i>	
Yellow-crowned night-heron	<i>Nyctanassa violacea</i>	FISH		
FISH				
Alewife			<i>Alosa pseudoharengus</i>	

SPECIES LIST

Common Name	Scientific Name	Common Name	Scientific Name
FISH, cont.		FISH, cont.	
American eel	<i>Anguilla rostrata</i>	Killifish	<i>Fundulus spp.</i>
American shad	<i>Alosa sapidissima</i>	Kingfishes	<i>Menticirrhus spp.</i>
Anchovies	-	Largemouth bass	<i>Micropterus salmoides</i>
Atlantic angel shark	<i>Squatina dumeril</i>	Lined seahorse	<i>Hippocampus erectus</i>
Atlantic croaker	<i>Micropogonias undulatus</i>	Longnose gar	<i>Lepisosteus osseus</i>
Atlantic herring	<i>Clupea harengus</i>	Mullet	<i>Mugil spp.</i>
Atlantic mackerel	<i>Scomber scombrus</i>	Northern puffer	<i>Sphoeroides maculatus</i>
Atlantic menhaden	<i>Brevoortia tyrannus</i>	Northern searobin	<i>Prionotus carolinus</i>
Atlantic needlefish	<i>Strongylura marina</i>	Northern stargazer	<i>Astroscopus guttatus</i>
Atlantic sharpnose shark	<i>Rhizoprionodon terraenovae</i>	Pigfish	<i>Orthopristis chrysoptera</i>
Atlantic spadefish	<i>Chaetodipterus faber</i>	Pinfish	<i>Lagodon rhomboides</i>
<u>Atlantic sturgeon</u>	<u><i>Acipenser oxyrinchus</i></u>	Pipefish	<i>Syngnathus spp.</i>
Black crappie	<i>Pomoxis nigromaculatus</i>	Red drum	<i>Sciaenops ocellatus</i>
Black drum	<i>Pogonias cromis</i>	Red hake	<i>Urophycis chuss</i>
Black sea bass	<i>Centropristes striata</i>	Redfin pickerel	<i>Esox americanus americanus</i>
Blacktip shark	<i>Carcharhinus limbatus</i>	Sand tiger shark	<i>Carcharias taurus</i>
Blue catfish	<i>Ictalurus furcatus</i>	Sandbar shark	<i>Carcharhinus plumbeus</i>
Blueback herring	<i>Alosa aestivalis</i>	Scalloped hammerhead	<i>Sphyraena lewini</i>
Bluefish	<i>Pomatomus saltatrix</i>	Sheepshead	<i>Archosargus probatocephalus</i>
Bullhead catfish	<i>Ameiurus spp.</i>	Shorthead redhorse	<i>Moxostoma macrolepidotum</i>
Butterfish	<i>Peprilus triacanthus</i>	<u>Shortnose sturgeon</u>	<u><i>Acipenser brevirostrum</i></u>
Chain pickerel	<i>Esox niger</i>	Silver hake	<i>Merluccius bilinearis</i>
Channel catfish	<i>Ictalurus punctatus</i>	Silversides	-
Chesapeake logperch	<i>Percina bimaculata</i>	Smallmouth bass	<i>Micropterus dolomieu</i>
Clearnose skate	<i>Raja eglanteria</i>	Smooth butterfly ray	<i>Gymnura micrura</i>
Cobia	<i>Rachycentron canadum</i>	Smooth dogfish	<i>Mustelus canis</i>
Common carp	<i>Cyprinus carpio</i>	Southern flounder	<i>Paralichthys lethostigma</i>
Cownose ray	<i>Rhinoptera bonasus</i>	Southern stingray	<i>Dasyatis americana</i>
Crevalle jack	<i>Caranx hippos</i>	Spanish mackerel	<i>Scomberomorus maculatus</i>
Dusky shark	<i>Carcharhinus obscurus</i>	Spinner shark	<i>Carcharhinus brevipinna</i>
Eastern creek chubsucker	<i>Erimyzon oblongus</i>	Spiny dogfish	<i>Squalus acanthias</i>
Gizzard shad	<i>Dorosoma cepedianum</i>	Spot	<i>Leiostomus xanthurus</i>
Golden shiner	<i>Notemigonus crysoleucas</i>	Spotfin shiner	<i>Cyprinella spiloptera</i>
Great hammerhead	<i>Sphyraena mokarran</i>	Spottail shiner	<i>Notropis hudsonius</i>
Great white shark	<i>Carcharodon carcharias</i>	Spotted hake	<i>Urophycis regia</i>
Harvestfish	<i>Peprilus alepidotus</i>	Spotted seatrout	<i>Cynoscion nebulosus</i>
Hickory shad	<i>Alosa mediocris</i>	Striped bass	<i>Morone saxatilis</i>

SPECIES LIST

Common Name	Scientific Name	Common Name	Scientific Name
FISH, cont.		INVERTEBRATES	
Striped burrfish	<i>Cyclichthys schoepfi</i>	BIVALVE	
Striped cusk-eel	<i>Ophidion marginatum</i>	Alewife floater	<i>Anodonta implicata</i>
Summer flounder	<i>Paralichthys dentatus</i>	Atlantic surfclam	<i>Spisula solidissima</i>
Sunfish	<i>Lepomis spp.</i>	Bay scallop	<i>Argopecten irradians</i>
Tautog	<i>Tautoga onitis</i>	Carolina slabshell	<i>Elliptio congenera</i>
Tessellated darter	<i>Etheostoma olmstedi</i>	BIVALVE	
Thresher shark	<i>Alopias vulpinus</i>	Carolina spike	<i>Elliptio raveneli</i>
Walleye	<i>Stizostedion vitreum vitreum</i>	Creeper	<i>Strophitus undulatus</i>
Weakfish	<i>Cynoscion regalis</i>	<u>Dwarf wedge mussel</u>	<u><i>Alasmidonta heterodon</i></u>
White crappie	<i>Pomoxis annularis</i>	Eastern elliptio	<i>Elliptio complanata</i>
White perch	<i>Morone americana</i>	Eastern floater	<i>Pyganodon cataracta</i>
Winter flounder	<i>Pleuronectes americanus</i>	Eastern lampmussel	<i>Lampsilis radiata</i>
Yellow perch	<i>Perca flavescens</i>	Eastern oyster	<i>Crassostrea virginica</i>
HABITATS		Eastern pondmussel	<i>Ligumia nasuta</i>
PLANT		Elliptio complex	<i>Elliptio sp.</i>
<u>Endangered plant</u>	-	Hard clam	<i>Mercenaria mercenaria</i>
Rare plant community	-	Northern lance	<i>Elliptio fisheriana</i>
<u>Threatened plant</u>	-	Paper pondshell	<i>Utterbackia imbecillis</i>
WETLAND		Softshell clam	<i>Mya arenaria</i>
<u>Seabeach amaranth</u>	<i>Amaranthus pumilus</i>	Stout razor clam	<i>Tagelus plebeius</i>
<u>Sensitive joint-vetch</u>	<i>Aeschynomene virginica</i>	Tidewater mucket	<i>Leptodea ochracea</i>
HERPETOFAUNA		<u>Triangle floater</u>	<u><i>Alasmidonta undulata</i></u>
AMPHIBIAN		Variable spike	<i>Elliptio icterina</i>
<u>Barking treefrog</u>	<i>Hyla gratiosa</i>	Yellow lampmussel	<i>Lampsilis cariosa</i>
TURTLE		CEPHALOPOD	
<u>Chicken turtle</u>	<i>Deirochelys reticularia</i>	Squid	<i>Loligo spp.</i>
Diamondback terrapin	<i>Malaclemys terrapin</i>	CRAB	
<u>Green sea turtle</u>	<i>Chelonia mydas</i>	Atlantic rock crab	<i>Cancer irroratus</i>
<u>Kemp's ridley sea turtle</u>	<i>Lepidochelys kempii</i>	Blue crab	<i>Callinectes sapidus</i>
<u>Leatherback sea turtle</u>	<i>Dermochelys coriacea</i>	Hermit crab	<i>Pagurus spp.</i>
<u>Loggerhead sea turtle</u>	<i>Caretta caretta</i>	Horseshoe crab	<i>Limulus polyphemus</i>
Snapping turtle	<i>Chelydra serpentine</i>	Lady crab	<i>Ovalipes ocellatus</i>
		Knobbed whelk	<i>Busycon carica</i>
Darkling beetle		Darkling beetle	<i>Schoenicus puberulus</i>
<u>Federally threatened insect 1</u>			-
<u>Federally threatened insect 2</u>			-

SPECIES LIST

Common Name	Scientific Name	Common Name	Scientific Name	
INVERTEBRATES, cont.			TERRESTRIAL MAMMALS	
<u>Helops darkling beetle</u>	<i>Helops cisteloides</i>	BAT		
<u>Little white tiger beetle</u>	<i>Ellipsoptera lepida</i>	Rafinesque's eastern big-eared bat	<i>Corynorhinus rafinesquii macrotis</i>	
Monarch butterfly	<i>Danaus plexippus</i>	SMALL MAMMAL		
<u>Northeastern beach tiger beetle</u>	<i>Cicindela dorsalis dorsalis</i>	<u>Delmarva fox squirrel</u>	<i>Sciurus niger cinereus</i>	
<u>Palamedes swallowtail</u>	<i>Papilio palamedes</i>	Dismal swamp southeastern shrew	<i>Sorex longirostris fisheri</i>	
<u>Selys' sunfly</u>	<i>Helocordulia selysi</i>	Northern river otter	<i>Lontra canadensis</i>	
BIVALVE		UNGULATE		
<u>White tiger beetle</u>	<i>Habroscelimorpha dorsalis media</i>	Horse	<i>Equus caballus</i>	
LOBSTER		* Underlined species are listed as either threatened or endangered under the federal ESA, and/or are listed as threatened, endangered, or special concern by Maryland and Virginia		
American lobster	<i>Homarus americanus</i>			
SHRIMP				
Brown shrimp	<i>Farfantepenaeus aztecus</i>			
Grass shrimp	<i>Palaemonetes spp.</i>			
Mantis shrimp	<i>Squilla empusa</i>			
Sand shrimp	<i>Crangon septemspinosa</i>			
White shrimp	<i>Litopenaeus setiferus</i>			
MARINE MAMMALS				
DOLPHIN				
Bottlenose dolphin	<i>Tursiops truncatus</i>			
Harbor porpoise	<i>Phocoena phocoena</i>			
Short-beaked common dolphin	<i>Delphinus delphis</i>			
MANATEE				
<u>West Indian manatee</u>	<i>Trichechus manatus</i>			
PINNIPED				
Gray seal	<i>Halichoerus grypus</i>			
Harbor seal	<i>Phoca vitulina</i>			
WHALE				
<u>Fin whale</u>	<i>Balaenoptera physalus</i>			
<u>Humpback whale</u>	<i>Megaptera novaeangliae</i>			
<u>North Atlantic right whale</u>	<i>Eubalaena glacialis</i>			

SHORELINE DESCRIPTIONS

EXPOSED, ROCKY SHORES

ESI = 1A

DESCRIPTION

- The intertidal zone is steep (greater than 30° slope), with very little width
- Sediment accumulations are uncommon and usually ephemeral, because waves remove the debris that has slumped from the eroding cliffs
- There are few attached plants or animals
- Less than 1% of shoreline

PREDICTED OIL BEHAVIOR

- Oil is held offshore by waves reflecting off the steep cliffs
- Any oil that is deposited is rapidly removed from exposed faces
- The most resistant oil would remain as a patchy band at or above the high-tide line

RESPONSE CONSIDERATIONS

- Cleanup is usually not required
- Access can be difficult and dangerous



Photo by S. Godfrey. © Calvert Marine Museum


SHORELINE DESCRIPTIONS

EXPOSED, SOLID MAN-MADE STRUCTURES

ESI = 1B

DESCRIPTION

- These structures are solid, man-made structures such as seawalls, groins, revetments, piers, and port facilities
- Many structures are constructed of concrete, wood, or metal
- Often there is no exposed substrate at low tide, but multiple habitats are indicated if present
- They are built to protect the shore from erosion by waves, boat wakes, and currents, and thus are exposed to rapid natural removal processes
- Attached animals and plants are sparse to moderate
- They are common in highly developed industrial and naval ports such as Norfolk and scattered along residential waterfronts, occurring along approximately 1% of the shoreline



PREDICTED OIL BEHAVIOR

- Oil is held offshore by waves reflecting off the steep, hard surface in exposed settings
- Oil readily adheres to the dry, rough surfaces, but it does not adhere to wet substrates
- The most resistant oil would remain as a patchy band at or above the high-tide line

RESPONSE CONSIDERATIONS

- Cleanup is usually not required
- High-pressure water spraying may be conducted to:
 - remove persistent oil in crevices;
 - improve aesthetics; or
 - prevent leaching of oil

SHORELINE DESCRIPTIONS

EXPOSED SCARPS AND STEEP SLOPES (CLAY)

ESI = 2B

DESCRIPTION

- These habitats generally occur along tidal channels and major river tributaries in the marsh where the currents cut a steep bank into the marsh soils
- Scarp heights vary from about 1 to 3 feet and usually consist of a heavily rooted, peaty soil
- May be fronted by a narrow beach of fine to medium-grained sand and/or shell fragments
- Low biological utilization because of strong currents and erosion
- Typically backed by wetland vegetation
- Very uncommon, comprising less than 1% of the shoreline



PREDICTED OIL BEHAVIOR

- Oil is not expected to adhere to the wet, impermeable clay surface
- There may be a thin band of oil left at or above the high water line

RESPONSE CONSIDERATIONS

- Cleanup is usually not required, because any stranded oil is quickly removed by wave action
- Access may be difficult


SHORELINE DESCRIPTIONS

FINE- TO MEDIUM-GRAINED SAND BEACHES

ESI = 3A

DESCRIPTION

- These beaches are flat to moderately sloping and relatively hard packed
- They are composed of predominantly quartz sand
- There can be heavy accumulations of wrack present
- They are utilized by birds and turtles
- Upper beach fauna include ghost crabs and amphipods; lower beach fauna can be moderate, but highly variable
- They are generally areas of heavy recreational use
- This shoreline type occurs along the outer coast and sporadic in the bay, comprising about 2% of the shoreline



PREDICTED OIL BEHAVIOR

- Light oil accumulations will be deposited as oily swashes or bands along the upper intertidal zone
- Heavy oil accumulations will cover the entire beach surface; oil will be lifted off the lower beach with the rising tide
- Maximum penetration of oil into fine- to medium-grained sand is about 10-15 cm
- Burial of oiled layers by clean sand within the first week after a spill typically will be less than 30 cm along the upper beach face
- Organisms living in the beach sediment may be killed by smothering or lethal oil concentrations in the interstitial water
- Biological impacts include temporary declines in infauna, which can affect important shorebird foraging areas

RESPONSE CONSIDERATIONS

- These beaches are among the easiest shoreline types to clean
- Cleanup should concentrate on removing oil and oily debris from the upper swash zone once oil has come ashore
- Traffic through both oiled and dune areas should be severely limited, to prevent contamination of clean areas
- Manual cleanup, rather than mechanical removal using road graders and front-end loaders, is advised to minimize the volume of sand removed from the shore and requiring disposal and replacement
- All efforts should focus on preventing the mixing of oil deeper into the sediments by vehicular and foot traffic
- Mechanical reworking of lightly oiled sediments from the high-tide line to the upper intertidal zone can be effective along outer beaches

SHORELINE DESCRIPTIONS

SCARPS AND STEEP SLOPES (SAND)

ESI = 3B

DESCRIPTION

- This shoreline type occurs where sandy bluffs are undercut by waves or currents and slump
- Some scarps are fronted by narrow beaches, if the erosion rates are moderate and episodic
- Trees growing at the top of these slopes are eventually undercut and the logs can accumulate at the base of the scarp
- Biological utilization by birds and infauna is low
- They are uncommon, comprising less than 1% of the shoreline



PREDICTED OIL BEHAVIOR

- Any stranded oil will concentrate at the high-water line and may penetrate sandy sediments
- Oil will also adhere to the dry surfaces of any logs that have accumulated at the base of the scarp
- There is little potential for burial except when major slumping of the bluff occurs
- Active erosion of the scarp will remove the oil

RESPONSE CONSIDERATIONS

- In most cases, removal of oiled sediments is not necessary because of the short residence time of the oil
- The need for removal of oiled sediments and debris should be carefully evaluated because of the potential for increased erosion
- Closely supervised manual labor should be used so that the minimal amount of material is removed during cleanup

SHORELINE DESCRIPTIONS

COARSE-GRAINED SAND BEACHES

ESI = 4

DESCRIPTION

- These beaches have moderate slopes, variable widths, and soft sediments. These characteristics combine to lower their trafficability
- Generally species density and diversity is lower than on fine-grained sand beaches
- They occur along 3% of the shoreline, mostly along banks of major rivers



PREDICTED OIL BEHAVIOR

- During small spills, oil will be deposited primarily as a band along the high-tide line
- Under very heavy accumulations, oil may spread across the entire beach face, though the oil will be lifted off the lower part of the beach with the rising tide
- Penetration of oil into coarse-grained sand can reach 25 cm
- Burial of oiled layers by clean sand can be as rapid as one tidal cycle and to depths of 60 cm or more
- Burial to depths over one meter is possible if the oil comes ashore at the start of a depositional period
- Biological impacts include temporary declines in infaunal populations, which can also affect important shorebird foraging areas

RESPONSE CONSIDERATIONS

- Remove oil primarily from the high tide line
- Removal of sediment should be limited to avoid erosion problems
- Mechanical reworking of the sediment into the surf zone may be used to release the oil without sediment removal
- Activity in the oiled sand should be limited to prevent mixing oil deeper into the beach
- Use of heavy equipment for oil/sand removal may result in the removal of excessive amounts of sand; manual cleanup may be more effective

SHORELINE DESCRIPTIONS

MIXED SAND AND GRAVEL BEACHES

ESI = 5

DESCRIPTION

- Moderately sloping beach composed of a mixture of sand and shell (shell component should comprise between 20 to 80 percent of total sediments)
- Because of the mixed sediment sizes and shapes, there may be zones of pure sand or shell
- They are uncommon, comprising approximately 1% of the shoreline



PREDICTED OIL BEHAVIOR

- During small spills, oil will be deposited along and above the high-tide swash
- Large spills will spread across the entire intertidal area
- Oil penetration into the beach sediments may be up to 50 cm; however, the sand fraction can be quite mobile, and oil behavior is much like on a sand beach if the sand fraction exceeds about 40%
- Burial of oil may be deep at and above the high-tide line, where oil tends to persist, particularly where beaches are only intermittently exposed to waves
- In sheltered pockets on the beach, pavements of asphaltated sediments can form if there is no removal of heavy oil accumulations because most of the oil remains on the surface
- Once formed, these asphalt pavements can persist for years

RESPONSE CONSIDERATIONS

- Remove oil primarily from the high tide line
- Removal of sediment should be limited to avoid erosion problems
- Mechanical reworking of the sediment into the surf zone may be used to release the oil without sediment removal
- Activity in the oiled sediments should be limited to prevent mixing oil deeper into the beach
- Use of heavy equipment for oiled sediment removal may result in the removal of excessive amounts of sediment; manual cleanup may be more effective

SHORELINE DESCRIPTIONS

GRAVEL BEACHES

ESI = 6A

DESCRIPTION

- The gravel fraction is composed of shell fragments; they occur as lag deposits where the finer-grained sediments have been eroded away
- The shells tend to form steep, narrow berms or washover deposits
- Shell beaches occur along areas exposed to high waves or boat wakes, and they comprise less than 1% of the shoreline



PREDICTED OIL BEHAVIOR

- Deep penetration of stranded oil is likely on shell beaches because of their high permeability
- Long-term persistence will be controlled by the depth of routine reworking by waves or boat wakes
- Deeply penetrated oil can leach out for long periods

RESPONSE CONSIDERATIONS

- Heavy accumulations of pooled oil should be removed quickly from the upper beachface
- All oiled debris should be removed
- Sediment removal should be limited as much as possible
- Low-pressure flushing can be used to float fresh oil away from the sediments for recovery by skimmers or sorbents
- Heavily oiled shells may have to be removed and replaced with clean shells

SHORELINE DESCRIPTIONS

RIPRAP

ESI = 6B

DESCRIPTION

- Riprap structures are composed of cobble- to boulder-sized blocks of bedrock or concrete
- Riprap structures are used for shoreline protection and tidal-inlet stabilization
- Attached biota are sparse on exposed riprap and in fresh- to brackish-water areas
- Present along highly developed commercial waterfronts and residential areas, comprising about 3% of the shoreline



PREDICTED OIL BEHAVIOR

- Deep penetration of oil between the blocks is likely
- Oil adheres readily to the rough surfaces of the blocks
- Uncleaned oil can cause chronic leaching until the oil hardens

RESPONSE CONSIDERATIONS

- When the oil is fresh and liquid, high pressure spraying and/or water flooding may be effective, making sure to recover all mobilized oil
- Heavy and weathered oils are more difficult to remove, requiring scraping and/or hot-water spraying
- It may be necessary to remove heavily oiled blocks and replace them

SHORELINE DESCRIPTIONS

BOULDER RUBBLE

ESI = 6D

DESCRIPTION

- Relatively steep rocky shores with accumulations of angular boulder-sized rubble displaying limited evidence of re-working by waves or sediment transport
- Attached biota may be common at lower intertidal levels, whereas biota along the upper intertidal zones is sparse
- Can co-occur with gravel beaches or exposed rocky shorelines; associated gravel beaches can be either at the upper or the lower half of the intertidal zone, depending on the nature of the rock outcrop
- Relatively uncommon and found mostly on the Potomac River upstream from Washington, D.C., comprising less than 1% of the shoreline



Ausra Huntington nee Paulauskaite /Pixels.com

PREDICTED OIL BEHAVIOR

- Oil tends to adhere to the high-water line where the rock surface dries out during low water
- On solid bedrock surfaces, the oil will occur as a surface coating
- Oil will pool and penetrate crevices in the surface rubble
- Where the rubble is loosely packed, oil can penetrate deeply, causing long-term contamination of the subsurface

RESPONSE CONSIDERATIONS

- Thick accumulations of pooled oil should be of high priority for removal, to prevent remobilization and/or penetration
- Manual removal of heavy oil is likely to leave significant residues, but may be useful for thick oil in crevices or sediment pockets

SHORELINE DESCRIPTIONS

EXPOSED TIDAL FLATS

ESI = 7

DESCRIPTION

- Exposed tidal flats are broad, flat intertidal areas composed primarily of sand and minor amounts of shell
- The presence of sand indicates that tidal currents and waves are strong enough to mobilize the sediments
- They are usually associated with another shoreline type on the landward side of the flat, though they can occur as separate shoals; they are commonly associated with tidal inlets
- Biological utilization can be very high, with large numbers of infauna, heavy use by birds for roosting and foraging, and use by foraging fish
- They occur along about 4% of the shoreline, most commonly on the eastern shore



PREDICTED OIL BEHAVIOR

- Oil does not usually adhere to the surface of exposed tidal flats, but rather moves across the flat and accumulates at the high-tide line
- Deposition of oil on the flat may occur on a falling tide if concentrations are heavy
- Oil does not penetrate water-saturated sediments
- Biological damage may be severe, primarily to infauna, thereby reducing food sources for birds and other predators

RESPONSE CONSIDERATIONS

- Currents and waves can be very effective in natural removal of the oil
- Cleanup is very difficult (and possible only during low tides)
- The use of machinery should be restricted to prevent mixing of oil into the sediments

SHORELINE DESCRIPTIONS

SHELTERED SCARPS (BEDROCK/MUD/CLAY)

ESI = 8A

DESCRIPTION

- This shoreline type is sheltered from wave activity and strong currents
- The slope of the intertidal zone is generally moderate to steep (greater than 15°) with little width
- These habitats are not common, occurring along the upper stretches of major rivers and comprising less than 1% of the shoreline



PREDICTED OIL BEHAVIOR

- Stranded oil will persist because of low energy setting
- There will be little potential for penetration into the clayey substrate

RESPONSE CONSIDERATIONS

- Low-pressure flushing at ambient temperatures is most effective when the oil is fresh and still liquid
- Where the high-tide area is accessible, it may be feasible to remove heavy oil accumulations and oiled debris

SHORELINE DESCRIPTIONS

SHELTERED, SOLID MAN-MADE STRUCTURES

ESI = 8B

DESCRIPTION

- These structures are solid man-made structures such as seawalls, groins, revetments, piers, and port facilities
- Most of the structures are designed to protect a single lot, thus their composition, design, and condition are highly variable
- Most structures are constructed of concrete, wood, or metal
- Often there is no exposed beach at low tide, but multiple habitats are indicated if present
- Attached animal and plant life can be moderate to high in salt-water areas
- They are common in highly developed commercial areas such as Norfolk and along residential waterfronts throughout the bay, comprising about 5% of the shoreline



PREDICTED OIL BEHAVIOR

- Oil will adhere readily to rough surfaces, particularly along the high-tide line, forming a distinct oil band
- The lower intertidal zone usually stays wet (particularly if algae covered), preventing oil from adhering to the surface

RESPONSE CONSIDERATIONS

- Cleanup of seawalls is usually conducted for aesthetic reasons or to prevent leaching of oil
- Low- to high-pressure spraying and high-volume deluge at ambient water temperatures is most effective when the oil is fresh

SHORELINE DESCRIPTIONS

SHELTERED RIPRAP

ESI = 8C

DESCRIPTION

- Riprap structures are composed of cobble- to boulder-sized blocks of bedrock or concrete
- These structures are found inside harbors and bays in developed areas, sheltered from direct exposure to waves
- Attached animal and plant life can be present
- They are common in highly developed commercial and residential waterfront areas, comprising 5% of the shoreline



PREDICTED OIL BEHAVIOR

- Deep penetration of oil between the boulders is likely
- Oil adheres readily to the rough surfaces
- If oil is left uncleaned, it may cause chronic leaching until the oil hardens

RESPONSE CONSIDERATIONS

- High-pressure spraying and high-volume deluge may be required to remove oil for aesthetic reasons and to prevent leaching of oil from the structure
- Cleanup crews should make sure to recover all released oil


SHORELINE DESCRIPTIONS

SHELTERED, ROCKY, RUBBLE SHORES

ESI = 8D

DESCRIPTION

- Relatively steep and narrow rocky shore that is covered by a veneer of angular rubble without any evidence of re-working by waves or sediment transport
- The surface rubble is highly variable in size and packing, but there is always some permeability in the surface materials
- Can co-occur with gravel beaches; the gravel beach can be either at the upper or the lower half of the intertidal zone, depending on the nature of the rock outcrop
- This shoreline type is sheltered from significant wave activity and strong currents
- They are uncommon, occurring along less than 1% of the shoreline



PREDICTED OIL BEHAVIOR

- Oil tends to adhere to the upper intertidal zone where the rock surface dries out during low tide, and the algal cover is sparse
- On solid bedrock surfaces, the oil will occur as a surface coating
- Oil will pool and penetrate crevices in the surface rubble
- Where the rubble is loosely packed, oil can penetrate deeply, causing long-term contamination of the subsurface

RESPONSE CONSIDERATIONS

- Thick accumulations of pooled oil should be of high priority for removal, to prevent re-mobilization and/or penetration
- Manual removal of heavy oil is likely to leave significant residues, but may be useful for oil in crevices or sediment pockets
- Flushing techniques will be most effective when oil is still fresh and liquid
- Expect to increase temperature and pressure over time as the oil weathers

SHORELINE DESCRIPTIONS

SHELTERED TIDAL FLATS

ESI = 9A

DESCRIPTION

- Sheltered tidal flats are composed primarily of mud with minor amounts of sand and shell
- They are present in calm-water habitats, sheltered from major wave activity, and are usually backed by marshes
- The sediments are very soft and cannot support even light foot traffic in many areas
- Sheltered tidal flats can be sparsely to heavily covered with algae and/or seagrasses
- They can have heavy wrack deposits along the upper fringe
- Large concentrations of shellfish, worms, and snails can be found on and in the sediments
- They are heavily utilized by birds for feeding
- Sheltered flats are common along most tidal channels, comprising 29% of the shoreline (the second most common habitat type)



PREDICTED OIL BEHAVIOR

- Oil does not usually adhere to the surface of sheltered tidal flats, but rather moves across the flat and accumulates at the high-tide line
- Deposition of oil on the flat may occur on a falling tide if concentrations are heavy
- Oil will not penetrate the water-saturated sediments, but could penetrate burrows or other crevices in muddy sediments
- In areas of high suspended sediments, sorption of oil can result in deposition of contaminated sediments on the flats
- Biological damage may be severe

RESPONSE CONSIDERATIONS

- These are high-priority areas necessitating the use of spill protection devices to limit impacts from both oil exposure and damage during response; deflection or sorbent booms and open water skimmers should be used
- Cleanup of the flat surface is very difficult because of the soft substrate; many methods may be restricted
- Low-pressure flushing and deployment of sorbents from shallow-draft boats may be attempted

SHORELINE DESCRIPTIONS

VEGETATED LOW BANKS

ESI = 9B

DESCRIPTION

- These habitats are either low banks with grasses or trees and tree roots exposed to the water
- They are flooded occasionally by high water
- These shorelines are most common along the major river tributaries and upper sections of tidal creeks to the bay and comprise 16% of the shoreline



PREDICTED OIL BEHAVIOR

- During low-water stages, oil will form a narrow band at the water level
- During high water, the oil will cover and coat the grasses and base of trees
- May cause loss of the grasses, but the trees should survive unless oil penetrates and persists in the substrate

RESPONSE CONSIDERATIONS

- Low-pressure flushing of oiled areas is effective in removing moderate to heavy accumulations of liquid oil from along the banks
- Sorbent and containment boom should be placed on the water side of the cleanup operations to contain and collect oil outflow
- Low- to high-pressure flushing can be used to remove oil from tree roots and trunks, if deemed necessary in high-use areas

SHORELINE DESCRIPTIONS

SALT- AND BRACKISH-WATER MARSHES

ESI = 10A

DESCRIPTION

- These marshes contain emergent, herbaceous vegetation that is salt tolerant; they are often dominated by one species, such as *Spartina alterniflora*
- Width of the marsh can vary widely, from a narrow fringe to extensive areas
- Sediments are composed of organic muds except on the margins of islands where sand is abundant
- Exposed areas are located along bays with wide fetches and along heavily trafficked waterways
- Sheltered areas are not exposed to significant wave or boat wake activity
- Resident flora and fauna are abundant with numerous species with high utilization by birds, fish, and shellfish
- They are the most common shoreline type present, comprising 65% of the shoreline



PREDICTED OIL BEHAVIOR

- Oil adheres readily to intertidal vegetation
- The band of coating will vary widely, depending upon the water level at the time oil slicks are in the vegetation; there may be multiple bands
- Large slicks will persist through multiple tidal cycles and coat the entire stem from the high-tide line to the base
- If the vegetation is thick, heavy oil coating will be restricted to the outer fringe, although lighter oils can penetrate deeper, to the limit of tidal influence
- Medium to heavy oils do not readily adhere to or penetrate the fine sediments, but can pool on the surface or in burrows
- Light oils can penetrate the top few centimeters of sediment and deeply into burrows and cracks (up to one meter)

RESPONSE CONSIDERATIONS

- Under light oiling, the best practice is natural recovery; natural removal processes and rates should be evaluated prior to conducting cleanup
- Heavy accumulations of pooled oil can be removed by vacuum, sorbents, or low-pressure flushing; in situ burning maybe be considered in remote areas
- Cleanup activities should be carefully supervised to avoid vegetation damage
- Any cleanup activity must not mix the oil deeper into the sediments; trampling of the roots must be minimized
- Cutting of oiled vegetation should only be considered when other resources present are at great risk from leaving the oiled vegetation in place

SHORELINE DESCRIPTIONS

FRESHWATER MARSHES

ESI = 10B

DESCRIPTION

- These are grassy wetlands composed of emergent herbaceous vegetation; there can be a mix of species
- They occur upstream of brackish vegetation in the upper estuary and along creeks and rivers
- Those along major channels are exposed to strong currents and boat wakes; smaller channels tend to be sheltered
- Resident flora and fauna are abundant
- Present along tidal freshwater sections of rivers and large creeks, comprising 6% of the shoreline



PREDICTED OIL BEHAVIOR

- Oil adheres readily to the vegetation
- The band of coating will vary widely, depending upon the water level at the time oil slicks are in the vegetation; there may be multiple bands
- Most of the time, there will be a narrow band because of the small tidal range at the head of the estuary; the band can be very large during high-water events
- If the vegetation is thick, heavy oil coating will be restricted to the outer fringe, although lighter oils can penetrate deeper

RESPONSE CONSIDERATIONS

- Under light oiling, the best practice is natural recovery; natural removal processes and rates should be evaluated prior to conducting cleanup
- Heavy accumulations of pooled oil can be removed by vacuum, sorbents, or low-pressure flushing; in situ burning maybe be considered in remote areas
- Cleanup activities should be carefully supervised to avoid vegetation damage
- Any cleanup activity must not mix the oil deeper into the sediments. Trampling of the roots must be minimized
- Cutting of oiled vegetation should only be considered when other resources present are at great risk from leaving the oiled vegetation in place

SHORELINE DESCRIPTIONS

SWAMPS

ESI = 10C

DESCRIPTION

- Swamps consist of shrubs and hardwood forested wetlands, essentially flooded forests. Vegetation is taller, on average, than 20 feet
- The sediment tends to be silty clay with large amounts of organic debris
- They are seasonally flooded, though there are many low, permanently flooded areas
- Resident flora and fauna are abundant with numerous species
- This shoreline type occurs along low areas adjacent to the major tributaries and comprises about 7% of the shoreline



PREDICTED OIL BEHAVIOR

- Oil behavior depends on whether the swamp is flooded or not
- During floods, most of the oil passes through the forest, coating the vegetation at the waterline, which changes levels throughout the flood event
- Oiled woody vegetation is less sensitive than grasses to oil coating
- Some oil can be trapped and pooled on the floodplain as water levels drop
- Penetration into the floodplain soils is usually limited because of high water levels, saturated soils, muddy composition, surface organic debris, and vegetation cover
- Large amounts of oily debris can remain
- During dry periods, terrestrial spills flow downhill and accumulate in depressions or reach waterbodies

RESPONSE CONSIDERATIONS

- Under light oiling, the best practice is to let the area recover naturally
- Heavy accumulations of pooled oil can be removed by vacuum, sorbents, or low-pressure flushing. Care must be taken to prevent transporting oil to sensitive areas down slope or along shore during flushing operations
- Under stagnant water conditions, herding of oil with water spray may be needed to push oil to collection areas
- Oily debris can be removed where there is access
- Any cleanup activity must not mix the oil deeper into the sediments
- Live woody vegetation should not be cut


SHORELINE DESCRIPTIONS

SCRUB AND SHRUB WETLANDS

ESI = 10D

DESCRIPTION

- Scrub-shrub wetlands consist of woody vegetation less than 20 feet tall including true shrubs, small trees, and trees and shrubs that are stunted due to environmental conditions
- The sediments are silty clay mixed with organic debris
- They are seasonally flooded, though there are many low, permanently flooded areas
- Resident flora and fauna are abundant
- This habitat type occurs along about 2% of the shoreline



PREDICTED OIL BEHAVIOR


- Oil behavior depends on water level
- During high water, most of the oil passes through the forest, coating the vegetation above the waterline
- Woody vegetation is less sensitive than grasses to oil
- Some oil can be trapped and pooled on the floodplain as water levels drop
- Penetration into the floodplain soils is usually limited because of high water levels, muddy composition, surface organic debris, and vegetation cover
- Large amounts of oily debris can remain in the wetland
- During dry periods, terrestrial spills flow downhill and accumulate in depressions or reach waterbodies

RESPONSE CONSIDERATIONS

- Under light oiling, the best practice is natural recovery
- Heavy accumulations of pooled oil can be removed by vacuum, sorbents, or low-pressure flushing
- Under stagnant water conditions, herding of oil with water spray may be needed to push oil to collection areas
- Oily debris can be removed where there is access
- Any cleanup activity must not mix the oil deeper into the sediments. Trampling of the roots must be minimized
- Live woody vegetation should not be cut

ENVIRONMENTAL SENSITIVITY INDEX MAPS





Map 49 Central Bay

HUMAN USE RESOURCES

DISPLAYED ON MAP (POINTS)

Map ID	Type	Name	Contact	Phone
1	AIRPORT	ANDERSON FARM		
2	AIRPORT	BLUERMEL FIELD		
3	AIRPORT	CRISFIELD MUNI		
4	BOAT RAMP	BOAT RAMP		
5	BOAT RAMP	COULBOURN CREEK	SOMERSET COUNTY - ROADS DEPARTMENT	
6	BOAT RAMP	JANES ISLAND STATE PARK	MARYLAND DNR - PARK SERVICE	
7	BOAT RAMP	LITTLE BOAT HARBOR	SOMERSET COUNTY - ROADS DEPARTMENT	
8	BOAT RAMP	RUMBLEY	SOMERSET COUNTY - ROADS DEPARTMENT	
9	BOAT RAMP	RUMBLY POINT	SOMERSET COUNTY - ROADS DEPARTMENT	
10	BOAT RAMP	SAINT PETERS CREEK	SOMERSET COUNTY - ROADS DEPARTMENT	
11	BOAT RAMP	SOMERS COVE	MARYLAND DEPARTMENT OF NATURAL RESOURCES	
12	FERRY	CRISFIELD FERRY	MARYLAND DEPARTMENT OF NATURAL RESOURCES	877-620-8367
13	MARINA	CRISFIELD MARINE TERMINAL	MARYLAND DEPARTMENT OF NATURAL RESOURCES	877-620-8367
14	MARINA	SOMERS COVE MARINA	MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
15	REPEATED MEASUREMENT SITE	WQ - SHELLFISH HARVEST WATERS, BIG ANNAMESSEX RIV*	MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
16	REPEATED MEASUREMENT SITE	WQ - SHELLFISH HARVEST WATERS, BIG ANNAMESSEX	MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
17	REPEATED MEASUREMENT SITE	WQ - SHELLFISH HARVEST WATERS, LITTLE ANNAMESSEX	MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
18	REPEATED MEASUREMENT SITE	WQ - SHELLFISH HARVEST WATERS, MANOKIN RIVER	MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
19	REPEATED MEASUREMENT SITE	WQ - SHELLFISH HARVEST WATERS, TANGER SOUND	MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
20	REPEATED MEASUREMENT SITE	WQ - STATION ET8.1	CHESAPEAKE BAY PROGRAM	800-968-7229
21	REPEATED MEASUREMENT SITE	WQ - STATION ET9.1	CHESAPEAKE BAY PROGRAM	800-968-7229

DISPLAYED ON MAP (POLYGONS)

Map ID	Type	Name	Contact	Phone
22	MANAGEMENT AREA	DEAL ISLAND WMA	WILDLIFE AND HERITAGE SERVICE	
23	MANAGEMENT AREA	FAIRMOUNT WMA	WILDLIFE AND HERITAGE SERVICE	
24	MANAGEMENT AREA	MARYLAND MARINE PROPERTIES WMA	WILDLIFE AND HERITAGE SERVICE	
25	MANAGEMENT AREA	POCOMOKE SOUND WMA	WILDLIFE AND HERITAGE SERVICE	
26	MILITARY	COAST GUARD STATION CRISFIELD	MARYLAND PARK SERVICE	
27	PARK	JANES ISLAND SP	MARYLAND PARK SERVICE	

ALSO PRESENT IN MAPPED AREA (POINTS)

Type	Name	Contact	Phone
COAST GUARD	ANT CRISFIELD		
COAST GUARD	CGC CHOKEBERRY		301-968-0971
COAST GUARD	STA CRISFIELD		410-968-0323

ALSO PRESENT IN MAPPED AREA (POLYGONS)

Type	Name	Contact	Phone
STATE PROTECTED AREA	BIG ANNEMESSEX SANCTUARY	MARYLAND DEPARTMENT OF NATURAL RESOURCES	877-620-8367
STATE PROTECTED AREA	KITTS CREEK SANCTUARY	MARYLAND DEPARTMENT OF NATURAL RESOURCES	877-620-8367
STATE PROTECTED AREA	MANOKIN RIVER SANCTUARY	MARYLAND DEPARTMENT OF NATURAL RESOURCES	877-620-8367

JURISDICTIONS

COUNTY:	SOMERSET COUNTY	FEMA:	REGION I
COAST GUARD:	DISTRICT 5, SECTOR BALTIMORE	EPA:	REGION 3
USACE:	NORTH ATLANTIC DIVISION, BALTIMORE DISTRICT		

SHORELINE RESOURCES

ESI POLYGON HABITAT TYPES		
ESI Rank	Habitat Classification	Area (Acres)
10A	Salt and Brackish Water Marshes	24,809.73
10B	Freshwater Marshes	1,041.36
10C	Swamps	6,056.84
10D	Scrub and Shrub Wetlands	1,201.55
9A	Sheltered Tidal Flats	2.85
7	Exposed Tidal Flats	130.11

ESI SHORELINE HABITAT TYPES


ESI Rank	Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
10A	Salt and Brackish Water Marshes	623,792.74	387.61	79%
10C	Swamps	281.85	0.18	< 1%
10D	Scrub and Shrub Wetlands	27,149.24	16.87	3%
9A	Sheltered Tidal Flats	17,138.82	10.65	2%
9B	Vegetated Low Banks	22,534.89	14.00	3%
8B	Sheltered, Solid Man-Made Structures	12,118.46	7.53	2%
8C	Sheltered Ripprap	12,592.64	7.82	2%
7	Exposed Tidal Flats	26,306.18	16.35	3%
6A	Gravel Beaches	156.75	0.10	< 1%
6B	Ripprap	1,560.07	0.97	< 1%
5	Mixed Sand and Gravel Beaches	70.73	0.04	< 1%
4	Coarse Grained Sand Beaches	42,844.70	26.62	5%
1B	Exposed, Solid Man-Made Structures	68.56	0.04	< 1%

Total ESI Shoreline: 786,615.62 Total ESI Shoreline: 488.78
 Total Shoreline: 651,265.76 Total Shoreline: 404.68

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%.

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 49A
Central Bay



Map 49A Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	Mapping Qualifier			S	F	Concentration	Monthly Presence										
		High Ecological Value	High Ecological Value	High Ecological Value				J	F	M	A	M	J	J	A	S	O	N
SAV	Submerged aquatic veg	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
SAV	Submerged aquatic veg	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
SAV	Submerged aquatic veg	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

BIRDS

Map ID	Subelement	Mapping Qualifier			S	F	Concentration	Monthly Presence											
		Great blue heron	Nesting	Nesting				J	F	M	A	M	J	J	A	S	O	N	D
		Bald eagle	Nesting	Nesting				5 Nests	Feb	Feb	Mar	Apr	Feb	Apr	Feb	Apr	Mar	Aug	-
418	Raptor	Bald eagle	Nesting	Nesting	-	-	-	Last Obs. 2004	-	-	-	-	-	-	-	-	-	-	
427	Raptor	Bald eagle	Nesting	Nesting	-	-	-	Last Obs. 1994	-	-	-	-	-	-	-	-	-	-	
440	Waterfowl	American black duck	Nesting	Nesting	-	-	-	Moderate	-	-	-	-	-	-	-	-	-	-	
533	Waterfowl	Bufflehead	Wintering	Wintering	-	-	-	1,000S	-	-	-	-	-	-	-	-	-	-	
	Waterfowl	Canada goose	Wintering	Wintering	-	-	-	1,000S	-	-	-	-	-	-	-	-	-	-	
	Waterfowl	Canvasback	Wintering	Wintering	-	-	-	1,000S	-	-	-	-	-	-	-	-	-	-	
	Waterfowl	Common goldeneye	Wintering	Wintering	-	-	-	10S	-	-	-	-	-	-	-	-	-	-	
	Waterfowl	Mergansers	Wintering	Wintering	-	-	-	100S	-	-	-	-	-	-	-	-	-	-	
	Waterfowl	Redhead	Wintering	Wintering	-	-	-	1,000S	-	-	-	-	-	-	-	-	-	-	
	Waterfowl	Ring-necked duck	Wintering	Wintering	-	-	-	10S	-	-	-	-	-	-	-	-	-	-	
	Waterfowl	Ruddy duck	Wintering	Wintering	-	-	-	1,000S	-	-	-	-	-	-	-	-	-	-	
	Waterfowl	Scaup	Wintering	Wintering	-	-	-	100S	-	-	-	-	-	-	-	-	-	-	
	Waterfowl	Snow goose	Wintering	Wintering	-	-	-	1,000S	-	-	-	-	-	-	-	-	-	-	
	Waterfowl	Tundra swan	Wintering	Wintering	-	-	-	100S	-	-	-	-	-	-	-	-	-	-	
1169	Waterfowl	American black duck	Nesting	Nesting	-	-	-	Moderate-High	-	-	-	-	-	-	-	-	-	-	

FISH

Map ID	Subelement	Mapping Qualifier			S	F	Concentration	Monthly Presence											
		Nursery Area	Nursery Area	Nursery Area				J	F	M	A	M	J	J	A	S	O	N	D
		Spawning Area	Spawning Area	Spawning Area				-	-	-	-	-	-	-	-	-	-	-	
1416	Diadromous	Alewife	Alewife	Alewife	-	-	-	Mar-May	Mar-May	Mar-May	Mar-May	Mar-May	May-Jul	May-Jul	May-Oct	Apr-Jun	Apr-Oct	-	
	Diadromous	American shad	American shad	American shad	-	-	-	Apr-Jun	Apr-Jun	Apr-Jun	Apr-Jun	Apr-Jun	-	-	-	-	-	Mar-May	
	Diadromous	American shad	American shad	American shad	-	-	-	Apr-Jun	Apr-Jun	Apr-Jun	Apr-Jun	Apr-Jun	-	-	-	-	-	Mar-Jun	
	Diadromous	Blueback herring	Blueback herring	Blueback herring	-	-	-	Apr-Jun	Apr-Jun	Apr-Jun	Apr-Jun	Apr-Jun	-	-	-	-	-	Feb-Jun	
	Diadromous	Hickory shad	Hickory shad	Hickory shad	-	-	-	Apr-May	Apr-May	Apr-May	Apr-May	Apr-May	-	-	-	-	-	Feb-Jun	
	Diadromous	Striped bass	Striped bass	Striped bass	-	-	-	Apr-Jun	Apr-Jun	Apr-Jun	Apr-Jun	Apr-Jun	-	-	-	-	-	Mar-Jun	
	Diadromous	Spawning Area	Spawning Area	Spawning Area	-	-	-	Apr-Jun	Apr-Jun	Apr-Jun	Apr-Jun	Apr-Jun	-	-	-	-	-	Mar-Jun	

FISH (continued)

Map ID	Subelement	Species	MD/VA		Mapping Qualifier	S	F	Concentration	Monthly Presence						
			J	F					J	F	M	A	M	J	
	Freshwater	Yellow perch	-	-	Spawning Area	-	-	-	-	-	Feb-Mar	Feb-Mar	Eggs	Larvae	Juveniles
1417	Diaxromous	Alewife	Alewife	-	Spawning Area	Nursery Area	-	-	Mar-May	Mar-May	-	Apr-Jun	Mar-Apr	Mar-Apr	Adults
	Diaxromous	American shad	American shad	-	Spawning Area	Nursery Area	-	-	-	-	May-Oct	-	-	-	Feb-Mar
	Diaxromous	American shad	American shad	-	Spawning Area	Nursery Area	-	-	Apr-Jun	Apr-Jun	-	-	-	-	Mar-May
	Diaxromous	Blueback herring	Blueback herring	-	Spawning Area	Nursery Area	-	-	Apr-Jun	Apr-Jun	-	May-Jul	May-Oct	-	Mar-Jun
	Diaxromous	Blueback herring	Hickory shad	-	Spawning Area	Nursery Area	-	-	Apr-Jun	Apr-Jun	-	Apr-Jun	Apr-Oct	-	Feb-Jun
	Diaxromous	Hickory shad	Hickory shad	-	Spawning Area	Nursery Area	-	-	Apr-May	Apr-May	-	-	-	-	Feb-Jun
	Diaxromous	Striped bass	Striped bass	-	Spawning Area	Nursery Area	-	-	Apr-Jun	Apr-Jun	-	-	-	-	Mar-Jun
1521	Freshwater	Yellow perch	Yellow perch	-	Spawning Area	Nursery Area	-	-	Feb-Mar	Feb-Mar	Feb-Mar	Feb-Mar	Mar-Apr	Mar-Apr	Feb-Mar
1539	Diaxromous	Striped bass	Striped bass	-	Spawning Area	Nursery Area	-	-	-	-	Apr-Jun	Apr-Jun	Jan-Dec	-	-
	Freshwater	Yellow perch	Yellow perch	-	Spawning Area	Nursery Area	-	-	Feb-Mar	Feb-Mar	Feb-Mar	Feb-Mar	Mar-Apr	Mar-Apr	Feb-Mar
1670	Diaxromous	Alewife	Alewife	-	Spawning Area	Nursery Area	-	-	-	-	Apr-Jun	Apr-Jun	Apr-Oct	Apr-Oct	Mar-May
	Diaxromous	American shad	American shad	-	Spawning Area	Nursery Area	-	-	-	-	May-Jul	May-Jul	May-Oct	May-Oct	Mar-Jun
	Diaxromous	Blueback herring	Blueback herring	-	Spawning Area	Nursery Area	-	-	-	-	May-Jul	May-Jul	May-Oct	Feb-Jun	Feb-Mar
	Diaxromous	Hickory shad	Hickory shad	-	Spawning Area	Nursery Area	-	-	-	-	Apr-Jun	Apr-Jun	Apr-Oct	Apr-Oct	Feb-Jun
	Diaxromous	Striped bass	Striped bass	-	Spawning Area	Nursery Area	-	-	-	-	Apr-Jun	Apr-Jun	-	-	Mar-Jun
1681	Diaxromous	Alewife	Blueback herring	-	Spawning Area	Nursery Area	-	-	-	-	Apr-Jun	Apr-Jun	Apr-Oct	Apr-Oct	Mar-May
	Diaxromous	Blueback herring	Blueback herring	-	Spawning Area	Nursery Area	-	-	-	-	May-Jul	May-Oct	Feb-Jun	-	-

HABITATS & RARE PLANTS

Map ID	Subelement	Species	MD/VA		Mapping Qualifier	S	F	Concentration	Monthly Presence					
			J	F					J	F	M	A	M	J
	Plant	Endangered plant	Vulnerable Occurrence	E/-	2 Species Present	-	-	-	-	-	-	-	-	-
1748	Plant	Endangered plant	Vulnerable Occurrence	E/-	2 Species Present	-	-	-	-	-	-	-	-	-

REPTILES & AMPHIBIANS

Map ID	Subelement	Species	MD/VA		Mapping Qualifier	S	F	Concentration	Monthly Presence					
			J	F					J	F	M	A	M	J
1775	Turtle	K. ridley sea turtle	General Distribution	E/E	Uncommon	-	-	-	-	-	-	-	-	-
	Turtle	K. ridley sea turtle	General Distribution	E/E	Common	-	-	-	-	-	-	-	-	-
	Turtle	Loggerhead sea turtle	General Distribution	T/T	Common	-	-	-	-	-	-	-	-	-
	Turtle	Loggerhead sea turtle	General Distribution	T/T	Abundant	-	-	-	-	-	-	-	-	-
1781	Turtle	Diamondback terrapin	Nesting	T/T	Possible Nest	May-Aug	-	-	-	-	-	-	-	-

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

BIRDS

Subelement	Species	MD\VA				Monthly Presence															
		Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
Diving	Common loon	Wintering	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	D. crested cormorant	Migration	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Grebes	Wintering	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Red-throated loon	Wintering	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Pelagic	Northern gannet	Nesting	Low	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Waterfowl	American black duck	Wintering	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	American coot	Wintering	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
American wigeon	Wintering	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Canada goose	Nesting	Moderate-High	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Gadwall	Wintering	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Green-winged teal	Wintering	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Long-tailed duck	Wintering	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Mallard	Nesting	Moderate-High	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Wintering	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Mallard	Wintering	Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Northern pintail	Wintering	10,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Northern shoveler	Wintering	Moderate-High	10,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Scoters	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Wood duck	Nesting	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Subelement	Species	MD\VA				Monthly Presence															
		Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles
Diacromous	Atlantic sturgeon	General Distribution	-E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Shortnose sturgeon	General Distribution	E/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Estuarine Nursery	White perch	Nursery Area	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

INVERTEBRATES

Subelement	Species	MD\VA				Monthly Presence															
		Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles
Crab	Blue crab	Concentration Area	Highly Abundant-Summ	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Oct
		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Oct

MARINE MAMMALS

Subelement	Species	MD\VA				Monthly Presence															
		Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Matting	Calving	Pupping	Molt
Manatee	West Indian manatee	General Distribution	-E	E	Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

TERRESTRIAL MAMMALS

Subelement	Species	MD\VA				Monthly Presence															
		Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Matting	Calving	Pupping	Molt
Small Mammal	Delmarva fox squirrel	General Distribution	E/E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

BIRDS

Sublement	Species	<i>Monthly Presence</i>												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
Diving	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-	-
Gull/Tern	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-	-
Gulls	Gulls	-	-	-	-	-	-	-	-	-	-	-	-	-
Terns	Terns	-	-	-	-	-	-	-	-	-	-	-	-	-
Passerine	Seaside sparrow	-	-	-	-	-	-	-	-	-	-	-	-	-
Wading	Clapper rail	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Sublement	Species	<i>Monthly Presence</i>												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Diadromous	Alewife	-	-	-	-	-	-	-	-	-	-	-	-	Mar-May
	American eel	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct
	American shad	-	-	-	-	-	-	-	-	-	-	-	-	Jan-May
	Blueback herring	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Gizzard shad	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Jun
	Hickory shad	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Striped bass	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Estuarine Nursery	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Atlantic menhaden	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Black drum	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Black sea bass	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Bluefish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Red drum	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Spot	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Dec
	Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-	Dec-Apr
	Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-	Feb-May
	Weakfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Aug
	White perch	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Estuarine Resident	Silversides	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
Fish	Anchoovies	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Striped burrfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Freshwater	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Yellow perch	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
Marine Benthic	Atlantic spadefish	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Marine Pelagic	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Sep
	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov

INVERTEBRATES

Subelement	Species	Monthly Presence											
		J	F	M	A	M	J	J	A	S	O	N	D
Crab	Blue crab	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Horseshoe crab	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec

MARINE MAMMALS

Subelement	Species	Monthly Presence											
		J	F	M	A	M	J	J	A	S	O	N	D
Dolphin	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES

ESI Rank	Habitat Classification	Area (Acres)											
		3,206.56	5.01	484.78	0.76	2,277.78	3.56	372.58	0.58	1.65	0.00	1.65	0.00
10A	Salt and Brackish Water Marshes												
10B	Freshwater Marshes												
10C	Swamps												
10D	Scrub and Shrub Wetlands												
7	Exposed Tidal Flats												

ESI SHORELINE HABITAT TYPES

ESI Rank	Shoreline Habitat Classification	Length (Meters)												
		103,740.07	64,46	85%	10,440.00	6,49	9%	3,649.32	2,27	3%	343.34	<1%	1,655.26	1,03
10A	Salt and Brackish Water Marshes													
10D	Scrub and Shrub Wetlands													
9B	Vegetated Low Banks													
8B	Sheltered, Solid Man-Made Structures													
8C	Sheltered Riprap													
7	Exposed Tidal Flats													
4	Coarse Grained Sand Beaches													


Total ESI Shoreline: 122,214.85
Total Shoreline: 105,921.90

Total ESI Shoreline: 75.94
Total Shoreline: 65.82

Note: A shoreline segment may include multiple habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 49B
Central Bay



SEE BACK OF MAP
for details about mapped species and
other species that occur in mapped area.
Data Published: September 2016



Map 49B Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	Mapping Qualifier			Concentration	Monthly Presence										
		S	F	MD\VA		J	F	M	A	M	J	J	A	S	O	N
SAV	Submerged aquatic veg	High Ecological Value			Ephemeral	-	-	-	-	-	-	-	-	-	-	-
SAV	Submerged aquatic veg	High Ecological Value			High	-	-	-	-	-	-	-	-	-	-	-
SAV	Submerged aquatic veg	High Ecological Value			Low	-	-	-	-	-	-	-	-	-	-	-

BIRDS

Map ID	Subelement	Species	Mapping Qualifier			Concentration	Monthly Presence										
			S	F	MD\VA		J	F	M	A	M	J	J	A	S	O	N
341	Gull/Tern	Common tern	Nesting			1 Pair	-	-	-	-	-	-	-	-	-	-	-
	Gull/Tern	Forster's tern	Nesting			20 Pairs	-	-	-	-	-	-	-	-	-	-	-
351	Gull/Tern	Forster's tern	Nesting			3 Pairs	-	-	-	-	-	-	-	-	-	-	-
418	Raptor	Bald eagle	Nesting			Last Obs. 2004	-	-	-	-	-	-	-	-	-	-	-
1283	Gull/Tern	Forster's tern	Nesting			109 Pairs	-	-	-	-	-	-	-	-	-	-	-
	Gull/Tern	G. black-backed gull	Nesting			1 Pair	-	-	-	-	-	-	-	-	-	-	-

FISH

Map ID	Subelement	Species	Mapping Qualifier			Concentration	Monthly Presence										
			S	F	MD\VA		J	F	M	A	M	J	J	A	S	O	N
1521	Freshwater	Yellow perch	Spawning Area			-	-	-	-	-	-	-	-	-	-	-	-
1631	Estuarine Nursery	White perch	Nursery Area			-	-	-	-	-	-	-	-	-	-	-	-
1661	Diadromous	Alewife	Migration			-	-	-	-	-	-	-	-	-	-	-	-
	Diadromous	American shad	Migration			-	-	-	-	-	-	-	-	-	-	-	-
	Diadromous	Blueback herring	Migration			-	-	-	-	-	-	-	-	-	-	-	-
	Diadromous	Hickory shad	Migration			-	-	-	-	-	-	-	-	-	-	-	-

REPTILES & AMPHIBIANS

Map ID	Subelement	Species	Mapping Qualifier			Concentration	Monthly Presence										
			S	F	MD\VA		J	F	M	A	M	J	J	A	S	O	N
1781	Turtle	Diamondback terrapin	Nesting			Possible Nest	-	-	-	-	-	-	-	-	-	-	-
1782	Turtle	Diamondback terrapin	Nesting			Probable Nest	-	-	-	-	-	-	-	-	-	-	-


INVERTEBRATES

Map ID	Subelement	Species	Mapping Qualifier			Concentration	Monthly Presence										
			S	F	MD\VA		J	F	M	A	M	J	J	A	S	O	N
1827	Insect	Vulnerable Occurrence	E/T	T	-	-	-	-	-	-	-	-	-	-	-	-	-
	Federally threatened insect 1																

MARINE MAMMALS

<i>Map ID</i>	<i>Subelement</i>	<i>Species</i>	<i>Mapping Qualifier</i>	<i>MD/VA</i>	<i>S F</i>	<i>Concentration</i>	<i>Monthly Presence</i>															
		Humpback whale	General Distribution	E/E	E	Very Rare	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping	Molt
WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)																						

BIRDS



FISH

Subelement	Species	MD/VA												Monthly Presence											
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults	Apr-Jun	Apr-Oct	Mar-May	Mar-Jun
Diadromous	Alewife	Nursery Area	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	American shad	Nursery Area	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Atlantic sturgeon	General Distribution	-/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Blueback herring	Nursery Area	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Hickory shad	Nursery Area	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Shortnose sturgeon	General Distribution	E/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Striped bass	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Striped bass	Nursery Area	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

REPTILES & AMPHIBIANS

Subelement	Species	MD/VA												Monthly Presence											
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles	Adults	Apr-Oct	May-Oct	Mar-Jun	Jan-Dec
Turtle	K. ridley sea turtle	General Distribution	E/E	E	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	K. ridley sea turtle	General Distribution	E/E	E	Uncommon	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Loggerhead sea turtle	General Distribution	T/T	T	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Loggerhead sea turtle	General Distribution	T/T	T	Abundant	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

INVERTEBRATES

Subelement	Species	MD/VA												Monthly Presence											
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles	Adults	Apr-Oct	May-Oct	Mar-Jun	Jun-Oct
Crab	Blue crab	Concentration Area	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		Highly Abundant-Summ	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

MARINE MAMMALS

Subelement	Species	MD/VA												Monthly Presence											
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles	Adults	Apr-Oct	May-Oct	Mar-Jun	Jan-Dec
Manatee	West Indian manatee	General Distribution	-/E	E	Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

TERRESTRIAL MAMMALS

Subelement	Species	MD/VA												Monthly Presence											
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles	Adults	Apr-Oct	May-Oct	Mar-Jun	Jan-Dec
Small Mammal	Delmarva fox squirrel	General Distribution	E/E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)

BIRDS

Subelement	Species	MD/VA												Monthly Presence											
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt	Adults	Apr-Jun	Apr-Oct	May-Aug	May-Jul
Diving	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Gull/Tern	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Gulls	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Passerine	Terns	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Shorebird	Saltmarsh sharp-tailed sparrow	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Wading	Seaside sparrow	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Willet	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Clapper rail	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

		Monthly Presence															
		Species		J F M A M J J A S O N D			Spawn			Eggs		Larvae		Juveniles		Adults	
Subelement	Subelement	Alewife								Mar-May	Mar-May	Apr-Jun	Apr-Jun	Jan-May	Jan-May	Apr-Oct	Mar-May
Diadromous	Alewife	American eel								Mar-May	Mar-May	Apr-Jun	Apr-Jun	Jan-May	Jan-May	Jan-Dec	Jan-Dec
		Blueback herring								Apr-Jun	Apr-Jun	May-Jul	May-Jul	Jan-Dec	Jan-Dec	Feb-Jun	Feb-Jun
		Gizzard shad								Mar-Jun	Mar-Jun	-	-	Jan-Dec	Jan-Dec	Jan-Dec	Jan-Dec
		Striped bass								-	-	-	-	Jan-Dec	Jan-Dec	Jan-Dec	Jan-Dec
Estuarine Nursery	Atlantic menhaden	Atlantic croaker								Apr-Jun	Apr-Jun	-	-	Dec-Apr	Jan-Dec	May-Nov	May-Nov
		Black drum								-	-	-	-	Apr-Dec	Apr-Dec	Apr-Dec	Apr-Dec
		Black sea bass								-	-	-	-	Apr-Nov	Apr-Nov	Apr-Nov	Apr-Nov
		Bluefish								-	-	-	-	May-Oct	May-Oct	May-Oct	May-Oct
		Northern puffer								-	-	-	-	Mar-Oct	Mar-Oct	Mar-Nov	Mar-Nov
		Red drum								-	-	-	-	-	-	May-Nov	May-Nov
		Spot								-	-	-	-	Nov-Apr	Nov-Apr	Apr-Dec	Apr-Dec
		Spotted seatrout								-	-	-	-	Aug-Dec	Aug-Dec	Apr-Dec	Apr-Dec
		Summer flounder								-	-	-	-	Feb-May	Feb-May	Apr-Dec	Jan-Dec
		Weakfish								-	-	-	-	May-Aug	May-Aug	Apr-Dec	Apr-Dec
		White perch								-	-	-	-	Jun-Dec	Jun-Dec	Jan-Dec	Jan-Dec
Estuarine Resident	Silversides	Anchovies								Mar-Jul	Mar-Jul	-	-	Mar-Nov	Mar-Nov	Mar-Nov	Mar-Nov
		Striped burrfish								Apr-Sep	Apr-Sep	-	-	-	-	May-Oct	May-Oct
		Yellow perch								-	-	-	-	Jan-Dec	Jan-Dec	Jan-Dec	Jan-Dec
		Atlantic spadefish								-	-	-	-	Jun-Sep	Jun-Sep	May-Oct	May-Oct
		Cownose ray								Jun-Jul	Jun-Jul	-	-	-	-	May-Oct	May-Oct
		Northern searobin								May-Jul	May-Jul	-	-	Apr-Dec	Apr-Dec	Jan-Dec	Jan-Dec
		Spotted hake								-	-	-	-	Jan-Dec	Jan-Dec	Jan-Dec	Jan-Dec
		Butterfish								-	-	-	-	Jul-Sep	Jul-Sep	Mar-Nov	Mar-Nov
		Harvestfish								-	-	-	-	-	-	May-Nov	May-Nov

INVERTEBRATES

		Monthly Presence															
		Species		J F M A M J J A S O N D			Spawn			Eggs		Larvae		Juveniles		Adults	
Subelement	Subelement	Eastern oyster								Jun-Sep	Jun-Sep	Jun-Oct	Jun-Oct	Jan-Dec	Jan-Dec	Jan-Dec	Jan-Dec
		Hard clam								Jun-Sep	Jun-Sep	-	-	Jan-Dec	Jan-Dec	Jan-Dec	Jan-Dec
		Softshell clam								May-Dec*	-	-	-	Jan-Dec	Jan-Dec	Jan-Dec	Jan-Dec
		Blue crab								-	-	-	-	Jan-Dec	Jan-Dec	Jan-Dec	Jan-Dec
		Horseshoe crab								-	-	-	-	Jan-Dec	Jan-Dec	Jan-Dec	Jan-Dec

MARINE MAMMALS

		Monthly Presence															
		Species		J F M A M J J A S O N D			Spawn			Eggs		Larvae		Juveniles		Adults	
Subelement	Subelement	Eastern oyster								Jun-Sep	Jun-Sep	Jun-Oct	Jun-Oct	Jan-Dec	Jan-Dec	Jan-Dec	Jan-Dec
		Bottlenose dolphin								May-Aug	-	-	-	-	-	-	-
		Bottlenose dolphin								-	-	-	-	-	-	-	-
		Harbor porpoise								-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

ESI POLYGON HABITAT TYPES		
	ESI Rank	Habitat Classification
	10A	Salt and Brackish Water Marshes
	10B	Freshwater Marshes
	10C	Swamps
	10D	Scrub and Shrub Wetlands
		Sheltered Tidal Flats
	9A	Exposed Tidal Flats
	7	


ESI SHORELINE HABITAT TYPES		
	ESI Rank	Shoreline Habitat Classification
	10A	Salt and Brackish Water Marshes
	10D	Scrub and Shrub Wetlands
	9A	Sheltered Tidal Flats
	9B	Vegetated Low Banks
	8B	Sheltered, Solid Man-Made Structures
	8C	Sheltered Riprap
	7	Exposed Tidal Flats
	6B	Riprap
	5	Mixed Sand and Gravel Beaches
	4	Coarse Grained Sand Beaches
	1B	Exposed, Solid Man-Made Structures

Total ESI Shoreline: 279,408.67 Total ESI Shoreline: 173.62
 Total Shoreline: 233,762.71 Total Shoreline: 145.25

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 49C
Central Bay



SEE BACK OF MAP
for details about mapped species and
other species that occur in mapped area.
Data Published: September 2016



Map 49C Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	Mapping Qualifier			S	F	Concentration	Monthly Presence										
		High Ecological Value	High Ecological Value	High Ecological Value				J	F	M	A	M	J	J	A	S	O	N
SAV	Submerged aquatic veg							-	-	-	-	-	-	-	-	-	-	-
SAV	Submerged aquatic veg							-	-	-	-	-	-	-	-	-	-	-
SAV	Submerged aquatic veg							-	-	-	-	-	-	-	-	-	-	-

BIRDS

Map ID	Subelement	Species	Mapping Qualifier			S	F	Concentration	Monthly Presence										
			J	F	M				J	F	M	A	M	J	J	A	S	O	N
418	Raptor	Bald eagle	Nesting					Last Obs. 2004											
484	Watertowl	Wood duck	Nesting																
487	Watertowl	American black duck	Nesting																
	Watertowl	Wood duck	Nesting																
533	Watertowl	Bufflehead	Wintering																
	Watertowl	Canada goose	Wintering																
	Watertowl	Canvasback	Wintering																
	Watertowl	Common goldeneye	Wintering																
	Watertowl	Mergansers	Wintering																
	Watertowl	Redhead	Wintering																
	Watertowl	Ring-necked duck	Wintering																
	Watertowl	Ruddy duck	Wintering																
	Watertowl	Scaup	Wintering																
	Watertowl	Snow goose	Wintering																
	Watertowl	Tundra swan	Wintering																
707	Raptor	Bald eagle	Roosting																
954	Watertowl	Bufflehead	Wintering																
	Watertowl	Canvasback	Wintering																
	Watertowl	Mergansers	Wintering																
	Watertowl	Redhead	Wintering																
	Watertowl	Ring-necked duck	Wintering																
	Watertowl	Ruddy duck	Wintering																
	Watertowl	Scaup	Wintering																
	1089	Common loon	Wintering																
	Diving	D. crested cormorant	Migration																
	Diving	Grebes	Wintering																
	Diving	Red-throated loon	Wintering																
	Pelagic	Northern gannet	Wintering																
	Watertowl	Long-tailed duck	Wintering																
	Watertowl	Scoter	Wintering																

BIRDS (continued)

Map ID	Subelement	Species	MD/VA						Monthly Presence											
			S	F	Concentration	J	F	M	A	M	J	J	A	S	N	D	Nest	Mig.(S)	Mig.(F)	Molt
1164	Diving	Common loon	Wintering														-	-	-	-
	Diving	Grebes	Wintering														-	-	-	-
	Diving	Red-throated loon	Wintering														-	-	-	-
	Pelagic	Northern gannet	Wintering														-	-	-	-
	Watertowl	Long-tailed duck	Wintering														-	-	-	-
	Watertowl	Scoters	Wintering														-	-	-	-

FISH

Map ID	Subelement	Species	MD/VA						Monthly Presence											
			S	F	Concentration	J	F	M	A	M	J	J	A	S	N	D	Spawn	Eggs	Larvae	Juveniles
1575	Diadromous	Atlantic sturgeon	General Distribution	-/E	E												-	-	-	Jan-Dec
	Diadromous	Shortnose sturgeon	General Distribution	E/E	E												-	Apr-Jun	Apr-Oct	Jan-Dec
1583	Diadromous	Alewife	Nursery Area	-													-	-	-	Mar-May
	Diadromous	Atlantic sturgeon	General Distribution	-/E	E												-	-	-	Jan-Dec
	Diadromous	Blueback herring	Nursery Area	-													-	May-Jul	May-Oct	Feb-Jun
	Diadromous	Shortnose sturgeon	General Distribution	E/E	E												-	Jan-Dec	Jan-Dec	Jan-Dec
1631	Estuarine Nursery	White perch	Nursery Area	-													-	Apr-Jun	Jan-Dec	-
1664	Diadromous	Alewife	Migration	-													-	Aug-Nov	Aug-Nov	Mar-May
	Diadromous	Blueback herring	Migration	-													-	Aug-Nov	Feb-Jun	Aug-Nov
1666	Diadromous	American shad	Migration	-													-	Jun-Oct	Mar-Jun	Feb-Jun
	Diadromous	Hickory shad	Migration	-													-	-	-	Mar-Jun
	Diadromous	Striped bass	Migration	-													-	-	-	-

HABITATS & RARE PLANTS

Map ID	Subelement	Species	MD/VA						Monthly Presence											
			S	F	Concentration	J	F	M	A	M	J	J	A	S	N	D	Spawn	Eggs	Larvae	Juveniles
1727	Plant	Endangered plant	Vulnerable Occurrence	E/-	-											-	-	-	-	-

REPTILES & AMPHIBIANS

Map ID	Subelement	Species	MD/VA						Monthly Presence											
			S	F	Concentration	J	F	M	A	M	J	J	A	S	N	D	Nest	Hatch	Internest	Juveniles
1775	Turtle	K. ridley sea turtle	General Distribution	E/E	E	Uncommon	-									-	-	-	Nov-Dec	-
	Turtle	K. ridley sea turtle	General Distribution	E/E	E	Common	-									-	-	-	Apr-Oct	-
	Turtle	Loggerhead sea turtle	General Distribution	T/T	T	Common	-									-	-	-	Nov-Dec	Nov-Dec
	Turtle	Loggerhead sea turtle	General Distribution	T/T	T	Abundant	-									-	-	-	Apr-Oct	Apr-Oct
1781	Turtle	Diamondback terrapin	Nesting	-		Possible Nest	-									May-Aug	-	-	Apr-Oct	Apr-Oct

INVERTEBRATES

Map ID	Subelement	Species	MD/VA						Monthly Presence											
			S	F	Concentration	J	F	M	A	M	J	J	A	S	N	D	Spawn	Eggs	Larvae	Juveniles
1788	Crab	Blue crab	Highly Abundant-Summ	-												-	-	-	Jun-Oct	Jun-Oct

MARINE MAMMALS

<i>Map ID</i>	<i>Subelement</i>	<i>Species</i>	<i>Mapping Qualifier</i>	<i>MD/VA</i>	<i>S F</i>	<i>Concentration</i>	<i>J F M A M J J A S O N D</i>	<i>Mating</i>	<i>Calving</i>	<i>Pupping</i>	<i>Molt</i>
1844	Manatee	West Indian manatee	General Distribution	-IE	E	Rare					-

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

BIRDS

<i>Subelement</i>	<i>Species</i>	<i>Mapping Qualifier</i>	<i>MD/VA</i>	Monthly Presence												
				<i>J F M A M J J A S O N D</i>	<i>Nest</i>	<i>Mig.(S)</i>	<i>Mig.(F)</i>	<i>Molt</i>								
Waterfowl	American black duck	Nesting	High						Mar-Aug						Jul-Aug	
	American black duck	Wintering	1,000S													
	American coot	Wintering	10S													
	American wigeon	Wintering	100S													
	Canada goose	Nesting	Moderate-High						Mar-Jun							Jul-Aug
	Gadwall	Wintering	1,000S													
	Green-winged teal	Wintering	1,000S													
	Mallard	Wintering	1,000S													
	Northern pintail	Wintering	10S													
	Northern shoveler	Wintering	Present													

TERRESTRIAL MAMMALS


<i>Subelement</i>	<i>Species</i>	<i>Mapping Qualifier</i>	<i>MD/VA</i>	Monthly Presence												
				<i>J F M A M J J A S O N D</i>	<i>Nest</i>	<i>Mig.(S)</i>	<i>Mig.(F)</i>	<i>Molt</i>								
Small Mammal	Delmarva fox squirrel	General Distribution	E/E													

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)

BIRDS

<i>Subelement</i>	<i>Species</i>	<i>Mapping Qualifier</i>	<i>MD/VA</i>	Monthly Presence												
				<i>J F M A M J J A S O N D</i>	<i>Nest</i>	<i>Mig.(S)</i>	<i>Mig.(F)</i>	<i>Molt</i>								
Diving	Brown pelican															
	Brown pelican															
	D. crested cormorant															
	D. crested cormorant															
Gull/Tern	Gulls															
	Terns															
	Passerine	Saltmarsh sharp-tailed sparrow														
	Seaside sparrow															
Shorebird	Willet															
	Clapper rail															
Wading																

FISH



INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Bivalve	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Cephalopod	Squid	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
Crab	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Shrimp	Horseshoe crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Brown shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Mantis shrimp	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	White shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct

MARINE MAMMALS

Subelement	Species	Monthly Presence												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
Dolphin	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-
	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-
	Harbor porpoise	-	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES

ESI POLYGON HABITAT TYPES	Habitat Classification	Area (Sq. Miles)												% of ESI Shoreline
		J	F	M	A	M	J	J	A	S	O	N	D	
10A	Salt and Brackish Water Marshes	6,685.54												88%
10B	Freshwater Marshes	522.84												<1%
10C	Swamps	3,297.45												2%
10D	Scrub and Shrub Wetlands	610.90												2%
7	Exposed Tidal Flats	5.77												3%

ESI SHORELINE HABITAT TYPES	Shoreline Habitat Classification	Length (Meters)												% of ESI Shoreline
		J	F	M	A	M	J	J	A	S	O	N	D	
10A	Salt and Brackish Water Marshes	109,070.48												88%
10C	Swamps	281.85												<1%
10D	Scrub and Shrub Wetlands	2,795.58												2%
9A	Sheltered Tidal Flats	2,088.93												2%
9B	Vegetated Low Banks	3,245.84												3%
8B	Sheltered, Solid Man-Made Structures	13.39												<1%
8C	Sheltered Riprap	1,362.78												1%
7	Exposed Tidal Flats	911.41												1%
6B	Riprap	110.54												<1%
4	Coarse Grained Sand Beaches	4,009.08												3%

Total ESI Shoreline: 123,889.88
Total Shoreline: 111,151.13

Total ESI Shoreline:

Total Shoreline:


76.98

69.07

Note: A shoreline segment may include multiple habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 49D
Central Bay



SEE BACK OF MAP
for details about mapped species and
other species that occur in mapped area.
Data Published: September 2016

Map 49D Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	<i>Mapping Qualifier</i>						Concentration	<i>Monthly Presence</i>							
		S	F	MDVA	J	F	M	A	M	J	J	A	S	O	N	D
SAV	Submerged aquatic veg	High Ecological Value														
SAV	Submerged aquatic veg	High Ecological Value														
SAV	Submerged aquatic veg	High Ecological Value														

BIRDS

Map ID	Subelement	Species	<i>Mapping Qualifier</i>						Concentration	<i>Monthly Presence</i>							
			S	F	MDVA	J	F	M	A	M	J	J	A	S	O	N	D
84	Gull/Tern	Forster's tern	Nesting						5 Pairs								
120	Wading	Great blue heron	Nesting						5 Nests								
422	Raptor	Bald eagle	Nesting						Last Obs. 2003								
533	Watervowl	Bufflehead	Wintering						1,000S								
	Watervowl	Canada goose	Wintering						100S								
	Watervowl	Canvasback	Wintering						1,000S								
	Watervowl	Common goldeneye	Wintering						10S								
	Watervowl	Mergansers	Wintering						100S								
	Watervowl	Redhead	Wintering						1,000S								
	Watervowl	Ring-necked duck	Wintering						10S								
	Watervowl	Ruddy duck	Wintering						1,000S								
	Watervowl	Scaup	Wintering						100S								
	Watervowl	Snow goose	Wintering						1,000S								
	Watervowl	Tundra swan	Wintering						100S								
707	Raptor	Bald eagle	Roosting						-								
1169	Watervowl	American black duck	Nesting						Moderate-High								

FISH

Map ID	Subelement	Species	<i>Mapping Qualifier</i>						Concentration	<i>Monthly Presence</i>							
			S	F	MDVA	J	F	M	A	M	J	J	A	S	O	N	D
1656	Diadromous	American shad	Nursery Area						-								
	Diadromous	Hickory shad	Nursery Area						-								
	Diadromous	Striped bass	Nursery Area						-								

HABITATS & RARE PLANTS

Map ID	Subelement	Species	<i>Mapping Qualifier</i>						Concentration	<i>Monthly Presence</i>							
			S	F	MDVA	J	F	M	A	M	J	J	A	S	O	N	D
1727	Plant	Endangered plant	Vulnerable Occurrence	E-					-								

REPTILES & AMPHIBIANS

<i>Map ID</i>	<i>Subelement</i>	<i>Species</i>	<i>Mapping Qualifier</i>	<i>MD/V/A</i>	<i>S</i>	<i>F</i>	<i>Concentration</i>	<i>J F M A M J J A S O N D</i>	<i>Nest</i>	<i>Hatch</i>	<i>Internest</i>	<i>Juveniles</i>	<i>Adults</i>
1754	Turtle	Diamondback terrapin	Nesting	-							-		Apr-Oct
1781	Turtle	Diamondback terrapin	Nesting	-	Possible Nest						-		Apr-Oct
1782	Turtle	Diamondback terrapin	Nesting	-	Probable Nest						-		Apr-Oct

INVERTEBRATES

<i>Map ID</i>	<i>Subelement</i>	<i>Species</i>	<i>Mapping Qualifier</i>	<i>MD/V/A</i>	<i>S</i>	<i>F</i>	<i>Concentration</i>	<i>J F M A M J J A S O N D</i>	<i>Spawn</i>	<i>Eggs</i>	<i>Larvae</i>	<i>Juveniles</i>	<i>Adults</i>
1827	Insect	Federally threatened insect 1	Vulnerable Occurrence	E/T	T	-	-			Jun-Jul	Jan-Dec	-	Jun-Aug

MARINE MAMMALS

<i>Map ID</i>	<i>Subelement</i>	<i>Species</i>	<i>Mapping Qualifier</i>	<i>MD/V/A</i>	<i>S</i>	<i>F</i>	<i>Concentration</i>	<i>J F M A M J J A S O N D</i>	<i>Mating</i>	<i>Calving</i>	<i>Pupping</i>	<i>Molt</i>
1849	Whale	Humpback whale	General Distribution	E/E	E	Very Rare	-				-	-

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

BIRDS

<i>Subelement</i>	<i>Species</i>	<i>Mapping Qualifier</i>	<i>MD/V/A</i>	<i>S</i>	<i>F</i>	<i>Concentration</i>	<i>J F M A M J J A S O N D</i>	<i>Nest</i>	<i>Mig.(S)</i>	<i>Mig.(F)</i>	<i>Molt</i>
Diving	Common loon	Wintering	-								-
	D. crested cormorant	Migration	1,000S								-
	Grebes	Wintering	100S								-
	Red-throated loon	Wintering	100S								-
Pelagic	Northern gannet	Wintering	100S								-
Waterfowl	American black duck	Nesting	Moderate								-
	American black duck	Wintering	1,000S								-
	American coot	Wintering	10S								-
	American wigeon	Wintering	100S								-
	Canada goose	Nesting	Moderate-High								-
	Gadwall	Wintering	1,000S								-
	Green-winged teal	Wintering	1,000S								-
	Long-tailed duck	Wintering	100S								-
	Mallard	Nesting	Moderate-High								-
	Mallard	Wintering	1,000S								-
	Northern pintail	Wintering	10S								-
	Northern shoveler	Wintering	Present								-
	Scoters	Wintering	10,000S								-
	Wood duck	Nesting	Moderate-High								-

FISH

Subelement	Species	Mapping Qualifier				MD\VA				Monthly Presence											
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
Diadromous	Alewife	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Mar-May
	Alewife	Nursery Area	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Jun	Apr-Oct	Mar-Jun
	American shad	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Mar-Dec
	Atlantic sturgeon	General Distribution	-/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Feb-Jun
	Blueback herring	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Feb-Jun
	Blueback herring	Nursery Area	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Oct	Feb-Jun
	Hickory shad	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec
	Shortnose sturgeon	General Distribution	E/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Jun	Mar-Jun
	Striped bass	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

REPTILES & AMPHIBIANS

Subelement	Species	Mapping Qualifier				MD\VA				Monthly Presence										
		S	F	Concentration	J	F	M	J	J	A	S	O	N							
Turtle	K. ridley sea turtle	E/E	E	Uncommon	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	K. ridley sea turtle	E/E	E	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Loggerhead sea turtle	T/T	T	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Dec
	Loggerhead sea turtle	T/T	T	Abundant	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Dec

INVERTEBRATES

Subelement	Species	Mapping Qualifier				MD\VA				Monthly Presence			
		S	F	Concentration	J	F	M	J	J	A	S	O	N
Crab	Blue crab	Highly Abundant-Summ				-				-			

MARINE MAMMALS


Subelement	Species	Mapping Qualifier				MD\VA				Monthly Presence			
		S	F	Concentration	J	F	M	J	J	A	S	O	N
Manatee	West Indian manatee	-/E	E	Rare	-	-	-	-	-	-	-	-	-

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)


BIRDS

Subelement	Species	Mapping Qualifier				MD\VA				Monthly Presence			
		S	F	Concentration	J	F	M	J	J	A	S	O	N
Diving	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-
	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-
Gull/Tern	Gulls	-	-	-	-	-	-	-	-	-	-	-	-
	Terns	-	-	-	-	-	-	-	-	-	-	-	-
Passerine	Saltmarsh sharp-tailed sparrow	-	-	-	-	-	-	-	-	-	-	-	-
	Seaside sparrow	-	-	-	-	-	-	-	-	-	-	-	-
Shorebird	Willet	-	-	-	-	-	-	-	-	-	-	-	-
	Clapper rail	-	-	-	-	-	-	-	-	-	-	-	-
Wading	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH



INVERTEBRATES




MARINE MAMMALS



For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES




Total ESI Shoreline: 261,102.23 Total ESI Shoreline: 162.24
Total Shoreline: 200,430.03 Total Shoreline: 124.54

Note: A shoreline segment may include multiple habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 50
Central Bay



SEE BACK OF MAP
for details about mapped resources and
other resources that occur in mapped area.
Data Published: September 2016

Map 50 Central Bay

HUMAN USE RESOURCES

DISPLAYED ON MAP (POINTS)

<i>Map ID</i>	<i>Type</i>	<i>Name</i>	<i>Contact</i>	<i>Phone</i>
1	BOAT RAMP	DEAL ISLAND	SOMERSET COUNTY - ROADS DEPARTMENT	
2	BOAT RAMP	EWELL	SOMERSET COUNTY - ROADS DEPARTMENT	
3	BOAT RAMP	TYLERTON	SOMERSET COUNTY - ROADS DEPARTMENT	
4	BOAT RAMP	WENONA HARBOR PUBLIC BOAT RAMP	SOMERSET COUNTY - ROADS DEPARTMENT	
5	FERRY	EWELL FERRY	MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
6	REPEATED MEASUREMENT SITE	WQ - SHELLFISH HARVEST WATERS, CHESAPEAKE BAY	MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
7	REPEATED MEASUREMENT SITE	WQ - SHELLFISH HARVEST WATERS, MANOKIN RIVER	MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
8	REPEATED MEASUREMENT SITE	WQ - SHELLFISH HARVEST WATERS, TANGER SOUND	CHESAPEAKE BAY PROGRAM	800-968-7229
9	REPEATED MEASUREMENT SITE	WQ - STATION EE3.2		

DISPLAYED ON MAP (POLYGONS)

<i>Map ID</i>	<i>Type</i>	<i>Name</i>	<i>Contact</i>	<i>Phone</i>
10	MANAGEMENT AREA	DEAL ISLAND WMA	WILDLIFE AND HERITAGE SERVICE	
11	MANAGEMENT AREA	SOUTH MARSH ISLAND WMA	WILDLIFE AND HERITAGE SERVICE	
12	PARK	JANES ISLAND SP	MARYLAND PARK SERVICE	
13	WILDLIFE REFUGE	BLACKWATER NATIONAL WILDLIFE REFUGE	UNITED STATES FISH AND WILDLIFE SERVICE	410-228-2677
14	WILDLIFE REFUGE	MARTIN NATIONAL WILDLIFE REFUGE	UNITED STATES FISH AND WILDLIFE SERVICE	410-228-2692

ALSO PRESENT IN MAPPED AREA (POLYGONS)

<i>Type</i>	<i>Name</i>	<i>Contact</i>	<i>Phone</i>
STATE PROTECTED AREA	LOWER MAIN STEM BAY SANCTUARY	MARYLAND DEPARTMENT OF NATURAL RESOURCES	877-620-8367
STATE PROTECTED AREA	MANOKIN RIVER SANCTUARY	MARYLAND DEPARTMENT OF NATURAL RESOURCES	877-620-8367
STATE PROTECTED AREA	SOMERSET SANCTUARY	MARYLAND DEPARTMENT OF NATURAL RESOURCES	877-620-8367

JURISDICTIONS

COUNTY: DORCHESTER COUNTY, SOMERSET COUNTY
COAST GUARD: DISTRICT 5, SECTOR BALTIMORE
USACE: NORTH ATLANTIC DIVISION, BALTIMORE DISTRICT

SHORELINE RESOURCES**ESI POLYGON HABITAT TYPES**

ESI Rank	Habitat Classification	Area (Acres)	Area (Sq. Miles)
10A	Salt and Brackish Water Marshes	12,843.99	20.07
10B	Freshwater Marshes	2.36	0.00
10C	Swamps	34.27	0.05
10D	Scrub and Shrub Wetlands	118.11	0.18
9A	Sheltered Tidal Flats	1.10	0.00
7	Exposed Tidal Flats	88.52	0.14


ESI SHORELINE HABITAT TYPES

ESI Rank	Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
10A	Salt and Brackish Water Marshes	477,996.56	297.01	72%
10C	Swamps	338.62	0.21	< 1%
10D	Scrub and Shrub Wetlands	9,996.67	6.21	2%
9A	Sheltered Tidal Flats	6,559.80	4.08	1%
9B	Vegetated Low Banks	4,709.95	2.93	1%
8B	Sheltered, Solid Man-Made Structures	5,759.02	3.58	1%
8C	Sheltered Riprap	1,920.79	1.19	< 1%
7	Exposed Tidal Flats	120,721.03	75.01	18%
6B	Riprap	5,717.28	3.55	1%
5	Mixed Sand and Gravel Beaches	58.76	0.04	< 1%
4	Coarse Grained Sand Beaches	10,493.42	6.52	2%
3A	Fine to Medium Grained Sand Beaches	20,206.68	12.56	3%
1B	Exposed, Solid Man-Made Structures	638.63	0.40	< 1%
Total ESI Shoreline:		665,117.19	Total ESI Shoreline:	413.28
Total Shoreline:		490,708.62	Total Shoreline:	304.91

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%.

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 50A Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	Mapping Qualifier						Concentration	Monthly Presence							
		S	F	MD\VA	J	F	M	A	M	J	J	A	S	O	N	D
SAV	Submerged aquatic veg	High Ecological Value														
SAV	Submerged aquatic veg	High Ecological Value														
SAV	Submerged aquatic veg	High Ecological Value														

BIRDS

Map ID	Subelement	Species	Mapping Qualifier						Concentration	Monthly Presence							
			S	F	MD\VA	J	F	M	A	M	J	J	A	S	O	N	D
92	Gull/Tern	Forster's tern	Nesting			188 Pairs											
167	Wading	Great blue heron	Nesting			8 Nests											
351	Gull/Tern	Forster's tern	Nesting			3 Pairs											
414	Watertowl	Brant	Wintering			10S											
	Watertowl	Canvasback	Wintering			100S											
	Watertowl	Common goldeneye	Wintering			100S											
	Watertowl	Redhead	Wintering			10,000S											
	Watertowl	Ruddy duck	Wintering			100S											
	Watertowl	Scaup	Wintering			1,000S											
	Watertowl	Tundra swan	Wintering			1,000S											
1302	Diving	Common loon	Wintering			100S											

REPTILES & AMPHIBIANS

Map ID	Subelement	Species	Mapping Qualifier						Concentration	Monthly Presence							
			S	F	MD\VA	J	F	M	A	M	J	J	A	S	O	N	D
1781	Turtle	Diamondback terrapin	Nesting														

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

Subelement	Species	Mapping Qualifier						Concentration	Monthly Presence							
		S	F	MD\VA	J	F	M	A	M	J	J	A	S	O	N	D
Diving	Common loon	Wintering			100S											
	D. crested cormorant	Migration			1,000S											
	Grebes	Wintering			100S											
	Red-throated loon	Wintering			100S											
Pelagic	Northern gannet	Wintering			100S											
Waterfowl	American black duck	Nesting			Moderate-High											
	American black duck	Wintering			1,000S											
	American coot	Wintering			10S											

BIRDS (continued)

Subelement	Species	MDVA			Monthly Presence												Nest	Mig.(S)	Mig.(F)	Molt
		Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D			
American wigeon	Wintering	100S				-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Bufflehead	Wintering	1,000S				-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Canada goose	Nesting	Moderate-High				-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Aug
Canada goose	Wintering	100S				-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Canvasback	Wintering	1,000S				-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Common goldeneye	Wintering	10S				-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Gadwall	Wintering	1,000S				-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Green-winged teal	Wintering	1,000S				-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Long-tailed duck	Wintering	100S				-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Mallard	Wintering	100S				-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Mergansers	Wintering	10S				-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Northern pintail	Wintering	Present				-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Northern shoveler	Wintering	1,000S				-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Redhead	Wintering	10S				-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Ring-necked duck	Wintering	1,000S				-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Ruddy duck	Wintering	100S				-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Scaup	Wintering	10,000S				-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Scoters	Wintering	1,000S				-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Snow goose	Wintering	100S				-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Tundra swan	Wintering	Moderate-High				-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Wood duck	Nesting	Mar-Jun				-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Aug

FISH

Subelement	Species	MDVA			Monthly Presence												Eggs	Larvae	Juveniles	Adults	
		Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D				
Diadromous	Alewife	Migration	-			-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Mar-May	Mar-May	
	Alewife	Nursery Area	-			-	-	-	-	-	-	-	-	-	-	-	-	Apr-Jun	Apr-Oct	Mar-May	
	American shad	Migration	-			-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Mar-Jun	Mar-Jun	
	American shad	Nursery Area	-			-	-	-	-	-	-	-	-	-	-	-	-	May-Jul	May-Oct	Jan-Dec	
	Atlantic sturgeon	General Distribution	-IE	E	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Feb-Jun	Feb-Jun	
	Blueback herring	Migration	-			-	-	-	-	-	-	-	-	-	-	-	-	May-Jul	May-Oct	Feb-Jun	
	Blueback herring	Nursery Area	-			-	-	-	-	-	-	-	-	-	-	-	-	Jun-Oct	Feb-Jun	Feb-Jun	
	Hickory shad	Migration	-			-	-	-	-	-	-	-	-	-	-	-	-	Apr-Jun	Apr-Oct	Feb-Jun	
	Hickory shad	Nursery Area	-			-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec	Jan-Dec	
	Shortnose sturgeon	General Distribution	E/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	Striped bass	Migration	-			-	-	-	-	-	-	-	-	-	-	-	-	Apr-Jun	Jan-Dec	Mar-Jun	
	Striped bass	Nursery Area	-			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

REPTILES & AMPHIBIANS

Subelement	Species	MDVA			Monthly Presence												Nest	Hatch	Internest	Juveniles	Adults
		Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D				
Turtle	K. ridley sea turtle	General Distribution	E/E	E	Uncommon	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Dec	Apr-Oct	Apr-Oct	Apr-Oct
	K. ridley sea turtle	General Distribution	E/E	E	Common	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Dec	Nov-Dec	Nov-Dec	Nov-Dec
	Loggerhead sea turtle	General Distribution	T/T	T	Abundant	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Jun	Mar-Jun	Mar-Jun	Mar-Jun
	Loggerhead sea turtle	General Distribution	T/T	T	Common	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Jun	Jan-Dec	Jan-Dec	Jan-Dec

Central Bay: Map 50A

INVERTEBRATES

Subelement	Species	Mapping Qualifier	Monthly Presence																		
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawning	Eggs	Larvae	Juveniles
Crab	Blue crab	Highly Abundant-Summer	-/E	E	Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Oct

MARINE MAMMALS

Subelement	Species	Mapping Qualifier	Monthly Presence																	
			J	F	M	A	M	J	J	A	S	O	N	D	Spawning	Eggs	Larvae	Juveniles	Adults	
Manatee	West Indian manatee	General Distribution	-/E	E	Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Whale	Humpback whale	General Distribution	E/E	E	Very Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)

BIRDS

Subelement	Species	Monthly Presence																			
		J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Eggs	Larvae	Juveniles	Adults	
Diving	Brown pelican	-/E	E	Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Gull/Tern	Brown pelican	-/E	E	Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Gulls	Brown pelican	-/E	E	Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Terns	Brown pelican	-/E	E	Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Passerine	Saltmarsh sharp-tailed sparrow	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Pelagic	Seaside sparrow	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Shorebird	Storm-petrels	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Wading	Willet	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	Clapper rail	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

FISH

Subelement	Species	Monthly Presence																			
		J	F	M	A	M	J	J	A	S	O	N	D	Spawning	Eggs	Larvae	Juveniles	Adults			
Diadromous	Alewife	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-May	Apr-Jun	
	American eel	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-May	Jan-Dec	
	Blueback herring	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun	Feb-Jun	
	Gizzard shad	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec	
	Striped bass	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec	
	Estuarine Nursery	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	May-Nov	
	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec	
	Atlantic menhaden	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec	Apr-Dec	
	Black drum	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec	Apr-Dec	
	Black sea bass	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov	Apr-Dec	
	Bluefish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct	May-Oct	
	Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Oct	Mar-Nov	
	Red drum	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov	May-Nov	
	Spot	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Apr	Apr-Dec	
	Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Dec	Apr-Dec	
	Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Feb-May	Jan-Dec	
	Weakfish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Aug	Apr-Dec	

FISH (continued)

Subelement	Species	Mapping Qualifier	S	F	Concentration	Monthly Presence												Jan-Dec			
						J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs		
	White perch					-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec		
	Estuarine Resident	Silversides				-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov		
	Fish	Anchovies				-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec		
	Marine Benthic	Striped burrfish				-	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct		
	Marine Pelagic	Atlantic spadefish				-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep		
	Cownose ray	Cownose ray				-	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct		
	Northern searobin	Northern searobin				-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec		
	Spotted hake	Spotted hake				-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec		
	Butterfish	Butterfish				-	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Sep		
	Harvestfish	Harvestfish				-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov		
INVERTEBRATES																					
Subelement	Species	Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Monthly Presence	
	Bivalve	Eastern oyster				-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep		
	Crab	Hard clam				-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Oct		
	Crab	Softshell clam				-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep		
	Crab	Blue crab				-	-	-	-	-	-	-	-	-	-	-	-	-	May-Dec*		
	Crab	Horseshoe crab				-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec		
MARINE MAMMALS																					
Subelement	Species	Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping	Molt
	Dolphin	Bottlenose dolphin				-	-	-	-	-	-	-	-	-	-	-	-	-	May-Aug	May-Aug	-
	Dolphin	Bottlenose dolphin				-	-	-	-	-	-	-	-	-	-	-	-	-	May-Aug	-	-
	Dolphin	Harbor porpoise				-	-	-	-	-	-	-	-	-	-	-	-	-	May-Aug	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

ESI POLYGON HABITAT TYPES

ESI Rank	Habitat Classification	Area (Acres)	Area (Sq. Miles)
10A	Salt and Brackish Water Marshes	3,286.30	5.13
10B	Freshwater Marshes	2.36	0.00
10C	Swamps	30.72	0.05
10D	Scrub and Shrub Wetlands	27.98	0.04
9A	Sheltered Tidal Flats	1.10	0.00
7	Exposed Tidal Flats	6.70	0.01


ESI SHORELINE HABITAT TYPES

ESI Rank	Shoreline Habitat Classification	Length (Meters)	% of ESI Shoreline
10A	Salt and Brackish Water Marshes	93,576.77	77%
10C	Swamps	274.38	< 1%
10D	Scrub and Shrub Wetlands	6,061.96	5%
9A	Sheltered Tidal Flats	3,959.79	3%
9B	Vegetated Low Banks	3,704.44	3%
8B	Sheltered, Solid Man-Made Structures	1,484.88	1%
8C	Sheltered Riprap	1,815.47	2%
7	Exposed Tidal Flats	947.31	1%
6B	Riprap	3,083.22	3%
5	Mixed Sand and Gravel Beaches	58.76	< 1%
4	Coarse Grained Sand Beaches	5,497.75	5%
1B	Exposed, Solid Man-Made Structures	335.14	< 1%
Total ESI Shoreline:		120,799.86	Total ESI Shoreline: 75.06
Total Shoreline:		100,746.71	Total Shoreline: 62.60

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 50B Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	Mapping Qualifier						Concentration	Monthly Presence										
		S	F	M	D	N	V		J	F	M	A	M	J	J	A	S	O	N
SAV	Submerged aquatic veg	High Ecological Value						Ephemeral	-	-	-	-	-	-	-	-	-	-	-
SAV	Submerged aquatic veg	High Ecological Value						High	-	-	-	-	-	-	-	-	-	-	-
SAV	Submerged aquatic veg	High Ecological Value						Low	-	-	-	-	-	-	-	-	-	-	-

BIRDS

Map ID	Subelement	Species	Mapping Qualifier						Concentration	Monthly Presence										
			S	F	M	D	N	V		J	F	M	A	M	J	J	A	S	O	N
50	Diving	Brown pelican	Nesting						1594 Pairs	-	-	-	-	-	-	-	-	-	-	-
	Diving	D. crested cormorant	Nesting						557 Pairs	May-Aug	-	-	-	-	-	-	-	-	-	-
	Gull/Tern	Common tern	Nesting						46 Pairs	May-Aug	-	-	-	-	-	-	-	-	-	-
	Gull/Tern	Forster's tern	Nesting						427 Pairs	May-Jul	-	-	-	-	-	-	-	-	-	-
	Gull/Tern	G. black-backed gull	Nesting						27 Pairs	Apr-Jun	-	-	-	-	-	-	-	-	-	-
	Gull/Tern	Herring gull	Nesting						300 Pairs	Apr-Jul	-	-	-	-	-	-	-	-	-	-
	Wading	Great egret	Nesting						8 Pairs	Apr-Aug	-	-	-	-	-	-	-	-	-	-
87	Gull/Tern	Herring gull	Nesting						1 Pair	Apr-Jul	-	-	-	-	-	-	-	-	-	-
155	Wading	Great blue heron	Nesting						4 Nests	Feb-Jun	-	-	-	-	-	-	-	-	-	-
	Wading	Great egret	Nesting						30 Nests	Apr-Aug	-	-	-	-	-	-	-	-	-	-
469	Watertowl	American black duck	Nesting						High	Mar-Aug	-	-	-	-	-	-	-	-	-	-
	Watertowl	Canada goose	Nesting						Moderate-High	Mar-Jun	-	-	-	-	-	-	-	-	-	-
	Watertowl	Mallard	Nesting						Moderate-High	Mar-Aug	-	-	-	-	-	-	-	-	-	-
494	Diving	Brown pelican	Nesting						17 Pairs	May-Sep	-	-	-	-	-	-	-	-	-	-
	Diving	D. crested cormorant	Nesting						12 Pairs	May-Aug	-	-	-	-	-	-	-	-	-	-
	Wading	Great blue heron	Nesting						4 Pairs	Feb-Jun	-	-	-	-	-	-	-	-	-	-
550	Raptor	Peregrine falcon	Nesting						-T	Feb-Jun	-	-	-	-	-	-	-	-	-	-
1302	Diving	Common loon	Wintering						100S	-	-	-	-	-	-	-	-	-	-	-
1353	Diving	Brown pelican	Nesting						24 Pairs	May-Sep	-	-	-	-	-	-	-	-	-	-
	Diving	D. crested cormorant	Nesting						5 Pairs	May-Aug	-	-	-	-	-	-	-	-	-	-
	Gull/Tern	G. black-backed gull	Nesting						18 Pairs	Apr-Jun	-	-	-	-	-	-	-	-	-	-
	Gull/Tern	Herring gull	Nesting						36 Pairs	Apr-Jul	-	-	-	-	-	-	-	-	-	-

REPTILES & AMPHIBIANS

Map ID	Subelement	Species	Mapping Qualifier						Concentration	Monthly Presence										
			S	F	M	D	N	V		J	F	M	A	M	J	J	A	S	O	N
1781	Turtle	Diamondback terrapin	Nesting						Possible Nest	-	-	-	-	-	-	-	-	-	-	-
1782	Turtle	Diamondback terrapin	Nesting						Probable Nest	May-Aug	-	-	-	-	-	-	-	-	-	-

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

BIRDS

Subelement	Species	Mapping Qualifier	MD/VA						Monthly Presence												
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
Diving	Common loon	Wintering	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	D. crested cormorant	Migration	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-May	Aug-Nov	-
Grebes		Wintering	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Red-throated loon	Northern gannet	Wintering	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Pelagic	American black duck	Nesting	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Aug	-	
Waterfowl	American wigeon	Wintering	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Brant		Wintering	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Bufflehead		Wintering	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Canada goose		Wintering	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Canvasback		Wintering	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Common goldeneye		Wintering	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Gadwall		Wintering	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Green-winged teal		Wintering	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Long-tailed duck		Wintering	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Mallard		Wintering	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Mergansers		Wintering	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Northern pintail		Wintering	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Redhead		Wintering	10,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Ring-necked duck		Wintering	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Ruddy duck		Wintering	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Scaup		Wintering	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Scoters		Wintering	10,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Tundra swan		Wintering	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

FISH

Subelement	Species	Mapping Qualifier	MD/VA						Monthly Presence												
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles
Diadromous	Alewife	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-May
	American shad	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Jun
	Atlantic sturgeon	General Distribution	-IE	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Blueback herring	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Hickory shad	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Oct
	Shortnose sturgeon	General Distribution	E/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Striped bass	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Jun

REPTILES & AMPHIBIANS

		Monthly Presence																						
Subelement	Species	Mapping Qualifier		S	F	Concentration		J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles	Adults
Turtle	K. ridley sea turtle	General Distribution	E/E	E	Uncommon			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Dec	-
	K. ridley sea turtle	General Distribution	E/E	E	Common			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct	-
	Loggerhead sea turtle	General Distribution	T/T	T	Common			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Dec	Nov-Dec
	Loggerhead sea turtle	General Distribution	T/T	T	Abundant			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct	Apr-Oct

INVERTEBRATES

		Monthly Presence																						
Subelement	Species	Mapping Qualifier		S	F	Concentration		J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
Crab	Blue crab	Concentration Area				Highly Abundant-Summ		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Oct	Jun-Oct

MARINE MAMMALS

		Monthly Presence																						
Subelement	Species	Mapping Qualifier		S	F	Concentration		J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping	Molt	
Manatee	West Indian manatee	General Distribution	-E/E	E	Rare			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)

BIRDS

		Monthly Presence															
Subelement	Species	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
Diving	Brown pelican		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Brown pelican		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Brown pelican		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	D. crested cormorant		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Gull/Tern	Gulls		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Terns		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Pelagic	Storm-petrels		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

		Monthly Presence																
Subelement	Species	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
Diadromous	Alewife		-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-May	Apr-Jun
	American eel		-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-May	Jan-Dec
	Blueback herring		-	-	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun	Jan-Dec
	Gizzard shad		-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec
	Striped bass		-	-	-	-	-	-	-	-	-	-	-	-	-	-	Dec-Apr	Jan-Dec
Estuarine Nursery	Atlantic croaker		-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov	Jan-Dec
	Atlantic menhaden		-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec	Apr-Dec
	Black drum		-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov	Apr-Nov
	Black sea bass		-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct	May-Oct
	Bluefish		-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Oct	Mar-Oct
	Northern puffer		-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov	May-Nov
	Red drum		-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Apr	Apr-Dec
	Spot		-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Apr	Apr-Dec

FISH (continued)

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Dec
	Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Weakfish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	White perch	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Silversides	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Estuarine Resident	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Fish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Anchoovies	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Striped burrfish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Marine Benthic	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Atlantic spadefish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Sep
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	-

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Hard clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Horseshoe crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec

MARINE MAMMALS

Subelement	Species	Monthly Presence												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-
	Harbor porpoise	-	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov


SHORELINE RESOURCES

ESI POLYGON HABITAT TYPES		Habitat Classification	Area (Acres)	Area (Sq. Miles)
		Salt and Brackish Water Marshes	4,046.85	6.32
		Swamps	3.55	0.01
		Exposed Tidal Flats	26.67	0.04
ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification	Length (Meters)	Length (Miles)
		Salt and Brackish Water Marshes	141,147.31	87.71
		Swamps	64.23	0.04
		Exposed Tidal Flats	87,742.19	54.52
		Coarse Grained Sand Beaches	2,513.77	1.56
		Fine to Medium Grained Sand Beaches	6,803.41	4.23
		Total ESI Shoreline:	238,270.91	148.05
		Total Shoreline:	141,523.47	87.94
		Total ESI Shoreline:	141,523.47	87.94

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 50C Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	Mapping Qualifier			S	F	Concentration	Monthly Presence										
		High Ecological Value	High Ecological Value	High Ecological Value				J	F	M	A	M	J	J	A	S	O	N
SAV	Submerged aquatic veg							-	-	-	-	-	-	-	-	-	-	-
SAV	Submerged aquatic veg							-	-	-	-	-	-	-	-	-	-	-
SAV	Submerged aquatic veg							-	-	-	-	-	-	-	-	-	-	-

BIRDS

Map ID	Subelement	Species	Mapping Qualifier			S	F	Concentration	Monthly Presence										
			J	F	M				J	F	M	A	M	J	J	A	S	O	N
335	Gull/Tern	G. black-backed gull	Nesting					13 Pairs											
380	Gull/Tern	G. black-backed gull	Nesting					15 Pairs											
442	Watertowl	American black duck	Nesting					Moderate											
	Watertowl	Canada goose	Nesting					Moderate-High											
	Watertowl	Mallard	Nesting					Moderate-High											
	Watertowl	Wood duck	Nesting					Moderate-High											
483	Watertowl	American black duck	Nesting					Very High											
	Watertowl	Mallard	Nesting					Moderate-High											
515	Wading	BC night-heron	Nesting					12 Pairs											
	Wading	Glossy ibis	Nesting					1 Pair											
	Wading	Little blue heron	Nesting					1 Pair											
	Wading	YC night-heron	Nesting					2 Pairs											
550	Raptor	Peregrine falcon	Nesting					-T											
557	Watertowl	American black duck	Wintering					1,000S											
	Watertowl	American coot	Wintering					10S											
	Watertowl	American wigeon	Wintering					100S											
	Watertowl	Gadwall	Wintering					1,000S											
	Watertowl	Green-winged teal	Wintering					1,000S											
	Watertowl	Mallard	Wintering					1,000S											
	Watertowl	Northern pintail	Wintering					10S											
	Watertowl	Northern shoveler	Wintering					Present											
565	Watertowl	American black duck	Wintering					1,000S											
	Watertowl	American wigeon	Wintering					100S											
	Watertowl	Gadwall	Wintering					100S											
	Watertowl	Green-winged teal	Wintering					100S											
	Watertowl	Mallard	Wintering					100S											
	Watertowl	Northern pintail	Wintering					100S											
839	Watertowl	Canada goose	Wintering					100S											
860	Watertowl	Canada goose	Wintering					100S											
	Watertowl	Canvasback	Wintering					1,000S											

BIRDS (continued)

Map ID	Subelement	Species	MD/VA			Monthly Presence																	
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt	
	Watertowl	Redhead	Wintering			1,000S													-	-	-	-	
	Watertowl	Ruddy duck	Wintering			1,000S													-	-	-	-	
	Watertowl	Scaup	Wintering			100S													-	-	-	-	
	Watertowl	Snow goose	Wintering			1,000S													-	-	-	-	
	Watertowl	Tundra swan	Wintering			100S													-	-	-	-	
1148	Watertowl	Brant	Wintering			10S													-	-	-	-	
	Watertowl	Common goldeneye	Wintering			100S													-	-	-	-	
	Watertowl	Ruddy duck	Wintering			100S													-	-	-	-	
1315	Gull/Tern	G. black-backed gull	Nesting			53 Pairs													Apr-Jun	-	-	-	
	Gull/Tern	Herring gull	Nesting			26 Pairs													Apr-Jul	-	-	-	

FISH

Map ID	Subelement	Species	MD/VA			Monthly Presence																	
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Eggs	Larvae	Juveniles	Adults	
1681	Diadromous	Alewife	Nursery Area			-													-	Apr-Jun	Apr-Oct	Mar-May	
	Diadromous	Blueback herring	Nursery Area			-													-	May-Jul	May-Oct	Feb-Jun	

HABITATS & RARE PLANTS

Map ID	Subelement	Species	MD/VA			Monthly Presence																	
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Eggs	Larvae	Juveniles	Adults	
1727	Plant	Endangered plant	Vulnerable Occurrence	E/-.		-													-	-	-	-	
						-												-	-	-	-		

INVERTEBRATES

Map ID	Subelement	Species	MD/VA			Monthly Presence																	
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Eggs	Larvae	Juveniles	Adults	
1827	Insect	Federally threatened insect 1	Vulnerable Occurrence	E/T	T	-													Jun-Aug	Jun-Jul	Jan-Dec	-	
						-												-	-	-	-		

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

BIRDS

Subelement	Species	MD/VA			Monthly Presence																	
		Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt	
★ Diving	Common loon	Wintering			100S													-	-	Mar-May	Aug-Nov	-
	D. crested cormorant	Migration			1,000S													-	-	-	-	-
	Grebies	Wintering			100S													-	-	-	-	-
	Red-throated loon	Wintering			100S													-	-	-	-	-
	Northern gannet	Wintering			100S													-	-	-	-	-
	Brant	Wintering			100S													-	-	-	-	-
	Bufflehead	Wintering			1,000S													-	-	-	-	-
	Canada goose	Wintering			1,000S													-	-	-	-	-
	Canvasback	Wintering			100S													-	-	-	-	-
	Common goldeneye	Wintering			10S													-	-	-	-	-
	Long-tailed duck	Wintering			100S													-	-	-	-	-
	Mergansers	Wintering			100S													-	-	-	-	-

BIRDS (continued)

Map ID	Subelement	Species	MD/VA		Monthly Presence												Molt			
			Mapping Qualifier	S F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)
	Redhead	Wintering			10,000S												-	-	-	-
	Ring-necked duck	Wintering			10S												-	-	-	-
	Ruddy duck	Wintering			10S												-	-	-	-
	Scaup	Wintering			1,000S												-	-	-	-
	Scooter	Wintering			10,000S												-	-	-	-
	Tundra swan	Wintering			1,000S												-	-	-	-

FISH

Subelement	Species	MD/VA		Monthly Presence												Molt					
		Mapping Qualifier	S F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
	Diadromous	Alewife	Migration	-												-	-	-	-	Aug-Nov	Mar-May
		American shad	Migration	-												-	-	-	-	Aug-Nov	Mar-Jun
		Atlantic sturgeon	General Distribution	-E	E											-	-	-	-	-	Jan-Dec
		Blueback herring	Migration	-												-	-	-	-	Aug-Nov	Feb-Jun
		Hickory shad	Migration	-												-	-	-	-	Jun-Oct	Feb-Jun
		Shortnose sturgeon	General Distribution	E/E	E											-	-	-	-	Jan-Dec	Jan-Dec
		Striped bass	Migration	-												-	-	-	-	-	Mar-Jun

REPTILES & AMPHIBIANS

Subelement	Species	MD/VA		Monthly Presence												Molt					
		Mapping Qualifier	S F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Hatch	Internest	Juveniles
	Turtle	K. ridley sea turtle	General Distribution	E/E	E	Uncommon										-	-	-	-	Nov-Dec	-
		K. ridley sea turtle	General Distribution	E/E	E	Common										-	-	-	-	Apr-Oct	-
		Loggerhead sea turtle	General Distribution	T/T	T	Common										-	-	-	-	Nov-Dec	Nov-Dec
		Loggerhead sea turtle	General Distribution	T/T	T	Abundant										-	-	-	-	Apr-Oct	Apr-Oct

INVERTEBRATES

Subelement	Species	MD/VA		Monthly Presence												Molt					
		Mapping Qualifier	S F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
	Crab	Blue crab	Concentration Area	Highly Abundant-Summ												-	-	-	-	Jun-Oct	Jun-Oct

MARINE MAMMALS

Subelement	Species	MD/VA		Monthly Presence												Molt				
		Mapping Qualifier	S F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping	Molt
	Manatee	General Distribution	-E	E	Rare											-	-	-	-	-
	Whale	General Distribution	E/E	E	Very Rare											-	-	-	-	-

BIRDS

Sublement	Species	Monthly Presence												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
Diving	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-	-
Gull/Tern	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-	-
Gulls	Gulls	-	-	-	-	-	-	-	-	-	-	-	-	-
Terns	Terns	-	-	-	-	-	-	-	-	-	-	-	-	-
Passerine	Seaside sparrow	-	-	-	-	-	-	-	-	-	-	-	-	-
Shorebird	Willet	-	-	-	-	-	-	-	-	-	-	-	-	-
Wading	Clapper rail	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Sublement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Diadromous	Alewife	-	-	-	-	-	-	-	-	-	-	-	-	Mar-May
	American eel	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Jun
	Blueback herring	-	-	-	-	-	-	-	-	-	-	-	-	Jan-May
	Gizzard shad	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Striped bass	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Estuarine Nursery	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Atlantic menhaden	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Black drum	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Black sea bass	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Bluefish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Red drum	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Spot	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Dec
	Weakfish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	White perch	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Estuarine Resident	Silversides	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Fish	Anchovies	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Striped burrfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
Marine Benthic	Atlantic spadefish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Marine Pelagic	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Sep
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
		-	-	-	-	-	-	-	-	-	-	-	-	May-Nov

INVERTEBRATES

		Monthly Presence															
		J	F	M	A	M	J	J	A	S	O	N	D	Eggs	Larvae	Juveniles	Adults
Subelement	Species													Jun-Sep	Jun-Oct	Jan-Dec	Jan-Dec
Bivalve	Eastern oyster	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	Jan-Dec
	Hard clam													Jun-Sep	Jun-Oct	Jan-Dec	Jan-Dec
	Softshell clam													May-Dec*	-	Jan-Dec	Jan-Dec
Crab	Blue crab													-	-	Jan-Dec	Jan-Dec
	Horseshoe crab													-	-	Jan-Dec	Jan-Dec

MARINE MAMMALS

		Monthly Presence															
		J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping	Molt
Subelement	Species													May-Aug	May-Aug	-	-
	Bottlenose dolphin													-	-	-	-
	Harbor porpoise													-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES

ESI POLYGON HABITAT TYPES

ESI Rank	Habitat Classification
10A	Salt and Brackish Water Marshes

ESI SHORELINE HABITAT TYPES


ESI Rank	Shoreline Habitat Classification	Length (Meters)	% of ESI Shoreline
10A	Salt and Brackish Water Marshes	24,417.38	70%
7	Exposed Tidal Flats	7,819.91	22%
6B	Riprap	70.18	<1%
4	Coarse Grained Sand Beaches	2,481.90	1.54

Total ESI Shoreline: 34,789.38 Total ESI Shoreline: 21.62
Total Shoreline: 24,417.38 Total Shoreline: 15.17

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 50D Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	Mapping Qualifier						Concentration	Monthly Presence						
		S	F	MDVA	J	F	M	A	M	J	J	A	S	O	N
SAV	Submerged aquatic veg	High Ecological Value													
SAV	Submerged aquatic veg	High Ecological Value													
SAV	Submerged aquatic veg	High Ecological Value													

BIRDS

Map ID	Subelement	Species	Mapping Qualifier						Concentration	Monthly Presence					
			S	F	MDVA	J	F	M	A	M	J	J	A	S	O
128	Wading	Great blue heron	Nesting						3 Nests						
	Wading	Great egret	Nesting						18 Nests						
169	Wading	Great blue heron	Nesting						15 Nests						
170	Wading	Great blue heron	Nesting						15 Nests						
	Wading	Great egret	Nesting						45 Nests						
183	Wading	Great blue heron	Nesting						9 Nests						
195	Wading	Great blue heron	Nesting						48 Nests						
199	Wading	Great egret	Nesting						8 Nests						
202	Wading	Great blue heron	Nesting						1 Nest						
	Wading	Great egret	Nesting						5 Nests						
203	Wading	Great egret	Nesting						15 Nests						
204	Wading	Great egret	Nesting						140 Nests						
366	Gull/Tern	G. black-backed gull	Nesting						25 Pairs						
	Gull/Tern	Herring gull	Nesting						55 Pairs						
418	Raptor	Bald eagle	Nesting						Last Obs. 2004						
495	Wading	Great blue heron	Nesting						3 Pairs						
515	Wading	BC night-heron	Nesting						12 Pairs						
	Wading	Glossy ibis	Nesting						1 Pair						
	Wading	Little blue heron	Nesting						1 Pair						
	Wading	YC night-heron	Nesting						2 Pairs						
550	Raptor	Peregrine falcon	Nesting						-T						
866	Watertowl	Brant	Wintering						10S						
	Watertowl	Canada goose	Wintering						100S						
	Watertowl	Common goldeneye	Wintering						100S						
	Watertowl	Ruddy duck	Wintering						100S						
1294	Shorebird	American oystercatcher	Nesting						26 Nests						
1295	Shorebird	American oystercatcher	Nesting						10 Nests						
1296	Wading	Great blue heron	Nesting						1 Pair						
1297	Gull/Tern	G. black-backed gull	Nesting						32 Pairs						
	Gull/Tern	Herring gull	Nesting						637 Pairs						

BIRDS (continued)

Map ID	Subelement	Species	MD/VA				Monthly Presence															
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
	Wading	BC night-heron	Nesting			15 Pairs													Apr-Aug	-	-	-
	Wading	Glossy ibis	Nesting			4 Pairs													Apr-Jul	-	-	-
	Wading	Great blue heron	Nesting			1 Pair													Feb-Jun	-	-	-
	Wading	Great egret	Nesting			2 Pairs													Apr-Aug	-	-	-
	Wading	Little blue heron	Nesting			8 Pairs													May-Jun	-	-	-
	Wading	Tricolored heron	Nesting			5 Pairs													Mar-Apr	-	-	-
	Wading	YC night-heron	Nesting			26 Pairs													Apr-Jul	-	-	-
1300	Gull/Tern	G. black-backed gull	Nesting			14 Pairs													Apr-Jun	-	-	-
	Gull/Tern	Herring gull	Nesting			148 Pairs													Apr-Jul	-	-	-
	Wading	BC night-heron	Nesting			34 Pairs													Apr-Aug	-	-	-
	Wading	Glossy ibis	Nesting			10 Pairs													Apr-Jul	-	-	-
	Wading	Great blue heron	Nesting			98 Pairs													Feb-Jun	-	-	-
	Wading	Great egret	Nesting			3 Pairs													Apr-Aug	-	-	-
	Wading	Little blue heron	Nesting			8 Pairs													May-Jun	-	-	-
	Wading	Snowy egret	Nesting			12 Pairs													Apr-Aug	-	-	-
	Wading	Tricolored heron	Nesting			10 Pairs													Mar-Apr	-	-	-
	Wading	YC night-heron	Nesting			17 Pairs													Apr-Jul	-	-	-
1315	Gull/Tern	G. black-backed gull	Nesting			53 Pairs													Apr-Jun	-	-	-
	Gull/Tern	Herring gull	Nesting			26 Pairs													Apr-Jul	-	-	-
1316	Wading	Great blue heron	Nesting			15 Pairs													Feb-Jun	-	-	-

REPTILES & AMPHIBIANS

Map ID	Subelement	Species	MD/VA				Monthly Presence															
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Internest	Juveniles	Adults
1781	Turtle	Diamondback terrapin	Nesting			Possible Nest												May-Aug	-	-	Apr-Oct	Apr-Oct
1782	Turtle	Diamondback terrapin	Nesting			Probable Nest												May-Aug	-	-	Apr-Oct	Apr-Oct

MARINE MAMMALS

Map ID	Subelement	Species	MD/VA				Monthly Presence															
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping	Molt
1849	Whale	Humpback whale	General Distribution	E/E	E	Very Rare													-	-	-	-
																			-	-	-	-

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

BIRDS

Subelement	Species	MD/VA				Monthly Presence															
		Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
Diving	Common loon	Wintering			100S													-	-	-	-
	Common loon	Wintering			100S													Mar-May	Aug-Nov	-	-
	D. crested cormorant	Migration			1,000S													Mar-May	Aug-Nov	-	-
	D. crested cormorant	Wintering			1,000S													-	-	-	-
	Grebes	Wintering			100S													-	-	-	-
	Grebes	Wintering			100S													-	-	-	-
	Red-throated loon	Wintering			100S													-	-	-	-

BIRDS (continued)

Map ID	Subelement	Species	MD\VA						Monthly Presence												
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
Pelagic		Northern gannet	Wintering			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Waterfowl		American black duck	Nesting			Very High												Mar-Aug	-	-	Jul-Aug
		American black duck	Wintering			1,000S												-	-	-	-
		American wigeon	Wintering			100S												-	-	-	-
		Brant	Wintering			100S												-	-	-	-
		Bufflehead	Wintering			1,000S												-	-	-	-
		Canada goose	Wintering			100S												-	-	-	-
		Canvasback	Wintering			10S												-	-	-	-
		Common goldeneye	Wintering			100S												-	-	-	-
		Gadwall	Wintering			100S												-	-	-	-
		Green-winged teal	Wintering			100S												-	-	-	-
		Long-tailed duck	Wintering			100S												-	-	-	-
		Mallard	Nesting			Moderate-High											Mar-Aug	-	-	Jul-Aug	-
		Mallard	Wintering			100S											-	-	-	-	-
		Mergansers	Wintering			100S											-	-	-	-	-
		Northern pintail	Wintering			100S											-	-	-	-	-
		Redhead	Wintering			10,000S											-	-	-	-	-
		Ring-necked duck	Wintering			10S											-	-	-	-	-
		Ruddy duck	Wintering			10S											-	-	-	-	-
		Scaup	Wintering			1,000S											-	-	-	-	-
		Scoter	Wintering			10,000S											-	-	-	-	-
		Tundra swan	Wintering			1,000S											-	-	-	-	-

FISH

Subelement	Species	MD\VA						Monthly Presence												
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Juveniles	Larvae	Adults
Diadromous	Alewife	Migration		-													-	Aug-Nov	Mar-May	-
	American shad	Migration		-													-	Aug-Nov	Mar-Jun	-
	Atlantic sturgeon	General Distribution	-E	E	-												-		Jan-Dec	-
	Blueback herring	Migration															-	Aug-Nov	Feb-Jun	-
	Hickory shad	Migration															-	Jun-Oct	Feb-Jun	-
	Shortnose sturgeon	General Distribution	E/E	E	-												-	Jan-Dec	Jan-Dec	-
	Striped bass	Migration															-	Mar-Jun	-	-

REPTILES & AMPHIBIANS

Subelement	Species	MD\VA						Monthly Presence											
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Juveniles	Internest
Turtle	K. ridley sea turtle	General Distribution	E/E	E	Uncommon	-										-	Nov-Dec	Apr-Oct	-
	K. ridley sea turtle	General Distribution	E/E	E	Common	-										-	Nov-Dec	Nov-Dec	-
	Loggerhead sea turtle	General Distribution	T/T	T	Common	-										-	Apr-Oct	Apr-Oct	-
	Loggerhead sea turtle	General Distribution	T/T	T	Abundant	-										-	Nov-Dec	Nov-Dec	-

INVERTEBRATES

Subelement	Species	Mapping Qualifier	MD\VA	S F	Concentration	J F M A M J J A S O N D	Monthly Presence
Crab	Blue crab	Concentration Area			Highly Abundant-Summer	-	Adults Jun-Oct
MARINE MAMMALS							
Subelement	Species	Mapping Qualifier	MD\VA	S F	Concentration	J F M A M J J A S O N D	Monthly Presence
Manatee	West Indian manatee	General Distribution		-/E	E	-	-
		Rare				-	-

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)

BIRDS

Subelement	Species	Mapping Qualifier	MD\VA	S F	Concentration	J F M A M J J A S O N D	Monthly Presence
Diving	Brown pelican					-	-
	Brown pelican					-	-
	Brown pelican					-	-
	D. crested cormorant					-	-
	D. crested cormorant					-	-
Gull/Tern	Gulls					-	-
	Terns					-	-
Pelagic	Storm-petrels					-	-

FISH

Subelement	Species	Mapping Qualifier	MD\VA	S F	Concentration	J F M A M J J A S O N D	Monthly Presence
Diadromous	Alewife					Mar-May	Eggs Apr-Jun
	American eel					-	Adults Mar-Oct
	Blueback herring					-	Jan-Dec
	Gizzard shad					-	Jan-Dec
	Striped bass					-	Feb-Jun
Estuarine Nursery	Atlantic croaker					-	Jan-Dec
	Atlantic menhaden					-	Jan-Dec
	Black drum					-	Jan-Dec
	Black sea bass					-	Jan-Dec
	Bluefish					-	Jan-Dec
	Northern puffer					-	Jan-Dec
	Red drum					-	Jan-Dec
	Spot					-	Jan-Dec
	Spotted seatrout					-	Jan-Dec
	Summer flounder					-	Jan-Dec
	Weakfish					-	Jan-Dec
	White perch					-	Jan-Dec
Estuarine Resident	Silversides					-	Jan-Dec
Fish	Anchoovies					-	Jan-Dec
	Striped burrfish					-	Jan-Dec

FISH (continued)

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Marine Benthic	Atlantic spadefish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Marine Pelagic	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Bivalve	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Hard clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec*
Crab	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Horseshoe crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec

MARINE MAMMALS

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Dolphin	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	May-Aug
	Harbor porpoise	-	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES


ESI POLYGON HABITAT TYPES		Habitat Classification	Area (Acres)	Area (Sq. Miles)
ESI Rank				
10A		Salt and Brackish Water Marshes	5,093.15	7.96
10D		Scrub and Shrub Wetlands	90.14	0.14
	7	Exposed Tidal Flats	55.15	0.09

ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
ESI Rank					
10A		Salt and Brackish Water Marshes	218,855.11	135.99	81%
10D		Scrub and Shrub Wetlands	3,934.71	2.44	1%
9A		Sheltered Tidal Flats	2,600.01	1.62	1%
9B		Vegetated Low Banks	1,005.51	0.62	< 1%
8B		Sheltered, Solid Man-Made Structures	4,274.13	2.66	2%
8C		Sheltered Riprap	105.32	0.07	< 1%
	7	Exposed Tidal Flats	24,211.61	15.04	9%
6B		Riprap	2,563.88	1.59	1%
3A		Fine to Medium Grained Sand Beaches	13,403.27	8.33	5%
1B		Exposed, Solid Man-Made Structures	303.49	0.19	< 1%
		Total ESI Shoreline:	271,257.04	Total ESI Shoreline:	168.55
		Total Shoreline:	224,021.06	Total Shoreline:	139.20

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%.

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 51
Central Bay



Map 51 Central Bay

HUMAN USE RESOURCES

DISPLAYED ON MAP (POINTS)

Map ID	Type	Name	Contact	Phone
1	AIRPORT	BONHAM		
2	AIRPORT	CHEERY FIELD		
3	AIRPORT	WEBSTER NOLF		
4	AIRPORT	WINGFIELD		
5	AQUACULTURE	OYSTER NON-COMM BAKER RF#01-2210	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
6	AQUACULTURE	OYSTER NON-COMM JEWELL, LLOYD M., JR. RF#00-0008	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
7	AQUACULTURE	OYSTER NON-COMM MCKENNEY RF#00-1951	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
8	AQUACULTURE	OYSTER NON-COMM SZENAS RF#SZENAS	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
9	AQUACULTURE	OYSTER NON-COMM TORBECK, KATHERINE M. RF#00-0095	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
10	AQUACULTURE	OYSTER NON-COMM TYBURSKI, ROBERT M. RF#98-1712	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
11	ARTIFICIAL REEF	POINT LOOKOUT		
12	BOAT RAMP	PINEY POINT RECREATION AREA	ST. MARY'S COUNTY - DEPT. OF REC. AND PARKS	
13	BOAT RAMP	POINT LOOKOUT STATE PARK - BOAT RAMPS	MARYLAND DNR - PARK SERVICE	
14	BOAT RAMP	POINT LOOKOUT STATE PARK - CAMP STORE/BOAT RENTAL	MARYLAND DNR - PARK SERVICE	
15	BOAT RAMP	ST. INIGOS LANDING	ST. MARY'S COUNTY - DEPT. OF REC. AND PARKS	
16	FERRY	POINT LOOKOUT STATE PARK FERRY		
17	REPEATED MEASUREMENT SITE	DB - MOORED BUOY, POTOMAC, MD	NATIONAL COASTAL DATA DEVELOPMENT CENTER	866-732-2382
18	REPEATED MEASUREMENT SITE	DB - NOS, 8635750 - LEWISETTA, VA	NATIONAL COASTAL DATA DEVELOPMENT CENTER	866-732-2382
19	REPEATED MEASUREMENT SITE	NWLON - LEWISETTA, POTOMAC RIVER	NATIONAL WATER LEVEL PROGRAM	301-713-2815
20	REPEATED MEASUREMENT SITE	WQ - SHELLFISH HARVEST WATERS, CHESAPEAKE BAY	MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
21	REPEATED MEASUREMENT SITE	WQ - SHELLFISH HARVEST WATERS, POTOMAC RIVER	MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
22	REPEATED MEASUREMENT SITE	WQ - SHELLFISH HARVEST WATERS, SMITH CREEK	MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
23	REPEATED MEASUREMENT SITE	WQ - SHELLFISH HARVEST WATERS, ST GEORGES CREEK	MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
24	REPEATED MEASUREMENT SITE	WQ - SHELLFISH HARVEST WATERS, ST JEROME CREEK	MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
25	REPEATED MEASUREMENT SITE	WQ - SHELLFISH HARVEST WATERS, ST MARYS RIVER	MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
26	REPEATED MEASUREMENT SITE	WQ - SHELLFISH HARVEST WATERS, TANNER CREEK	MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
27	REPEATED MEASUREMENT SITE	WQ - STATION LE2.3		
28	REPEATED MEASUREMENT SITE	WQ - STATION SMNT14	CHESAPEAKE BAY PROGRAM	800-968-7229
29	REPEATED MEASUREMENT SITE	WQ - STATION SMT06	CHESAPEAKE BAY PROGRAM	800-968-7229
30	REPEATED MEASUREMENT SITE	WQ - STATION SMT07	CHESAPEAKE BAY PROGRAM	800-968-7229
31	REPEATED MEASUREMENT SITE	WQ - STATION SMT08	CHESAPEAKE BAY PROGRAM	800-968-7229
32	REPEATED MEASUREMENT SITE	WQ - STATION SMT10	CHESAPEAKE BAY PROGRAM	800-968-7229
33	REPEATED MEASUREMENT SITE	WQ - STATION SMT10B	CHESAPEAKE BAY PROGRAM	800-968-7229
34	REPEATED MEASUREMENT SITE	WQ - STATION SMT11	CHESAPEAKE BAY PROGRAM	800-968-7229
35	REPEATED MEASUREMENT SITE	WQ - STATION SMT12	CHESAPEAKE BAY PROGRAM	800-968-7229
36	REPEATED MEASUREMENT SITE	WQ - STATION SMT13	CHESAPEAKE BAY PROGRAM	800-968-7229

DISPLAYED ON MAP (POLYGONS)			
Map ID	Type	Name	Contact
37	MILITARY	WEBSTER FLD	
38	PARK	POINT LOOKOUT SP	MARYLAND PARK SERVICE
ALSO PRESENT IN MAPPED AREA (POINTS)			
Type	Name	Contact	Phone
COAST GUARD	ANT POTOMAC	301-872-4036	
COAST GUARD	STA ST. INGOES	301-872-4344	
ALSO PRESENT IN MAPPED AREA (POLYGONS)			
Type	Name	Contact	Phone
ESSENTIAL HABITAT	RARE VULNERABLE SPECIES		
STATE PROTECTED AREA	POINT LOOKOUT SANCTUARY	MARYLAND DEPARTMENT OF NATURAL RESOURCES	877-620-8367
STATE PROTECTED AREA	ST INGOES SF	MARYLAND FOREST SERVICE	

JURISDICTIONS

COUNTY: NORTHUMBERLAND COUNTY, ST. MARY'S COUNTY
 COAST GUARD: DISTRICT 5, SECTOR BALTIMORE, SECTOR HAMPTON ROADS
 USACE: NORTH ATLANTIC DIVISION, BALTIMORE DISTRICT

SHORELINE RESOURCES**ESI POLYGON HABITAT TYPES**

ESI Rank	Habitat Classification	Area (Acres)	Area (Sq. Miles)
10A	Salt and Brackish Water Marshes	781.16	1.22
10B	Freshwater Marshes	22.78	0.04
10C	Swamps	965.22	1.51
10D	Scrub and Shrub Wetlands	71.77	0.11
7	Exposed Tidal Flats	33.71	0.05

ESI SHORELINE HABITAT TYPES


ESI Rank	Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
10A	Salt and Brackish Water Marshes	110,019.23	68.36	21%
10B	Freshwater Marshes	587.61	0.37	< 1%
10C	Swamps	10,633.09	6.61	2%
10D	Scrub and Shrub Wetlands	5,542.20	3.44	1%
9A	Sheltered Tidal Flats	609.04	0.38	< 1%
9B	Vegetated Low Banks	157,824.61	98.07	30%
8B	Sheltered, Solid Man-Made Structures	46,861.99	29.12	9%
8C	Sheltered Riprap	32,799.68	20.38	6%
7	Exposed Tidal Flats	71,755.27	44.59	14%
6B	Riprap	35,129.25	21.83	7%
5	Mixed Sand and Gravel Beaches	3,248.49	2.02	1%
4	Coarse Grained Sand Beaches	41,599.82	25.85	8%
3A	Fine to Medium Grained Sand Beaches	83.86	0.05	< 1%
3B	Scars and Steep Slopes (Sand)	2,246.20	1.40	< 1%
1B	Exposed, Solid Man-Made Structures	11,813.99	7.34	2%

Total ESI Shoreline: 530,754.34 Total ESI Shoreline: 329.80
 Total Shoreline: 390,812.69 Total Shoreline: 242.84

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%.

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 51A Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	Mapping Qualifier		S	F	Concentration	Monthly Presence								
		High Ecological Value	High Ecological Value				J	F	M	A	M	J	J	A	S
SAV	Submerged aquatic veg						-	-	-	-	-	-	-	-	-
SAV	Submerged aquatic veg						-	-	-	-	-	-	-	-	-

BIRDS

Map ID	Subelement	Species	Mapping Qualifier		S	F	Concentration	Monthly Presence							
			Nesting	Roosting				J	F	M	A	M	J	J	A
418	Raptor	Bald eagle	Nesting	Wintering	Last Obs.	2004	Low	-	-	-	-	-	-	-	-
422	Raptor	Bald eagle	Nesting	Wintering	Last Obs.	2003	Low	-	-	-	-	-	-	-	-
429	Raptor	Bald eagle	Nesting	Wintering	Last Obs.	2006	Low	-	-	-	-	-	-	-	-
490	Watervowl	American black duck	Nesting	Roosting	-	1,000S	1,000S	-	-	-	-	-	-	-	-
707	Raptor	Bald eagle	Roosting	Wintering	1,000S	1,000S	1,000S	-	-	-	-	-	-	-	-
970	Watervowl	Canada goose	Roosting	Wintering	1,000S	1,000S	1,000S	-	-	-	-	-	-	-	-
	Watervowl	Redhead	Roosting	Wintering	1,000S	1,000S	1,000S	-	-	-	-	-	-	-	-
	Watervowl	Ring-necked duck	Roosting	Wintering	1,000S	1,000S	1,000S	-	-	-	-	-	-	-	-
	Watervowl	Ruddy duck	Roosting	Wintering	1,000S	1,000S	1,000S	-	-	-	-	-	-	-	-
	Watervowl	Scaup	Roosting	Wintering	100S	100S	100S	-	-	-	-	-	-	-	-

FISH

Map ID	Subelement	Species	Mapping Qualifier		S	F	Concentration	Monthly Presence							
			Nursery Area	Moderate Run				J	F	M	A	M	J	J	A
1511	Diadromous	Alewife	Nursery Area	Moderate Run	-	-	-	-	-	-	-	-	-	-	-
	Diadromous	Blueback herring	Nursery Area	Moderate Run	-	-	-	-	-	-	-	-	-	-	-
	Estuarine Nursery	White perch	Spawning Area	Moderate Run	-	-	-	-	-	-	-	-	-	-	-
1674	Diadromous	Alewife	Nursery Area	Moderate Run	-	-	-	-	-	-	-	-	-	-	-
	Diadromous	Blueback herring	Nursery Area	Moderate Run	-	-	-	-	-	-	-	-	-	-	-

HABITATS & RARE PLANTS

Map ID	Subelement	Species	Mapping Qualifier		S	F	Concentration	Monthly Presence							
			Vulnerable Occurrence	Endangered plant				J	F	M	A	M	J	J	A
1738	Plant				E/-	3 Species Present		-	-	-	-	-	-	-	-
								-	-	-	-	-	-	-	-

REPTILES & AMPHIBIANS

Map ID	Subelement	Species	Mapping Qualifier		S	F	Concentration	Monthly Presence							
			Possible Nest	Probable Nest				J	F	M	A	M	J	J	A
1781	Turtle	Diamondback terrapin	Nesting	Nesting	-	-	-	-	-	-	-	-	-	-	-
1782	Turtle	Diamondback terrapin	Nesting	Nesting	-	-	-	-	-	-	-	-	-	-	-

MARINE MAMMALS

Map ID	Subelement	Species	Mapping Qualifier	MD/VA	MD/VA	Concentration	Monthly Presence
		Humpback whale	General Distribution	S F	E/E E	Very Rare	J F M A M J J A S O N D
WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)							

BIRDS

Subelement	Species	Mapping Qualifier	MD/VA	MD/VA	Concentration	Monthly Presence
WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)						
Diving	Common loon	Wintering			Present	-
	Common loon	Wintering			100S	-
	D. crested cormorant	Migration			1,000S	-
Grebes		Wintering			100S	-
Grebes		Wintering			100S	-
Red-throated loon		Wintering			Present	-
Red-throated loon		Wintering			100S	-
Pelagic	Northern gannet	Wintering			100S	-
	Northern gannet	Wintering			Present	-
Waterfowl	American black duck	Wintering			1,000S	-
	American black duck	Wintering			10S	-
	American coot	Wintering			10S	-
	American coot	Wintering			100S	-
	American wigeon	Wintering			10S	-
	Bufflehead	Wintering			1,000S	-
	Canada goose	Nesting			Moderate-High	-
	Canada goose	Wintering			10,000S	-
Canvasback		Wintering			1,000S	-
Common goldeneye		Wintering			100S	-
Gadwall		Wintering			10S	-
Green-winged teal		Wintering			100S	-
Long-tailed duck		Wintering			100S	-
Mallard		Nesting			Moderate-High	-
Mallard		Wintering			100S	-
Mallard		Wintering			1,000S	-
Mergansers		Wintering			100S	-
Northern pintail		Wintering			10S	-
Northern pintail		Wintering			100S	-
Redhead		Wintering			10S	-
Ring-necked duck		Wintering			10S	-
Ruddy duck		Wintering			10,000S	-
Scaup		Wintering			1,000S	-
Scoters		Wintering			100S	-
Snow goose		Wintering			10,000S	-
Tundra swan		Wintering			Present	-
Wood duck		Nesting			100S	-
					Moderate-High	-
					Mar-Jun	-
					Jul-Aug	-

FISH

Subelement	Species	MD\VA												Monthly Presence												
		S	F	Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults		
Diadromous	Alewife	Migration	Migration	Moderate	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Mar-May	
	Alewife	Migration	Migration	Run	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Mar-May
	American shad	Migration	Migration	Strong Run	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Mar-Jun
	American shad	General Distribution	-E	E	Moderate	Run	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Mar-Jun
	Atlantic sturgeon	Migration	Migration	Moderate	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Jan-Dec
	Blueback herring	Migration	Migration	Moderate	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun	Feb-Jun
	Blueback herring	Migration	Migration	Moderate	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun	Feb-Jun
	Hickory shad	Migration	Migration	Moderate	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Oct	Feb-Jun
	Hickory shad	General Distribution	E/E	E	Moderate	Run	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Oct	Feb-Jun
	Shortnose sturgeon	General Distribution	E/E	E	Moderate	Run	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec
	Shortnose sturgeon	Migration	Migration	Moderate	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Jun	Mar-Jun
	Striped bass	Migration	Migration	Moderate	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Jun	Mar-Jun
	Striped bass	Nursery Area	Nursery Area	Nursery Area	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Jun	Jan-Dec

REPTILES & AMPHIBIANS

Subelement	Species	MD\VA												Monthly Presence												
		S	F	Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Neat	Hatch	Internest	Juveniles	Adults		
Turtle	K. ridley sea turtle	General Distribution	E/E	E	Uncommon	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Dec	-
	K. ridley sea turtle	General Distribution	E/E	E	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct	-
	Loggerhead sea turtle	General Distribution	T/T	T	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Dec	Nov-Dec
	Loggerhead sea turtle	General Distribution	T/T	T	Abundant	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct	Apr-Oct

INVERTEBRATES

Subelement	Species	MD\VA												Monthly Presence												
		S	F	Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Neat	Hatch	Internest	Juveniles	Adults		
Crab	Blue crab	Concentration Area	Highly Abundant-Summ	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Oct	Jun-Oct

MARINE MAMMALS

Subelement	Species	MD\VA												Monthly Presence												
		S	F	Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Neat	Hatch	Internest	Juveniles	Adults		
Manatee	West Indian manatee	General Distribution	-E	E	Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-May	Aug-Nov

BIRDS

Subelement	Species	MD\VA												Monthly Presence											
		J	F	M	A	M	J	J	A	S	O	N	D	Neat	Mig.(S)	Mig.(F)	Molt								
Diving	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Gull/Tern	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Gulls	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Terns	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Pelagic	Storm-petrels	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

	Subelement	Species	Monthly Presence												Adults				
			J	F	M	A	M	J	J	A	S	O	N	D					
	Diadromous	Alewife	-	-	-	-	-	-	-	-	-	-	-	-	Mar-May	Apr-Jun	Larvae	Juveniles	Adults
		American eel	-	-	-	-	-	-	-	-	-	-	-	-	Jan-May	Apr-Oct	Mar-May	Jan-Dec	Mar-Dec
		Blueback herring	-	-	-	-	-	-	-	-	-	-	-	-	May-Jun	May-Jul	May-Oct	Feb-Jun	Jan-Dec
		Gizzard shad	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Jun	-	Jan-Dec	Jan-Dec	Jan-Dec
		Striped bass	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec	Jan-Dec
	Estuarine Nursery	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Jun	Apr-Jun	Jan-Dec	Jan-Dec	Jan-Dec
		Atlantic menhaden	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec	Apr-Dec	Apr-Dec
		Black drum	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec	Apr-Dec	Jan-Dec
		Black sea bass	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov	Apr-Nov	Apr-Nov
		Bluefish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct	May-Oct	May-Oct
		Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Oct	Mar-Oct	Mar-Oct
		Red drum	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov	May-Nov	May-Nov
		Spot	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Apr	Nov-Apr	Apr-Dec
		Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Dec	Aug-Dec	Apr-Dec
		Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Feb-May	Feb-May	Apr-Dec
		Weakfish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Aug	May-Aug	Jun-Dec
		White perch	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec	Jan-Dec
	Estuarine Resident	Silversides	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov	Mar-Nov	Mar-Nov
		Anchovies	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		Striped burrfish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct	May-Oct	May-Oct
		Atlantic spadefish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec	Jan-Dec
		Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep	Jun-Sep	Jun-Sep
		Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct	May-Oct	Apr-Dec
		Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec	Jan-Dec
	Marine Pelagic	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Sep	Jul-Sep	Mar-Nov
		Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov

INVERTEBRATES

	Subelement	Species	Monthly Presence												Adults				
			J	F	M	A	M	J	J	A	S	O	N	D					
	Bivalve	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep	Jun-Sep	Larvae	Juveniles	Adults
		Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Aug	May-Aug	Jan-Dec
		Horseshoe crab	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec

MARINE MAMMALS

	Subelement	Species	Monthly Presence												Adults				
			J	F	M	A	M	J	J	A	S	O	N	D					
	Dolphin	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	May-Aug	May-Aug	-	-	-
		Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		Harbor porpoise	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from [response.restoration.noaa.gov](http://restoration.noaa.gov)

SHORELINE RESOURCES

ESI POLYGON HABITAT TYPES		Habitat Classification		Area (Acres)	Area (Sq. Miles)
ESI Rank					
10A		Salt and Brackish Water Marshes		164.30	0.26
10B		Freshwater Marshes		9.67	0.02
10C		Swamps		341.23	0.53
10D		Scrub and Shrub Wetlands		11.54	0.02
7		Exposed Tidal Flats		23.79	0.04


ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification		Length (Meters)	Length (Miles)	% of ESI Shoreline
ESI Rank						
10A		Salt and Brackish Water Marshes		41,614.53	25.86	21%
10B		Freshwater Marshes		449.71	0.28	< 1%
10C		Swamps		4,535.74	2.82	2%
10D		Scrub and Shrub Wetlands		2,372.94	1.47	1%
9A		Sheltered Tidal Flats		422.63	0.26	< 1%
9B		Vegetated Low Banks		64,508.09	40.08	32%
8B		Sheltered, Solid Man-Made Structures		17,445.83	10.84	9%
8C		Sheltered Riprap		12,345.03	7.67	6%
7		Exposed Tidal Flats		30,788.27	19.13	15%
6B		Riprap		8,629.89	5.36	4%
5		Mixed Sand and Gravel Beaches		963.10	0.60	< 1%
4		Coarse Grained Sand Beaches		15,123.37	9.40	7%
3B		Scarps and Steep Slopes (Sand)		784.79	0.49	< 1%
1B		Exposed, Solid Man-Made Structures		2,629.45	1.63	1%

Total ESI Shoreline:	202,613.35	Total ESI Shoreline:	125.90
Total Shoreline:	143,469.25	Total Shoreline:	89.15

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 51B
Central Bay



SEE BACK OF MAP
for details about mapped species and
other species that occur in mapped area.
Data Published: September 2016



Map 51B Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

	Subelement	Species	Mapping Qualifier	MD/V/A	S	F	Concentration	J F M A M J J A S O N D	Monthly Presence
	SAV	Submerged aquatic veg	High Ecological Value	-	-	-	-	-	-
	SAV	Submerged aquatic veg	High Ecological Value	-	-	-	-	-	-
	SAV	Submerged aquatic veg	High Ecological Value	-	-	-	-	-	-

BIRDS

	Map ID	Subelement	Species	Mapping Qualifier	MD/V/A	S	F	Concentration	J F M A M J J A S O N D	Monthly Presence
	418	Raptor	Bald eagle	Nesting	Last Obs. 2004	-	-	-	Feb-Apr	-

BIRDS

	Subelement	Species	Mapping Qualifier	MD/V/A	S	F	Concentration	J F M A M J J A S O N D	Monthly Presence
	Diving	Common loon	Wintering	-	-	-	-	-	-
	Diving	Common loon	Wintering	-	-	-	-	-	-
	Diving	D. crested cormorant	Wintering	-	-	-	-	-	-
	Diving	Grebels	Wintering	-	-	-	-	-	-
	Diving	Grebels	Wintering	-	-	-	-	-	-
	Diving	Red-throated loon	Wintering	-	-	-	-	-	-
	Diving	Red-throated loon	Wintering	-	-	-	-	-	-
	Pelagic	Northern gannet	Wintering	-	-	-	-	-	-
	Pelagic	Northern gannet	Wintering	-	-	-	-	-	-
	Waterfowl	American black duck	Wintering	-	-	-	-	-	-
	Waterfowl	American coot	Wintering	-	-	-	-	-	-
	Waterfowl	American wigeon	Wintering	-	-	-	-	-	-
	Waterfowl	Bufflehead	Wintering	-	-	-	-	-	-
	Waterfowl	Canada goose	Wintering	-	-	-	-	-	-
	Waterfowl	Canvasback	Wintering	-	-	-	-	-	-
	Waterfowl	Common goldeneye	Wintering	-	-	-	-	-	-
	Waterfowl	Gadwall	Wintering	-	-	-	-	-	-
	Waterfowl	Long-tailed duck	Wintering	-	-	-	-	-	-
	Waterfowl	mallard	Wintering	-	-	-	-	-	-
	Waterfowl	Mergansers	Wintering	-	-	-	-	-	-
	Waterfowl	Northern pintail	Wintering	-	-	-	-	-	-
	Waterfowl	Redhead	Wintering	-	-	-	-	-	-
	Waterfowl	Ring-necked duck	Wintering	-	-	-	-	-	-

BIRDS (continued)

Map ID	Subelement	Species	MD\VA		Monthly Presence															
			Mapping Qualifier	S F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)
		Ruddy duck	Wintering		1,000S												-	-	-	-
		Scallop	Wintering		100S												-	-	-	-
		Scoters	Wintering		100S												-	-	-	-
		Tundra swan	Wintering		100S												-	-	-	-

FISH

Subelement	Species	MD\VA		Monthly Presence																	
		Mapping Qualifier	S F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
	Alewife	Migration		Moderate Run												-	-	-	-	Aug-Nov	Mar-May
	Alewife	Nursery Area	Migration	Moderate Run												-	-	Apr-Jun	Apr-Oct	Mar-May	
	American shad	General Distribution	-E E	Strong Run												-	-	-	-	Aug-Nov	Mar-Jun
	Atlantic sturgeon	Migration		Moderate Run												-	-	-	-	Jan-Dec	Feb-Jun
	Blueback herring	Migration		Moderate Run												-	-	May-Jul	May-Oct	Feb-Jun	
	Blueback herring	Nursery Area	Migration	Moderate Run												-	-	Jun-Oct	Feb-Jun	Jan-Dec	
	Hickory shad	General Distribution	E/E E	Moderate Run												-	-	-	-	Jan-Dec	Mar-Jun
	Shortnose sturgeon	Migration	E/E E	-												-	-	-	-	Jan-Dec	Apr-Oct
	Shortnose sturgeon	Nursery Area	-													-	-	-	-	Jan-Dec	Mar-May
	Striped bass	Spawning Area	-	-												-	-	Apr-Jun	Jan-Dec	-	Mar-May
	Estuarine Nursery	White perch	-	-												-	-	Apr-Jun	Apr-Jun	-	Mar-May

REPTILES & AMPHIBIANS

Subelement	Species	MD\VA		Monthly Presence																	
		Mapping Qualifier	S F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles	Adults
	K. ridley sea turtle	General Distribution	E/E E	Common												-	-	-	-	Apr-Oct	-
	K. ridley sea turtle	General Distribution	E/E E	Uncommon												-	-	-	-	Nov-Dec	-
	Loggerhead sea turtle	General Distribution	T/T T	Common												-	-	-	-	Nov-Dec	Apr-Oct
	Loggerhead sea turtle	General Distribution	T/T T	Abundant												-	-	-	-	Apr-Oct	Jan-Oct

INVERTEBRATES

Subelement	Species	MD\VA		Monthly Presence																	
		Mapping Qualifier	S F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
	Blue crab	Concentration Area	Highly Abundant-Summ													-	-	-	-	Jun-Oct	Jun-Oct

MARINE MAMMALS

Subelement	Species	MD\VA		Monthly Presence																	
		Mapping Qualifier	S F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping	Molt	
	West Indian manatee	General Distribution	-E E	Rare												-	-	-	-	Jun-Oct	Jun-Oct

BIRDS

Subelement	Species	<i>Monthly Presence</i>												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
Diving	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-	-
	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-	-
	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-	-
Gull/Tern	Gulls	-	-	-	-	-	-	-	-	-	-	-	-	-
	Terns	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Subelement	Species	<i>Monthly Presence</i>												Adults					
		J	F	M	A	M	J	J	A	S	O	N	D	Mar-May	Eggs	Larvae	Juveniles		
Diadromous	Alewife	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Jun	Apr-Oct	Mar-Dec	Jan-Dec		
	American eel	-	-	-	-	-	-	-	-	-	-	-	-	Jan-May	-	-	Jan-Dec		
	Blueback herring	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct	Feb-Jun	-	Jan-Dec		
	Gizzard shad	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec	-	Jan-Dec		
	Striped bass	-	-	-	-	-	-	-	-	-	-	-	-	Dec-Apr	Jan-Dec	May-Nov	May-Nov		
Estuarine Nursery	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec	Jan-Dec	
	Atlantic menhaden	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec	Apr-Dec	Apr-Dec	
	Black drum	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov	Apr-Nov	Apr-Nov	
	Black sea bass	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct	May-Oct	May-Oct	
	Bluefish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Oct	Mar-Oct	Mar-Oct	
	Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov	May-Nov	May-Nov	
	Red drum	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Apr	Nov-Apr	Nov-Apr	
	Spot	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Dec	Aug-Dec	Aug-Dec	
	Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Feb-May	Feb-May	Feb-May	
	Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Aug	May-Aug	May-Aug	
	Weakfish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Dec	Jun-Dec	Jun-Dec	
	White perch	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec	Jan-Dec	
Estuarine Resident	Silversides	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov	Mar-Nov	Mar-Nov	
	Anchoovies	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec	Jan-Dec
	Striped burrfish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct	May-Oct	May-Oct	
	Atlantic spadefish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep	Jun-Sep	Jun-Sep	
	Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct	May-Oct	May-Oct	
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec	Apr-Dec	Apr-Dec	
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec	Jan-Dec	
Marine Pelagic	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Sep	Jul-Sep	Jul-Sep	
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov	Mar-Nov	Mar-Nov	

INVERTEBRATES


Subelement	Species	<i>Monthly Presence</i>												Adults				
		J	F	M	A	M	J	J	A	S	O	N	D	Jun-Sep	Eggs	Larvae	Juveniles	
Bivalve	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec	Jan-Dec
Crab	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec	Jan-Dec
	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec	Jan-Dec
	Horseshoe crab	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec	Jan-Dec

MARINE MAMMALS

Subelement	Species	Monthly Presence
Dolphin	Bottlenose dolphin	J F M A M J J A S O N D
		-
		-
		-
		-
		-
		-
		-
		-
		-
		-
		-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov


SHORELINE RESOURCES



Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





SEE BACK OF MAP
for details about mapped species and
other species that occur in mapped area.
Data Published: September 2016

0 Not for Navigation 1 Miles
0 1 Kilometers
1:50,000



Map 51C Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BIRDS

Map ID	Subelement	Species	Mapping Qualifier	MD/VA	S	F	Concentration	Monthly Presence															
								J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
44	Wading	Great blue heron	Nesting	62 Pairs				-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun	Feb-Jun	-	-
185	Wading	Great blue heron	Nesting	56 Nests				-	-	-	-	-	-	-	-	-	-	-	-	Last Obs. 2004	Feb-Apr	-	-
418	Raptor	Bald eagle	Nesting	Moderate-High				-	-	-	-	-	-	-	-	-	-	-	-	Mar-Jun	Mar-Aug	Jul-Aug	Jul-Aug
452	Waterfowl	Canada goose	Nesting	Moderate-High				-	-	-	-	-	-	-	-	-	-	-	-	Mar-Aug	Mar-Aug	Jul-Aug	Jul-Aug
463	Waterfowl	Mallard	Nesting	Low				-	-	-	-	-	-	-	-	-	-	-	-	Mar-Aug	Mar-Aug	Jul-Aug	Jul-Aug
		American black duck	Nesting	Moderate-High				-	-	-	-	-	-	-	-	-	-	-	-	Mar-Jun	Mar-Aug	Jul-Aug	Jul-Aug
		Waterfowl	Canada goose	Moderate-High				-	-	-	-	-	-	-	-	-	-	-	-	Mar-Jun	Mar-Aug	Jul-Aug	Jul-Aug
		Waterfowl	Mallard	Moderate-High				-	-	-	-	-	-	-	-	-	-	-	-	Mar-Jun	Mar-Aug	Jul-Aug	Jul-Aug
484	Waterfowl	Wood duck	Nesting	Low				-	-	-	-	-	-	-	-	-	-	-	-	Mar-Aug	Mar-Aug	Jul-Aug	Jul-Aug
490	Waterfowl	American black duck	Nesting	Moderate-High				-	-	-	-	-	-	-	-	-	-	-	-	Mar-Aug	Mar-Aug	Jul-Aug	Jul-Aug
536	Waterfowl	Bufflehead	Wintering	1,000S				-	-	-	-	-	-	-	-	-	-	-	-	1,000S	1,000S	-	-
		Waterfowl	Canada goose	Wintering				-	-	-	-	-	-	-	-	-	-	-	-	1,000S	1,000S	-	-
		Waterfowl	Canvasback	Wintering				-	-	-	-	-	-	-	-	-	-	-	-	1,000S	1,000S	-	-
		Waterfowl	Common goldeneye	Wintering				-	-	-	-	-	-	-	-	-	-	-	-	1,000S	1,000S	-	-
		Waterfowl	Mergansers	Wintering				-	-	-	-	-	-	-	-	-	-	-	-	100S	100S	-	-
		Waterfowl	Redhead	Wintering				-	-	-	-	-	-	-	-	-	-	-	-	100S	100S	-	-
		Waterfowl	Ring-necked duck	Wintering				-	-	-	-	-	-	-	-	-	-	-	-	100S	100S	-	-
		Waterfowl	Ruddy duck	Wintering				-	-	-	-	-	-	-	-	-	-	-	-	1,000S	1,000S	-	-
		Waterfowl	Scaup	Wintering				-	-	-	-	-	-	-	-	-	-	-	-	100S	100S	-	-
		Waterfowl	Tundra swan	Wintering				-	-	-	-	-	-	-	-	-	-	-	-	100S	100S	-	-
562	Waterfowl	American black duck	Wintering	1,000S				-	-	-	-	-	-	-	-	-	-	-	-	1,000S	1,000S	-	-
		Waterfowl	American coot	Wintering				-	-	-	-	-	-	-	-	-	-	-	-	10S	10S	-	-
		Waterfowl	Green-winged teal	Wintering				-	-	-	-	-	-	-	-	-	-	-	-	100S	100S	-	-
		Waterfowl	Mallard	Wintering				-	-	-	-	-	-	-	-	-	-	-	-	1,000S	1,000S	-	-
		Waterfowl	Northern pintail	Wintering				-	-	-	-	-	-	-	-	-	-	-	-	100S	100S	-	-
621	Waterfowl	American black duck	Wintering	10S				-	-	-	-	-	-	-	-	-	-	-	-	100S	100S	-	-
		Waterfowl	American coot	Wintering				-	-	-	-	-	-	-	-	-	-	-	-	100S	100S	-	-
		Waterfowl	Mallard	Wintering				-	-	-	-	-	-	-	-	-	-	-	-	100S	100S	-	-
		Waterfowl	Northern pintail	Wintering				-	-	-	-	-	-	-	-	-	-	-	-	10S	10S	-	-
652	Waterfowl	American wigeon	Wintering	10S				-	-	-	-	-	-	-	-	-	-	-	-	100S	100S	-	-
		Waterfowl	Gadwall	Wintering				-	-	-	-	-	-	-	-	-	-	-	-	10S	10S	-	-
808	Waterfowl	American black duck	Wintering	10S				-	-	-	-	-	-	-	-	-	-	-	-	10S	10S	-	-
		Waterfowl	Bufflehead	Wintering				-	-	-	-	-	-	-	-	-	-	-	-	10S	10S	-	-
		Waterfowl	Canada goose	Wintering				-	-	-	-	-	-	-	-	-	-	-	-	100S	100S	-	-
		Waterfowl	Mallard	Wintering				-	-	-	-	-	-	-	-	-	-	-	-	Present	Present	-	-
		Waterfowl	Tundra swan	Wintering				-	-	-	-	-	-	-	-	-	-	-	-	10S	10S	-	-
1086	Diving	Common loon	Wintering	100S				-	-	-	-	-	-	-	-	-	-	-	-	100S	100S	-	-

Central Bay: Map 51C

BIRDS (continued)

Map ID	Subelement	Species	MD\VA			Monthly Presence												Molt		
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)
	Diving	D. crested cormorant	Migration			1,000S	-	-	-	-	-	-	-	-	-	-	-	-	Mar-May	Aug-Nov
	Diving	Grebes	Wintering			100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Diving	Red-throated loon	Wintering			100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pelagic	Northern gannet	Wintering			100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Scoters	Wintering			10,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Map ID	Subelement	Species	MD\VA			Monthly Presence												Spawn	Eggs	Larvae	Juveniles	Adults	
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
1662	Diadromous	Alewife	Migration			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Mar-May
	Diadromous	American shad	Migration			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Mar-Jun
	Diadromous	Blueback herring	Migration			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Feb-Jun
	Diadromous	Hickory shad	Migration			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Oct	Feb-Jun
	Diadromous	Striped bass	Migration			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Jun	

HABITATS & RARE PLANTS

Map ID	Subelement	Species	MD\VA			Monthly Presence												Mating	Calving	Pupping	Molt
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping
1727	Plant	Endangered plant	Vulnerable Occurrence	E-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

MARINE MAMMALS

Map ID	Subelement	Species	MD\VA			Monthly Presence												Mating	Calving	Pupping	Molt
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping
1849	Whale	Humpback whale	General Distribution	E/E	E	Very Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

BIRDS

Map ID	Subelement	Species	MD\VA			Monthly Presence												Nest	Mig.(S)	Mig.(F)	Molt
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)
	Diving	Common loon	Wintering			Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		Grebes	Wintering			100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		Red-throated loon	Wintering			Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pelagic	Northern gannet	Wintering			Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Long-tailed duck	Wintering			100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		Scoters	Wintering			100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Subelement	Species	MDVA		Monthly Presence												Adults									
		S	F	Mapping Qualifier		Concentration		J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles		
Diadromous	Alewife	Migration	Moderate Run	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Mar-May	
	American shad	Migration	Strong Run	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Mar-Jun
	Atlantic sturgeon	General Distribution	-E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	
	Blueback herring	Migration	Moderate Run	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun	
	Hickory shad	Migration	Moderate Run	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun	
	Shortnose sturgeon	General Distribution	E/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	
	Shortnose sturgeon	Migration	E/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Jun	
	Striped bass	Nursery Area	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec	

REPTILES & AMPHIBIANS

Subelement	Species	MDVA		Monthly Presence												Adults							
		S	F	Mapping Qualifier		Concentration		J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles
Turtle	K. ridley sea turtle	General Distribution	E/E	E	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct
	K. ridley sea turtle	General Distribution	E/E	E	Uncommon	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Dec
	Loggerhead sea turtle	General Distribution	T/T	T	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Dec
	Loggerhead sea turtle	General Distribution	T/T	T	Abundant	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct

INVERTEBRATES

Subelement	Species	MDVA		Monthly Presence												Adults							
		S	F	Mapping Qualifier		Concentration		J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles
Crab	Blue crab	Concentration Area	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Oct
		Highly Abundant-Summ	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

MARINE MAMMALS

Subelement	Species	MDVA		Monthly Presence												Adults							
		S	F	Mapping Qualifier		Concentration		J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping	Molt
Manatee	West Indian manatee	General Distribution	-E	E	Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
						-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)

BIRDS

Subelement	Species	MDVA		Monthly Presence												Adults							
		J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt						
Diving	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Gull/Tern	Gulls	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Terns	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Pelagic	Storm-petrels	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Diadromous	Alewife	-	-	-	-	-	-	-	-	-	-	-	-	Mar-May
	American eel	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct
	Blueback herring	-	-	-	-	-	-	-	-	-	-	-	-	Jan-May
	Gizzard shad	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Striped bass	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Estuarine Nursery	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Atlantic menhaden	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Black drum	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Black sea bass	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Bluefish	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Red drum	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Spot	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Apr
	Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Dec
	Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Weakfish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	White perch	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
Estuarine Resident	Silversides	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Fish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Anchovies	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Striped burrfish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Atlantic spadefish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Sep
	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov

INVERTEBRATES

	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Bivalve	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Oct
	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Crab	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Horseshoe crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec

MARINE MAMMALS

	Species	Monthly Presence												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
Dolphin	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-
	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-
Crab	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	-
	Harbor porpoise	-	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES

ESI POLYGON HABITAT TYPES		Habitat Classification	Area (Acres)	Area (Sq. Miles)
ESI Rank				
10A		Salt and Brackish Water Marshes	171.67	0.27
10B		Freshwater Marshes	2.95	0.00
10C		Swamps	171.95	0.27
10D		Scrub and Shrub Wetlands	25.95	0.04
7		Exposed Tidal Flats	2.53	0.00

ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
ESI Rank					
10A		Salt and Brackish Water Marshes	15,506.39	9.64	33%
10C		Swamps	1,915.11	1.19	4%
10D		Scrub and Shrub Wetlands	1,508.46	0.94	3%
9B		Vegetated Low Banks	1,762.74	1.10	4%
8B		Sheltered, Solid Man-Made Structures	862.24	0.54	2%
8C		Sheltered Riprap	1,127.42	0.70	2%
7		Exposed Tidal Flats	12,239.32	7.61	26%
6B		Riprap	6,637.60	4.12	14%
4		Coarse Grained Sand Beaches	3,945.10	2.45	9%
1B		Exposed, Solid Man-Made Structures	897.91	0.56	2%
Total ESI Shoreline:			46,402.29	Total ESI Shoreline: 28.83	
Total Shoreline:			29,668.26	Total Shoreline: 18.44	

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov



76° 30' W

76° 28' W


76° 26' W

Map 51B

SCATTERED ALONG SHORELINES



232



Map 51D Central Bay



SEE BACK OF MAP
for details about mapped species
and other species that occur in mapped area.
Data Published: September 2016

0 Not for Navigation 1 Miles
0 1 Kilometers

1:50,000



Map 51D Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BIRDS

Map ID	Subelement	Species	Mapping Qualifier	MD/VA	S	F	Concentration	Monthly Presence												
								J	F	M	A	M	J	J	A	S	O	N	D	Nest
44	Wading	Great blue heron	Nesting	62 Pairs				-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
120	Wading	Great blue heron	Nesting					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
148	Wading	Great blue heron	Nesting					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
194	Wading	Great blue heron	Nesting					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
232	Waterfowl	Canada goose	Nesting					-	-	-	-	-	-	-	-	-	-	-	-	Mar-Aug
	Waterfowl	Mallard	Nesting					-	-	-	-	-	-	-	-	-	-	-	-	Mar-Aug
496	Wading	Great blue heron	Nesting					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
751	Waterfowl	American black duck	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Waterfowl	American coot	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Waterfowl	American wigeon	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Waterfowl	Bufflehead	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Waterfowl	Canvasback	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Waterfowl	Common goldeneye	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Waterfowl	Gadwall	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Waterfowl	Mallard	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Waterfowl	Mergansers	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Waterfowl	Ring-necked duck	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Waterfowl	Ruddy duck	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Waterfowl	Scaup	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Waterfowl	Snow goose	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Waterfowl	Tundra swan	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
798	Raptor	Bald eagle	Nesting					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
808	Waterfowl	American black duck	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Waterfowl	Bufflehead	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Waterfowl	Canada goose	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Waterfowl	Mallard	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Waterfowl	Tundra swan	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
1034	Diving	Common loon	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Diving	D. crested cormorant	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Diving	Grebes	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Diving	Red-throated loon	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Pelagic	Northern gannet	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Waterfowl	Scoters	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
1365	Gull/Tern	G. black-backed gull	Nesting					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Gull/Tern	Herring gull	Nesting					-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun

FISH

Map ID	Subelement	Species	MD/V/A			Mapping Qualifier			MD/V/A			Monthly Presence									
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles
1620	Diadromous	Alewife	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Feb-Apr
	Diadromous	Blueback herring	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Mar-Jun

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

BIRDS

Subelement	Species	MD/V/A			Mapping Qualifier			MD/V/A			Monthly Presence			MD/V/A			Mapping Qualifier			MD/V/A		
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt		
Diving	Common loon	Wintering																				
	Grebes	Wintering																				
	Red-throated loon	Wintering																				
Pelagic	Northern gannet	Wintering																				
	Bufflehead	Wintering																				
Watertowl	Canada goose	Wintering																				
	Canvasback	Wintering																				
	Common goldeneye	Wintering																				
	Long-tailed duck	Wintering																				
	Mergansers	Wintering																				
	Redhead	Wintering																				
	Ring-necked duck	Wintering																				
	Ruddy duck	Wintering																				
	Scaup	Wintering																				
	Scoters	Wintering																				
	Snow goose	Wintering																				
	Tundra swan	Wintering																				

FISH

Subelement	Species	MD/V/A			Mapping Qualifier			MD/V/A			Monthly Presence			MD/V/A			Mapping Qualifier			MD/V/A		
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt		
Diadromous	Alewife	Migration																				
	American shad	Migration																				
	Atlantic sturgeon	General Distribution	-E	E	-																	
	Blueback herring	Migration																				
	Hickory shad	Migration																				
	Shortnose sturgeon	General Distribution	E/E	E	-																	
	Shortnose sturgeon	Migration	E/E	E	-																	
	Striped bass	Nursery Area	-																			

REPTILES & AMPHIBIANS

Subelement	Species	MD/V/A			Mapping Qualifier			MD/V/A			Monthly Presence			MD/V/A			Mapping Qualifier			MD/V/A		
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles	Adults	
Turtle	K. ridley sea turtle	General Distribution	E/E	E	-												-	-	-	Nov-Dec	-	
	K. ridley sea turtle	General Distribution	E/E	E	-												-	-	-	Apr-Oct	-	
	Loggerhead sea turtle	General Distribution	T/T	T	-												-	-	-	Nov-Dec	Nov-Dec	
	Loggerhead sea turtle	General Distribution	T/T	T	-												-	-	-	Apr-Oct	Apr-Oct	

Central Bay: Map 51D

INVERTEBRATES

Subelement	Species	Mapping Qualifier	MDVA	MDVA	Monthly Presence
Crab	Blue crab	Concentration Area	S F	Concentration	J F M A M J J A S O N D
		Highly Abundant-Summ			-

MARINE MAMMALS

Subelement	Species	Mapping Qualifier	MDVA	MDVA	Monthly Presence
Manatee	West Indian manatee	General Distribution	S F	Concentration	J F M A M J J A S O N D
		-/E E	Rare		-

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)

BIRDS

Subelement	Species	Mapping Qualifier	MDVA	MDVA	Monthly Presence
Diving	Brown pelican	General Distribution	S F	Concentration	J F M A M J J A S O N D
	D. crested cormorant				-
	D. crested cormorant				-
Gull/Tern	Gulls				-
	Terns				-

FISH

Subelement	Species	Mapping Qualifier	MDVA	MDVA	Monthly Presence
Diadromous	Alewife	General Distribution	S F	Concentration	J F M A M J J A S O N D
	American eel				-
	Blueback herring				-
	Gizzard shad				-
	Striped bass				-
Estuarine Nursery	Atlantic croaker	General Distribution	S F	Concentration	J F M A M J J A S O N D
	Atlantic menhaden				-
	Black drum				-
	Black sea bass				-
	Bluefish				-
	Northern puffer				-
	Red drum				-
	Spot				-
	Spotted seatrout				-
	Summer flounder				-
	Weakfish				-
	White perch				-
Estuarine Resident	Silversides	General Distribution	S F	Concentration	J F M A M J J A S O N D
	Anchovies				-
	Striped burrfish				-
	Atlantic spadefish				-
	Cownose ray				-
	Northern searobin				-

FISH (continued)

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Marine Pelagic	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Marine Pelagic	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Sep
Marine Pelagic	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov

INVERTEBRATES														
Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Bivalve	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Bivalve	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Crab	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Crab	Stout razor clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Crab	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Crab	Horseshoe crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec

MARINE MAMMALS														
Subelement	Species	Monthly Presence												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
Dolphin	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES

ESI POLYGON HABITAT TYPES		Habitat Classification	Area (Acres)	Area (Sq. Miles)
ESI Rank				
10A		Salt and Brackish Water Marshes	154.33	0.24
10B		Freshwater Marshes	4.31	0.01
10C		Swamps	79.64	0.12
10D		Scrub and Shrub Wetlands	15.92	0.02
7		Exposed Tidal Flats	2.69	0.00


ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
ESI Rank					
10A		Salt and Brackish Water Marshes	18,135.62	11.27	14%
10C		Swamps	808.99	0.50	1%
10D		Scrub and Shrub Wetlands	416.82	0.26	< 1%
9A		Sheltered Tidal Flats	152.00	0.09	< 1%
9B		Vegetated Low Banks	51,288.88	31.87	39%
8B		Sheltered, Solid Man-Made Structures	18,209.23	11.31	14%
8C		Sheltered Riprap	9,869.34	6.13	7%
7		Exposed Tidal Flats	8,838.43	5.49	7%
6B		Riprap	8,371.53	5.20	6%
4		Coarse Grained Sand Beaches	10,117.31	6.29	8%
3A		Fine to Medium Grained Sand Beaches	83.86	0.05	< 1%
3B		Scarps and Steep Slopes (Sand)	1,249.33	0.78	1%
1B		Exposed, Solid Man-Made Structures	4,716.70	2.93	4%

Total ESI Shoreline:	132,258.03	Total ESI Shoreline:	82.18
Total Shoreline:	114,091.71	Total Shoreline:	70.89

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 52
Central Bay



Map 52
Central Bay

HUMAN USE RESOURCES

DISPLAYED ON MAP (POINTS)

Map ID	Type	Name	Contact	Phone
1	AIRPORT	SANFORD FIELD		
2	AIRPORT	SHIVOK		
3	AIRPORT	WHITHALL FARM		
4	AQUACULTURE	OYSTER COMMERCIAL GROAT RF#020331	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
5	AQUACULTURE	OYSTER NON-COMM BORMAN RF#98-0761	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
6	AQUACULTURE	OYSTER NON-COMM CHES BAY GOVERNOR'S RF#99-2003	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
7	AQUACULTURE	OYSTER NON-COMM CORNWELL RF#98-1434	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
8	AQUACULTURE	OYSTER NON-COMM CORNWELL RF#981190	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
9	AQUACULTURE	OYSTER NON-COMM DENTON, STEPHEN J. RF#98-1504	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
10	AQUACULTURE	OYSTER NON-COMM DREW RF#99-1250	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
11	AQUACULTURE	OYSTER NON-COMM EVANS, JOHN W. RF#99-0177	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
12	AQUACULTURE	OYSTER NON-COMM FEUERSTEIN RF#98-1754	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
13	AQUACULTURE	OYSTER NON-COMM FOURNIER RF#98-1959	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
14	AQUACULTURE	OYSTER NON-COMM FREEMAN, ANNE H. RF#2000-1799	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
15	AQUACULTURE	OYSTER NON-COMM HICKOX RF#980775	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
16	AQUACULTURE	OYSTER NON-COMM HOFFMAN, KATHERINE RF#98-1770	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
17	AQUACULTURE	OYSTER NON-COMM JOHNSON RF#98-1148	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
18	AQUACULTURE	OYSTER NON-COMM PLOGER, WAYNE D. RF#98-1981	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
19	AQUACULTURE	OYSTER NON-COMM PULLEY RF#98-1864	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
20	AQUACULTURE	OYSTER NON-COMM RENFRO, JAMES G. RF#99-1955	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
21	AQUACULTURE	OYSTER NON-COMM RILEY RF#98-1979	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
22	AQUACULTURE	OYSTER NON-COMM SKAPARS RF#98-2252	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
23	AQUACULTURE	OYSTER NON-COMM STICKLER, SHAWN RF#99-1265	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
24	AQUACULTURE	OYSTER NON-COMM WARING, ROBERT L. RF#2001-0614	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
25	AQUACULTURE	OYSTER NON-COMM WOOLEY RF#98-1975	V/A DEPARTMENT OF GAME AND INLAND FISHERIES	757-247-2254
26	BOAT RAMP	BONUM'S CREEK		804-367-1000
27	BOAT RAMP	GARDY'S MILL POND		804-367-0058
28	BOAT RAMP	LOGDE LANDING		804-367-0058
29	BOAT RAMP	PINEY POINT LIGHTHOUSE MUSEUM AND PARK	ST. MARY'S COUNTY - DEPT. OF REC. AND PARKS	
30	REPEATED MEASUREMENT SITE	DB - NOS, 8578240 - PINY POINT, MD	NATIONAL COASTAL DATA DEVELOPMENT CENTER	866-732-2382
31	REPEATED MEASUREMENT SITE	MWS - SITE PRRP	GREG PINIAK, PROJECT MANAGER	301-713-3028
32	REPEATED MEASUREMENT SITE	WQ - SHELLFISH HARVEST WATERS, PINY POINT CREEK	MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
33	REPEATED MEASUREMENT SITE	WQ - SHELLFISH HARVEST WATERS, POTOMAC RIVER	MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
34	REPEATED MEASUREMENT SITE	WQ - SHELLFISH HARVEST WATERS, ST GEORGES CREEK	MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
ALSO PRESENT IN MAPPED AREA (POLYGONS)				
	Type	Name	Contact	Phone
	STATE PROTECTED AREA	PINEY POINT SANCTUARY	MARYLAND DEPARTMENT OF NATURAL RESOURCES	877-620-8367

JURISDICTIONS

COUNTY: NORTHUMBERLAND CO., RICHMOND CO., ST. MARY'S CO., WESTMORELAND CO.
 COAST GUARD: DISTRICT 5, SECTOR BALTIMORE, SECTOR HAMPTON ROADS
 USACE: NORTH ATLANTIC DIVISION, BALTIMORE DISTRICT, NORFOLK DISTRICT

FEMA:

REGION I
REGION 3

EPA:

REGION I
REGION 3**SHORELINE RESOURCES****ESI POLYGON HABITAT TYPES**

ESI Rank	Habitat Classification	Area (Acres)	Area (Sq. Miles)
10A	Salt and Brackish Water Marshes	672.64	1.05
10B	Freshwater Marshes	423.52	0.66
10C	Swamps	3,663.11	5.72
10D	Scrub and Shrub Wetlands	687.43	1.07

ESI SHORELINE HABITAT TYPES


ESI Rank	Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
10A	Salt and Brackish Water Marshes	64,625.60	40.16	18%
10B	Freshwater Marshes	8,874.61	5.51	2%
10C	Swamps	4,422.67	2.75	1%
10D	Scrub and Shrub Wetlands	2,513.89	1.56	1%
9A	Sheltered Tidal Flats	410.67	0.26	< 1%
9B	Vegetated Low Banks	127,586.26	79.28	35%
8B	Sheltered, Solid Man-Made Structures	45,949.10	28.55	13%
8C	Sheltered Riprap	27,537.06	17.11	8%
7	Exposed Tidal Flats	10,636.69	6.61	3%
6B	Riprap	23,507.67	14.61	6%
5	Mixed Sand and Gravel Beaches	1,089.08	0.68	< 1%
4	Coarse Grained Sand Beaches	23,452.13	14.57	6%
3B	Scars and Steep Slopes (Sand)	3,726.35	2.32	1%
1B	Exposed, Solid Man-Made Structures	21,956.83	13.64	6%

Total ESI Shoreline: 366,288.62 Total ESI Shoreline: 227.60
 Total Shoreline: 324,120.48 Total Shoreline: 201.40

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%.

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 52A
Central Bay



SEE BACK OF MAP
for details about mapped species and
other species that occur in mapped area.
Data Published: September 2016

0 Not for Navigation 1 Miles
0 1 Kilometers
1:50,000

Map 52A Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BIRDS

Map ID	Subelement	Species	Mapping Qualifier	S	F	Concentration	Monthly Presence																
							J	F	M	A	M	J	J	A	S	O	N	D	Nest	Feb-Jun	Mig.(S)	Mig.(F)	Molt
151	Wading	Great blue heron	Nesting																				
232	Waterfowl	Canada goose	Nesting																	Mar-Aug			
620	Waterfowl	American black duck	Wintering																		Jun-Jul		
	Waterfowl	American coot	Wintering																		Jul-Aug		
	Waterfowl	American wigeon	Wintering																				
	Waterfowl	Gadwall	Wintering																				
	Waterfowl	Mallard	Wintering																				
	Waterfowl	Northern pintail	Wintering																				
707	Raptor	Bald eagle	Roosting																				
752	Waterfowl	American black duck	Wintering																	100S			
	Waterfowl	American coot	Wintering																	10S			
	Waterfowl	American wigeon	Wintering																				
	Waterfowl	Bufflehead	Wintering																				
	Waterfowl	Canvasback	Wintering																				
	Waterfowl	Common goldeneye	Wintering																				
	Waterfowl	Gadwall	Wintering																				
	Waterfowl	Mallard	Wintering																				
	Waterfowl	Mergansers	Wintering																				
	Waterfowl	Ring-necked duck	Wintering																				
	Waterfowl	Ruddy duck	Wintering																				
	Waterfowl	Scaup	Wintering																				
	Waterfowl	Snow goose	Wintering																				
798	Raptor	Bald eagle	Nesting																1 Nest				
824	Waterfowl	Bufflehead	Wintering																				
	Waterfowl	Ruddy duck	Wintering																				
	Waterfowl	Scaup	Wintering																				
1163	Diving	Common loon	Wintering																				
	Diving	Grebes	Wintering																				
	Diving	Red-throated loon	Wintering																				
	Pelagic	Northern gannet	Wintering																				
	Waterfowl	Canada goose	Wintering																				
	Waterfowl	Scaup	Wintering																				
	Waterfowl	Scoters	Wintering																				

FISH

Map ID	Subelement	Species	Mapping Qualifier	MD/VA	S	F	Concentration	Monthly Presence							Adults	
1620	Diadromous	Alewife	Migration	-				-	-	-	-	-	-	-	-	Feb-Apr
	Diadromous	Blueback herring	Migration	Moderate Run				-	-	-	-	-	-	-	-	Mar-Jun
1674	Diadromous	Alewife	Nursery Area	Moderate Run				-	-	-	-	-	-	-	-	Mar-May
	Diadromous	Blueback herring	Nursery Area	Moderate Run				-	-	-	-	-	-	-	-	Feb-Jun

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

BIRDS

	Subelement	Species	Mapping Qualifier	MD/VA	S	F	Concentration	Monthly Presence							Adults				
		Common loon	Wintering	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S) <th>Mig.(F)</th> <th>Molt</th>	Mig.(F)	Molt
	Diving	D. crested cormorant	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-May	Aug-Nov	-
		Grebes	Wintering	10S	1,000S	10S	10S	10S	10S	10S	10S	10S	10S	10S	10S	-	-	-	-
	Pelagic	Red-throated loon	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Northern gannet	Wintering	100S	1,000S	100S	1,000S	100S	1,000S	100S	1,000S	100S	1,000S	100S	1,000S	-	-	-	-
		Bufflehead	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		Canada goose	Wintering	10,000S	10,000S	10,000S	10,000S	10,000S	10,000S	10,000S	10,000S	10,000S	10,000S	10,000S	10,000S	-	-	-	-
		Canvasback	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		Common goldeneye	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		Long-tailed duck	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		Mergansers	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		Redhead	Wintering	100S	100S	100S	100S	100S	100S	100S	100S	100S	100S	100S	100S	-	-	-	-
		Ring-necked duck	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		Ruddy duck	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		Scaup	Wintering	1,000S	1,000S	1,000S	1,000S	1,000S	1,000S	1,000S	1,000S	1,000S	1,000S	1,000S	1,000S	-	-	-	-
		Scoters	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		Snow goose	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		Tundra swan	Wintering	100S	100S	100S	100S	100S	100S	100S	100S	100S	100S	100S	100S	-	-	-	-

FISH

	Subelement	Species	Mapping Qualifier	MD/VA	S	F	Concentration	Monthly Presence							Adults									
	Diadromous	Alewife	Migration	Moderate Run				-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-May
		American shad	Migration	Strong Run				-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Jun
		Atlantic sturgeon	General Distribution	-E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		Blueback herring	Migration	Moderate Run				-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
		Hickory shad	Migration	General Distribution	E/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
		Shortnose sturgeon	Migration	General Distribution	E/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		Shortnose sturgeon	Nursery Area	Striped bass	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Jun

REPTILES & AMPHIBIANS

Subelement	Species	Mapping Qualifier	Monthly Presence											
			J	F	M	A	M	J	J	A	S	O	N	D
Turtle	K. ridley sea turtle	General Distribution	E/E	E	Concentration	-	-	-	-	Hatch	Internest	Juveniles	Adults	-
	K. ridley sea turtle	General Distribution	E/E	E	Common	-	-	-	-	-	-	Nov-Dec	-	Apr-Oct
	Loggerhead sea turtle	General Distribution	T/T	T	Common	-	-	-	-	-	-	Nov-Dec	Nov-Dec	Apr-Oct
	Loggerhead sea turtle	General Distribution	T/T	T	Abundant	-	-	-	-	-	-	Jun-Oct	Jun-Oct	Apr-Oct

INVERTEBRATES

Subelement	Species	Mapping Qualifier	Monthly Presence											
			J	F	M	A	M	J	J	A	S	O	N	D
Crab	Blue crab	Concentration Area	-	-	Highly Abundant-Summ	-	-	-	-	-	-	-	-	-
			-	-	-	-	-	-	-	-	-	-	-	-

MARINE MAMMALS

Subelement	Species	Mapping Qualifier	Monthly Presence											
			J	F	M	A	M	J	J	A	S	O	N	D
Manatee	West Indian manatee	General Distribution	-/E	E	Rare	-	-	-	-	-	-	-	-	-
			-	-	-	-	-	-	-	-	-	-	-	-

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)

BIRDS

Subelement	Species	Mapping Qualifier	Monthly Presence											
			J	F	M	A	M	J	J	A	S	O	N	D
Diving	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-	-
	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-	-
Gull/Tern	Gulls	-	-	-	-	-	-	-	-	-	-	-	-	-
	Terns	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Subelement	Species	Mapping Qualifier	Monthly Presence											
			J	F	M	A	M	J	J	A	S	O	N	D
Diadromous	Alewife	-	-	-	-	-	-	-	-	-	-	-	-	-
	American eel	-	-	-	-	-	-	-	-	-	-	-	-	-
	Blueback herring	-	-	-	-	-	-	-	-	-	-	-	-	-
	Gizzard shad	-	-	-	-	-	-	-	-	-	-	-	-	-
	Striped bass	-	-	-	-	-	-	-	-	-	-	-	-	-
Estuarine Nursery	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-	-
	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-	-
	Atlantic herring	-	-	-	-	-	-	-	-	-	-	-	-	-
	Atlantic menhaden	-	-	-	-	-	-	-	-	-	-	-	-	-
	Black drum	-	-	-	-	-	-	-	-	-	-	-	-	-
	Black sea bass	-	-	-	-	-	-	-	-	-	-	-	-	-
	Bluefish	-	-	-	-	-	-	-	-	-	-	-	-	-
	Kingfishes	-	-	-	-	-	-	-	-	-	-	-	-	-
	Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-	-
	Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH (continued)

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Pigfish	■												-
	Red drum													Apr-Nov
	Spot													May-Nov
	Spotted seatrout													Apr-Dec
	Spotted seatrout													Apr-Dec
	Summer flounder													Apr-Nov
	Weakfish													Apr-Dec
	White perch													Jan-Dec
	Pipefish													Jan-Dec
	Silversides													Jan-Dec
	Anchovies													Mar-Nov
	Striped burrfish													Jan-Dec
	Blue catfish													May-Oct
	Bullhead catfish													Jan-Dec
	Atlantic spadefish													Jan-Dec
	Cownose ray													Jun-Sep
	Northern searobin													May-Oct
	Northern searobin													May-Oct
	Spotted hake													May-Oct
	Butterfish													Apr-Sep
	Butterfish													Apr-Dec
	Harvestfish													Jan-Dec
	Lined seahorse													Jan-Dec
	Spanish mackerel													May-Sep

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Eastern oyster													Jan-Dec
	Softshell clam													Jan-Dec
	Softshell clam													Jan-Dec
	Stout razor clam													Jan-Dec
	Blue crab													Jan-Dec
	Horseshoe crab													Jan-Dec
	Mantis shrimp													Jan-Dec

MARINE MAMMALS

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Bottlenose dolphin													-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

ESI POLYGON HABITAT TYPES	
	Habitat Classification
10A	Salt and Brackish Water Marshes
10B	Freshwater Marshes
10C	Swamps
10D	Scrub and Shrub Wetlands

ESI Rank	Habitat Classification	Area (Acres)	Area (Sq. Miles)
10A	Salt and Brackish Water Marshes	167.49	0.26
10B	Freshwater Marshes	89.90	0.14
10C	Swamps	482.92	0.75
10D	Scrub and Shrub Wetlands	79.67	0.12


ESI SHORELINE HABITAT TYPES

ESI Rank	Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
10A	Salt and Brackish Water Marshes	17,158.15	10.66	20%
10B	Freshwater Marshes	321.35	0.20	< 1%
10C	Swamps	123.40	0.08	< 1%
10D	Scrub and Shrub Wetlands	78.31	0.05	< 1%
9B	Vegetated Low Banks	25,587.69	15.90	29%
8B	Sheltered, Solid Man-Made Structures	5,260.73	3.27	6%
8C	Sheltered Riprap	2,732.60	1.70	3%
7	Exposed Tidal Flats	5,810.14	3.61	7%
6B	Riprap	7,750.67	4.82	9%
5	Mixed Sand and Gravel Beaches	40.37	0.03	< 1%
4	Coarse Grained Sand Beaches	11,586.54	7.20	13%
3B	Scars and Steep Slopes (Sand)	2,076.88	1.29	2%
1B	Exposed, Solid Man-Made Structures	8,510.95	5.29	10%
Total ESI Shoreline:		87,037.77	Total ESI Shoreline:	54.08
Total Shoreline:		67,849.21	Total Shoreline:	42.16

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 52B
Central Bay



SEE BACK OF MAP
for details about mapped species
and other species that occur in mapped area.
Data Published: September 2016



Map 52B Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	Mapping Qualifier	MD\VA	S	F	Concentration	Monthly Presence
SAV	Submerged aquatic veg	High Ecological Value				Ephemeral	J F M A M J J A S O N D
							- - - - -

BIRDS

Map ID	Subelement	Species	Mapping Qualifier	MD\VA	S	F	Concentration	Monthly Presence
			Nesting				3 Nests	J F M A M J J A S O N D
126	Wading	Great blue heron	Nesting				Present	- - - - -
232	Watertowl	Canada goose	Nesting				Present	- - - - -
	Watertowl	Mallard	Nesting				-	- - - - -
707	Raptor	Bald eagle	Roosting				-	- - - - -
752	Watertowl	American black duck	Wintering				100S	- - - - -
	Watertowl	American coot	Wintering				10S	- - - - -
	Watertowl	American wigeon	Wintering				Present	- - - - -
	Watertowl	Bufflehead	Wintering				1,000S	- - - - -
	Watertowl	Canvasback	Wintering				1,000S	- - - - -
	Watertowl	Common goldeneye	Wintering				10S	- - - - -
	Watertowl	Gadwall	Wintering				10S	- - - - -
	Watertowl	Mallard	Wintering				100S	- - - - -
	Watertowl	Mergansers	Wintering				100S	- - - - -
	Watertowl	Ring-necked duck	Wintering				100S	- - - - -
	Watertowl	Ruddy duck	Wintering				1,000S	- - - - -
	Watertowl	Scaup	Wintering				1,000S	- - - - -
	Watertowl	Snow goose	Wintering				Present	- - - - -
	798	Raptor	Nesting				1 Nest	- - - - -
966	Watertowl	Bald eagle	Wintering				1,000S	- - - - -
	Watertowl	Canada goose	Wintering				10S	- - - - -
	Watertowl	Tundra swan	Wintering				-	- - - - -

FISH

Map ID	Subelement	Species	Mapping Qualifier	MD\VA	S	F	Concentration	Monthly Presence
			Migration				-	J F M A M J J A S O N D
1620	Diadromous	Alewife	Migration				-	Aug-Nov
	Diadromous	Blueback herring	Migration				-	Feb-Apr
							-	Mar-Jun

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

BIRDS

Subelement	Species	MD\VA						Monthly Presence													
		Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mg.(S)	Mg.(F)	Molt
Diving	Common loon	Wintering			10S														-	-	-
	D. crested cormorant	Wintering			1,000S														Mar-May	Aug-Nov	-
Grebes	Wintering			10S															-	-	-
Red-throated loon	Wintering			10S															-	-	-
Pelagic	Northern gannet	Wintering			100S														-	-	-
Waterfowl	Bufflehead	Wintering			1,000S														-	-	-
	Canada goose	Wintering			10,000S														-	-	-
Canvasback	Wintering			1,000S															-	-	-
Common goldeneye	Wintering			100S															-	-	-
Long-tailed duck	Wintering			100S															-	-	-
Mergansers	Wintering			100S															-	-	-
Redhead	Wintering			100S															-	-	-
Ring-necked duck	Wintering			1,000S															-	-	-
Ruddy duck	Wintering			1,000S															-	-	-
Scaup	Wintering			1,000S															-	-	-
Scoters	Wintering			Present															-	-	-
Snow goose	Wintering			100S															-	-	-
Tundra swan	Wintering			100S															-	-	-

FISH

Subelement	Species	MD\VA						Monthly Presence														
		Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
Diadromous	Alewife	Migration	Moderate Run																-	Aug-Nov	Mar-May	Mar-Jun
	American shad	Migration	Strong Run																-	Aug-Nov	Jan-Dec	Feb-Jun
	Atlantic sturgeon	General Distribution	-E	E															-	Aug-Nov	Jun-Oct	Jan-Dec
	Blueback herring	Migration	Moderate Run																-	Jun-Oct	Feb-Jun	Mar-Jun
	Hickory shad	Migration	Moderate Run																-	Jan-Dec	Apr-Jun	Jan-Dec
	Shortnose sturgeon	General Distribution	E/E	E															-	-	-	-
	Shortnose sturgeon	Migration	E/E	E															-	Apr-Jun	Jan-Dec	-
	Striped bass	Nursery Area																	-	-	-	-

INVERTEBRATES

Subelement	Species	MD\VA						Monthly Presence														
		Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
Crab	Blue crab	Concentration Area	Highly Abundant-Summ																-	Jun-Oct	Jun-Oct	Jun-Oct

MARINE MAMMALS

Subelement	Species	MD\VA						Monthly Presence														
		Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping	Molt	
Manatee	West Indian manatee	General Distribution	-E	E	Rare														-	-	-	-

BIRDS

Sublement	Species	<i>Monthly Presence</i>												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
Diving	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-	-
Gull/Tern	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-	-
	Gulls	-	-	-	-	-	-	-	-	-	-	-	-	-
	Terns	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Sublement	Species	<i>Monthly Presence</i>												Adults			
		J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles
Diadromous	Alewife	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Apr	Feb-Apr	Feb-Apr	Feb-Apr
	American eel	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Blueback herring	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Gizzard shad	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Striped bass	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Estuarine Nursery	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Atlantic herring	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Atlantic menhaden	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Black drum	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Black sea bass	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Bluefish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Kingfishes	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pigfish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Red drum	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Spot	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Weakfish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	White perch	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Estuarine Resident	Pipefish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Silversides	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Anchoovies	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Freshwater	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Blue catfish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Bullhead catfish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Marine Benthic	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Marine Pelagic	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Lined seahorse	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Spanish mackerel	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Bivalve	Eastern oyster	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Jan-Dec
	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Stout razor clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Crab	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Mantis shrimp	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES

ESI POLYGON HABITAT TYPES

ESI/Rank	Habitat Classification	Area (Acres)		% of ESI Shoreline
		Area (Sq. Miles)	Length (Miles)	
10A	Salt and Brackish Water Marshes	354.31	18.60	23%
10B	Freshwater Marshes	150.70	2.59	3%
10C	Swamps	1,890.72	1,136.47	1%
10D	Scrub and Shrub Wetlands	227.67	1,482.92	1%

ESI SHORELINE HABITAT TYPES


ESI/Rank	Shoreline Habitat Classification	Length (Meters)		% of ESI Shoreline
		Length (Miles)	Length (Meters)	
10A	Salt and Brackish Water Marshes	29,932.98	29,932.98	23%
10B	Freshwater Marshes	4,175.39	4,175.39	3%
10C	Swamps	1,136.47	1,136.47	1%
10D	Scrub and Shrub Wetlands	1,482.92	1,482.92	1%
9B	Vegetated Low Banks	37,656.52	37,656.52	29%
8B	Sheltered, Solid Man-Made Structures	17,306.92	17,306.92	13%
8C	Sheltered Riprap	7,761.63	7,761.63	6%
7	Exposed Tidal Flats	1,713.18	1,713.18	1%
6B	Riprap	9,305.29	9,305.29	7%
5	Mixed Sand and Gravel Beaches	1,048.71	1,048.71	1%
4	Coarse Grained Sand Beaches	7,789.58	7,789.58	6%
3B	Scarps and Steep Slopes (Sand)	207.49	207.49	< 1%
1B	Exposed, Solid Man-Made Structures	9,199.10	9,199.10	7%

Total ESI Shoreline: 128,716.17 Total ESI Shoreline: 79.98
 Total Shoreline: 117,908.23 Total Shoreline: 73.26

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 52C Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BIRDS

Map ID	Subelement	Species	Mapping Qualifier	S	F	Concentration	Monthly Presence															
							J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
107	Wading	Great blue heron	Nesting			22 Nests													Feb-Jun	-	-	-
120	Wading	Great blue heron	Nesting			5 Nests													Feb-Jun	-	-	-
123	Wading	Great blue heron	Nesting			14 Nests													Feb-Jun	-	-	-
150	Wading	Great blue heron	Nesting			10 Nests													Feb-Jun	-	-	-
151	Wading	Great blue heron	Nesting			28 Nests													Feb-Jun	-	-	-
153	Wading	Great blue heron	Nesting			4 Nests													Feb-Jun	-	-	-
232	Watertowl	Canada goose	Nesting			Present													Mar-Aug	-	Jun-Jul	-
	Watertowl	Mallard	Nesting			Present													Mar-Aug	-	Jul-Aug	-
751	Watertowl	American black duck	Wintering			100S													-	-	-	-
	Watertowl	American coot	Wintering			10S													-	-	-	-
	Watertowl	American wigeon	Wintering			Present													-	-	-	-
	Watertowl	Bufflehead	Wintering			1,000S													1,000S	-	-	-
	Watertowl	Canvasback	Wintering			1,000S													1,000S	-	-	-
	Watertowl	Common goldeneye	Wintering			10S													10S	-	-	-
	Watertowl	Gadwall	Wintering			100S													100S	-	-	-
	Watertowl	Mallard	Wintering			100S													100S	-	-	-
	Watertowl	Mergansers	Wintering			100S													100S	-	-	-
	Watertowl	Ring-necked duck	Wintering			1,000S													1,000S	-	-	-
	Watertowl	Ruddy duck	Wintering			1,000S													1,000S	-	-	-
	Watertowl	Scaup	Wintering			Present													Present	-	-	-
	Watertowl	Snow goose	Wintering			100S													100S	-	-	-
	Watertowl	Tundra swan	Wintering			Dec-Jun													Dec-Jun	-	-	-
798	Raptor	Bald eagle	Nesting			1 Nest													1 Nest	-	-	-
824	Watertowl	Bufflehead	Wintering			10S													10S	-	-	-
	Watertowl	Ruddy duck	Wintering			1,000S													1,000S	-	-	-
	Watertowl	Scaup	Wintering			100S													100S	-	-	-
956	Watertowl	Canada goose	Wintering			1,000S													1,000S	-	-	-
1034	Diving	Common loon	Wintering			10S													10S	-	-	-
	Diving	D. crested cormorant	Wintering			1,000S													1,000S	-	-	-
	Diving	Grebes	Wintering			10S													10S	-	-	-
	Diving	Red-throated loon	Wintering			100S													100S	-	-	-
	Pelagic	Northern gannet	Wintering			1,000S													1,000S	-	-	-
	Watertowl	Scoters	Wintering			Present													Present	-	-	-
1162	Diving	Common loon	Wintering			100S													Present	-	-	-
	Diving	Grebes	Wintering			Present													Present	-	-	-
	Diving	Red-throated loon	Wintering			Present													Present	-	-	-
	Pelagic	Northern gannet	Wintering			Present													Present	-	-	-

BIRDS (continued)

Map ID	Subelement	Species	MD\VA			Monthly Presence															
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
	Waterfowl	Bufflehead	Wintering		1,000S													-	-	-	-
	Waterfowl	Canvasback	Wintering		1,000S													-	-	-	-
	Waterfowl	Common goldeneye	Wintering		100S													-	-	-	-
	Waterfowl	Mergansers	Wintering		100S													-	-	-	-
	Waterfowl	Redhead	Wintering		10S													-	-	-	-
	Waterfowl	Ring-necked duck	Wintering		100S												-	-	-	-	-
	Waterfowl	Ruddy duck	Wintering		10,000S												-	-	-	-	-
	Waterfowl	Scaup	Wintering		1,000S												-	-	-	-	-
	Waterfowl	Scoters	Wintering		100S												-	-	-	-	-
	Waterfowl	Snow goose	Wintering		Present												-	-	-	-	-
	Waterfowl	Tundra swan	Wintering		1,000S												-	-	-	-	-

FISH

Map ID	Subelement	Species	MD\VA			Monthly Presence															
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
1620	Diadromous	Alewife	Migration		-												-	-	Aug-Nov	Feb-Apr	-
	Diadromous	Blueback herring	Migration		-												-	-	Aug-Nov	Mar-Jun	-

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

Map ID	Subelement	Species	MD\VA			Monthly Presence															
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
	Waterfowl	Canada goose	Wintering		10,000S												-	-	-	-	-
		Long-tailed duck	Wintering		100S												-	-	-	-	-

FISH

Map ID	Subelement	Species	MD\VA			Monthly Presence															
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
	Diadromous	Alewife	Migration		Moderate Run												-	-	Aug-Nov	Mar-May	-
		American shad	Migration		Strong Run												-	-	Aug-Nov	Mar-Jun	-
		Atlantic sturgeon	General Distribution	-E	E												-	-	-	Jan-Dec	-
		Blueback herring	Migration		Moderate Run												-	-	Aug-Nov	Feb-Jun	-
		Hickory shad	Migration		Moderate Run												-	-	Jun-Oct	Feb-Jun	-
		Shortnose sturgeon	General Distribution	E/E	E												-	-	Jan-Dec	Jan-Dec	-
		Shortnose sturgeon	Migration	E/E	E												-	-	Mar-Jun	Mar-Jun	-
		Striped bass	Nursery Area														-	-	Apr-Jun	Jan-Dec	-

REPTILES & AMPHIBIANS

Map ID	Subelement	Species	MD\VA			Monthly Presence															
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Adults
	Turtle	K. ridley sea turtle	General Distribution	E/E	Uncommon												-	-	Nov-Dec	-	-
		K. ridley sea turtle	General Distribution	E/E	Common												-	-	Apr-Oct	-	-
		Loggerhead sea turtle	General Distribution	T/T	Common												-	-	Nov-Dec	Nov-Dec	-
		Loggerhead sea turtle	General Distribution	T/T	Abundant												-	-	Apr-Oct	Apr-Oct	-

Central Bay: Map 52C

INVERTEBRATES

Subelement	Species	Concentration Area	MDVA												Monthly Presence													
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults						
Crab	Blue crab	Highly Abundant-Summ	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		

MARINE MAMMALS

Subelement	Species	General Distribution	MDVA												Monthly Presence													
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping	Molt	Eggs	Larvae	Juveniles	Adults			
Manatee	West Indian manatee	-/E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)

BIRDS

Subelement	Species	General Distribution	MDVA												Monthly Presence											
			J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt	Eggs	Larvae	Juveniles	Adults				
Diving	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Gull/Tern	Gulls	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Terns	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH


Subelement	Species	General Distribution	MDVA												Monthly Presence											
			J	F	M	A	M	J	J	A	S	O	N	D	Feb-Apr	Feb-Apr	Feb-Apr	Feb-Apr	Feb-Apr	Feb-Apr	Feb-Apr	Feb-Apr	Feb-Apr	Feb-Apr	Feb-Apr	Feb-Apr
Diadromous	Alewife	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	American eel	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Blueback herring	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Gizzard shad	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Striped bass	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Estuarine Nursery	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Atlantic menhaden	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Black drum	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Black sea bass	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Bluefish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Red drum	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Spot	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Weakfish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	White perch	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Estuarine Resident	Silversides	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Anchoovies	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Striped burrfish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Marine Benthic	Atlantic spadefish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH (continued)

Subelement	Species	Monthly Presence												Larvae	Juveniles	Adults		
		J	F	M	A	M	J	J	A	S	O	N	D					
Spotted hake														-	-	Jan-Dec		
Butterfish														-	-	Jul-Sep		
Marine Pelagic														-	-	May-Nov		
INVERTEBRATES																		
Subelement	Species	Monthly Presence												Eggs	Larvae	Juveniles	Adults	
		J	F	M	A	M	J	J	A	S	O	N	D	Spawning	Eggs	Larvae	Juveniles	Adults
Bivalve	Eastern oyster													Jun-Sep	Jun-Sep	Jun-Oct	Jan-Dec	Jan-Dec
	Softshell clam													May-Dec*	May-Dec*	May-Jan*	-	Jan-Dec
	Softshell clam													May-Dec*	-	-	-	Jan-Dec
	Stout razor clam													Jun-Nov	-	-	-	Jan-Dec
Crab	Blue crab													-	-	-	-	Jan-Dec
	Horseshoe crab													-	-	-	-	Jan-Dec
MARINE MAMMALS																		
Subelement	Species	Monthly Presence												Mating	Calving	Pupping	Molt	
		J	F	M	A	M	J	J	A	S	O	N	D	-	-	-	-	
Dolphin	Bottlenose dolphin													-	-	-	-	

For additional information about species locations and extent, reference the underlying GIS data available from response.noaa.gov

ESI POLYGON HABITAT TYPES	
	Habitat Classification
10A	Salt and Brackish Water Marshes
10B	Freshwater Marshes
10C	Swamps
10D	Scrub and Shrub Wetlands



ESI SHORELINE HABITAT TYPES	
	Shoreline Habitat Classification
10A	Salt and Brackish Water Marshes
10B	Freshwater Marshes
10C	Swamps
10D	Scrub and Shrub Wetlands
9A	Sheltered Tidal Flats
9B	Vegetated Low Banks
8B	Sheltered, Solid Man-Made Structures
8C	Sheltered Riprap
7	Exposed Tidal Flats
6B	Riprap
4	Coarse Grained Sand Beaches
3B	Scars and Steep Slopes (Sand)
1B	Exposed, Solid Man-Made Structures



	Area (Acres)	Area (Sq. Miles)
10A	150,84	0.24
10B	58,06	0.09
10C	743,84	1.16
10D	111,59	0.17


	Length (Meters)	Length (Miles)	% of ESI Shoreline
10A	17,534,47	10.90	12%
10B	4,377,88	2.72	3%
10C	3,162,80	1.97	2%
10D	952,66	0.59	1%
9A	410,67	0.26	< 1%
9B	64,342,05	39.98	43%
8B	23,381,46	14.53	16%
8C	17,042,84	10.59	11%
7	3,113,37	1.93	2%
6B	6,451,71	4.01	4%
4	4,076,01	2.53	3%
3B	1,441,98	0.90	1%
1B	4,246,79	2.64	3%

Total ESI Shoreline:	150,534.68	Total ESI Shoreline:	93.54
Total Shoreline:	138,363.04	Total Shoreline:	85.97

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 57
Central Bay

HUMAN USE RESOURCES

DISPLAYED ON MAP (POINTS)			Name	Contact	Phone
Map ID	Type				
1	AQUACULTURE		OYSTER COMMERCIAL DREWER HARVEY RF#2002-0453	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
2	AQUACULTURE		OYSTER COMMERCIAL DREWER HARVEY RF#2002-1813	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
3	BOAT RAMP	HAMMOCK		LENE PENNINGTON, VDGIF	804-367-0058
4	BOAT RAMP	JENKINS CREEK		SOMERSET COUNTY - ROADS DEPARTMENT	
5	HELIPORT	MCCREADY MEMORIAL HOSPITAL		MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
6	REPEATED MEASUREMENT SITE	WQ - SHELLFISH HARVEST WATERS, POCOMOKE SOUND		MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
7	REPEATED MEASUREMENT SITE	WQ - SHELLFISH HARVEST WATERS, POKOMOKE SOUND		CHESAPEAKE BAY PROGRAM	800-968-7229
8	REPEATED MEASUREMENT SITE	WQ - STATION EE3.3		CHESAPEAKE BAY PROGRAM	800-968-7229
9	REPEATED MEASUREMENT SITE	WQ - STATION EE3.4		CHESAPEAKE BAY PROGRAM	800-968-7229
10	REPEATED MEASUREMENT SITE	WQ - STATION EE3.5		CHESAPEAKE BAY PROGRAM	800-968-7229
DISPLAYED ON MAP (POLYGONS)			Name	Contact	Phone
Map ID	Type				
11	MANAGEMENT AREA	CEDAR ISLAND WMA		WILDLIFE AND HERITAGE SERVICE	
12	MANAGEMENT AREA	GUARD SHORE WILDLIFE MANAGEMENT AREA		VA DEPT OF GAME AND INLAND FISHERIES	
13	MANAGEMENT AREA	LEVEL PONDS		VA DEPT OF GAME AND INLAND FISHERIES	
14	MANAGEMENT AREA	MARYLAND MARINE PROPERTIES WMA		WILDLIFE AND HERITAGE SERVICE	
15	MANAGEMENT AREA	POCOMOKE SOUND WMA		WILDLIFE AND HERITAGE SERVICE	
16	MANAGEMENT AREA	SAXIS WILDLIFE MANAGEMENT AREA		VA DEPT OF GAME AND INLAND FISHERIES	
17	NATURE CONSERVANCY	BIG MARSH		THE NATURE CONSERVANCY	703-841-5300
18	NATURE CONSERVANCY	CLAM MARSHES		THE NATURE CONSERVANCY	703-841-5300
19	NATURE CONSERVANCY	DEEP CREEK FEE		THE NATURE CONSERVANCY	703-841-5300
20	NATURE CONSERVANCY	LONG RIDGE		THE NATURE CONSERVANCY	703-841-5300
21	NATURE CONSERVANCY	THE NATURE CONSERVANCY LAND HOLDING		THE NATURE CONSERVANCY	703-841-5300
ALSO PRESENT IN MAPPED AREA (POLYGONS)			Name	Contact	Phone
Type					
STATE PROTECTED AREA		KITTS CREEK SANCTUARY		MARYLAND DEPARTMENT OF NATURAL RESOURCES	877-620-8367
STATE PROTECTED AREA		MARKS AND JACKS ISLANDS		VA DEPT OF CONSERVATION AND RECREATION	
STATE PROTECTED AREA		TIDAL LANDS		VA MARINE RESOURCES COMMISSION	

JURISDICTIONS

COUNTY: ACCOMACK COUNTY, SOMERSET COUNTY
COAST GUARD: DISTRICT 5, SECTOR BALTIMORE, SECTOR HAMPTON ROADS
USACE: NORTH ATLANTIC DIVISION, BALTIMORE DISTRICT, NORFOLK DISTRICT

FEMA:
REGION I
REGION 3

EPA:
REGION 3

SHORELINE RESOURCES**ESI POLYGON HABITAT TYPES**

ESI Rank	Habitat Classification	Area (Acres)	Area (Sq. Miles)
10A	Salt and Brackish Water Marshes	14,066.27	21.98
10B	Freshwater Marshes	842.10	1.32
10C	Swamps	1,517.47	2.37
10D	Scrub and Shrub Wetlands	565.63	0.88
9A	Sheltered Tidal Flats	14.11	0.02
7	Exposed Tidal Flats	215.69	0.34

ESI SHORELINE HABITAT TYPES


ESI Rank	Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
10A	Salt and Brackish Water Marshes	620,987.97	385.86	56%
10B	Freshwater Marshes	31,89	0.02	< 1%
10C	Swamps	11,516.44	7.16	1%
10D	Scrub and Shrub Wetlands	9,068.82	5.64	1%
9A	Sheltered Tidal Flats	397,008.06	246.69	36%
9B	Vegetated Low Banks	4,361.90	2.71	< 1%
8B	Sheltered, Solid Man-Made Structures	3,614.97	2.25	< 1%
8C	Sheltered Riprap	3,910.84	2.43	< 1%
7	Exposed Tidal Flats	25,385.06	15.77	2%
6B	Riprap	641.21	0.40	< 1%
5	Mixed Sand and Gravel Beaches	113.01	0.07	< 1%
4	Coarse Grained Sand Beaches	1,523.30	0.95	< 1%
3A	Fine to Medium Grained Sand Beaches	37,725.69	23.44	3%
3B	Scars and Steep Slopes (Sand)	297.45	0.18	< 1%
1B	Exposed, Solid Man-Made Structures	910.79	0.57	< 1%

Total ESI Shoreline: 1,117,097.40 Total ESI Shoreline: 694.13
 Total Shoreline: 634,937.22 Total Shoreline: 394.53

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 57A
Central Bay



SEE BACK OF MAP
for details about mapped species and
other species that occur in mapped area.
Data Published: September 2016



Map 57A Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	<i>Mapping Qualifier</i>						Concentration	<i>Monthly Presence</i>								
		S	M	D/V/A	F	Ephemeral	J	F	M	A	M	J	J	A	S	O	N
SAV	Submerged aquatic veg	High Ecological Value	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
SAV	Submerged aquatic veg	High Ecological Value	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
SAV	Submerged aquatic veg	High Ecological Value	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

BIRDS

Map ID	Subelement	Species	<i>Mapping Qualifier</i>						Concentration	<i>Monthly Presence</i>								
			S	M	D/V/A	F	E	T		J	F	M	A	M	J	J	A	S
410	Waterfowl	Brant	Wintering	-	-	-	-	-	100S	-	-	-	-	-	-	-	-	-
	Waterfowl	Northern pintail	Wintering	-	-	-	-	-	100S	-	-	-	-	-	-	-	-	-
	Waterfowl	Tundra swan	Wintering	-	-	-	-	-	1,000S	-	-	-	-	-	-	-	-	-
477	Waterfowl	American black duck	Nesting	-	-	-	-	-	High	Mar-Aug	-	-	-	-	-	-	-	-
	Waterfowl	Canada goose	Nesting	-	-	-	-	-	Moderate-High	Mar-Jun	-	-	-	-	-	-	-	-
535	Waterfowl	Bufflehead	Wintering	-	-	-	-	-	1,000S	-	-	-	-	-	-	-	-	-
	Waterfowl	Canada goose	Wintering	-	-	-	-	-	1,000S	-	-	-	-	-	-	-	-	-
	Waterfowl	Canvasback	Wintering	-	-	-	-	-	1,000S	-	-	-	-	-	-	-	-	-
	Waterfowl	Common goldeneye	Wintering	-	-	-	-	-	10S	-	-	-	-	-	-	-	-	-
	Waterfowl	Mergansers	Wintering	-	-	-	-	-	100S	-	-	-	-	-	-	-	-	-
	Waterfowl	Redhead	Wintering	-	-	-	-	-	1,000S	-	-	-	-	-	-	-	-	-
	Waterfowl	Ring-necked duck	Wintering	-	-	-	-	-	10S	-	-	-	-	-	-	-	-	-
	Waterfowl	Ruddy duck	Wintering	-	-	-	-	-	1,000S	-	-	-	-	-	-	-	-	-
	Waterfowl	Scaup	Wintering	-	-	-	-	-	100S	-	-	-	-	-	-	-	-	-
	Waterfowl	Snow goose	Wintering	-	-	-	-	-	1,000S	-	-	-	-	-	-	-	-	-
557	Waterfowl	American black duck	Wintering	-	-	-	-	-	1,000S	-	-	-	-	-	-	-	-	-
	Waterfowl	American coot	Wintering	-	-	-	-	-	10S	-	-	-	-	-	-	-	-	-
	Waterfowl	American wigeon	Wintering	-	-	-	-	-	100S	-	-	-	-	-	-	-	-	-
	Waterfowl	Gadwall	Wintering	-	-	-	-	-	1,000S	-	-	-	-	-	-	-	-	-
	Waterfowl	Green-winged teal	Wintering	-	-	-	-	-	1,000S	-	-	-	-	-	-	-	-	-
	Waterfowl	Mallard	Wintering	-	-	-	-	-	1,000S	-	-	-	-	-	-	-	-	-
	Waterfowl	Northern pintail	Wintering	-	-	-	-	-	10S	-	-	-	-	-	-	-	-	-
	Waterfowl	Northern shoveler	Wintering	-	-	-	-	-	Present	-	-	-	-	-	-	-	-	-
798	Raptor	Bald eagle	Nesting	-	-	-	-	-	1 Nest	-	-	-	-	-	-	-	-	-
924	Wading	Black rail	Vulnerable Occurrence	E/E	-	-	-	-	-	Dec-Jun	-	-	-	-	-	-	-	-
									May-Jul	-	-	-	-	-	-	-	-	-

INVERTEBRATES

Map ID	Subelement	Species	<i>Mapping Qualifier</i>						Concentration	<i>Monthly Presence</i>								
			S	M	D/V/A	F	E/T	T		J	F	M	A	M	J	J	A	S
1833	Insect	NE beach tiger beetle	-	-	-	-	-	-	Jun-Aug	-	-	-	-	-	-	-	-	-

Central Bay: Map 57A

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

BIRDS

	Sublement	Species	Mapping Qualifier	MD\VA	Monthly Presence												Molt	
					S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N
Diving	Common loon	Wintering		Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Grebes	Wintering		10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Red-throated loon	Wintering		100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Pelagic	Northern gannet	Wintering		Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Waterfowl	American black duck	Nesting		Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	American black duck	Wintering		1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	American wigeon	Wintering		100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	American wigeon	Migration		1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Blue-winged teal	Migration		100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Blue-winged teal	Brant	Wintering	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Bufflehead	Wintering		1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Canada goose	Nesting		Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Canada goose	Wintering		1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Canvasback	Wintering		1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Gadwall	Wintering		1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Gadwall	Wintering		100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Green-winged teal	Wintering		1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Green-winged teal	Wintering		1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Long-tailed duck	Wintering		100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Mallard	Nesting		Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Mallard	Wintering		100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Mergansers	Wintering		1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Northern pintail	Wintering		100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Northern shoveler	Wintering		10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Redhead	Wintering		Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Redhead	Wintering		1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Ring-necked duck	Wintering		10,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Ruddy duck	Wintering		Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Scaup	Wintering		10,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Scaup	Wintering		1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Scoters	Wintering		1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Snow goose	Wintering		100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Tundra swan	Wintering		100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Subelement	Species	MD\VA												Monthly Presence											
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults				
Diadromous	Alewife	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Feb-Apr			
	Alewife	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Mar-May			
	American shad	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Feb-Jun			
	American shad	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Mar-Jun			
	Atlantic sturgeon	General Distribution	-/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Jan-Dec			
	Blueback herring	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Feb-Jun			
	Blueback herring	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Mar-Jun			
	Hickory shad	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep	Feb-May			
	Hickory shad	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Oct	Feb-Jun			
	Shortnose sturgeon	General Distribution	E/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec			
	Striped bass	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun	Feb-Jun			
	Striped bass	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Jun	Mar-Jun			

REPTILES & AMPHIBIANS

Subelement	Species	MD\VA												Monthly Presence											
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults				
Turtle	Diamondback terrapin	Concentration Area	High	-	-	-	-	-	-	-	-	-	-	May-Aug	-	-	-	-	-	-	Apr-Oct	Apr-Oct			
	K. ridley sea turtle	General Distribution	E/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Dec	-			
	K. ridley sea turtle	General Distribution	E/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct	Apr-Oct			
	Loggerhead sea turtle	General Distribution	T/T	T	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Dec	Nov-Dec			
	Loggerhead sea turtle	General Distribution	T/T	T	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct	Apr-Oct			

INVERTEBRATES

Subelement	Species	MD\VA												Monthly Presence											
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults				
Crab	Blue crab	Concentration Area	Highly Abundant-Summ	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Mar*	Jun-Mar*			
	Blue crab	Concentration Area	Highly Abundant	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Mar*	Jun-Mar*			

MARINE MAMMALS

Subelement	Species	MD\VA												Monthly Presence											
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults				
Manatee	West Indian manatee	General Distribution	-/E	E	Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

BIRDS

		Monthly Presence												Eggs		Larvae		Juveniles		Adults	
		Subelement		Species		J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
	Diving	Brown pelican																-	-	-	-
	Gull/Tern	D. crested cormorant																-	-	-	-
	Gulls	Gulls																-	-	-	-
	Passerine	Terns																-	-	-	-
		Saltmarsh sharp-tailed sparrow																May-Aug	-	-	-
		Seaside sparrow																May-Jul	-	-	-
	Shorebird	Willet																May-Jul	-	-	-
	Wading	Clapper rail																Apr-Sep	-	-	-
	Waterfowl	Wood duck																Apr-Jul	-	Jul-Aug	-

FISH

		Monthly Presence												Eggs	Larvae	Juveniles	Adults			
		Subelement		Species		J	F	M	A	M	J	J	A	S	O	N	D	Mar-May	Apr-May	Mar-May
	Diadromous	Alewife		Alewife													Feb-Apr	Feb-Apr	Feb-Apr	Feb-Apr
		American eel															-	Jan-May	Jan-May	Jan-Dec
		Blueback herring															Apr-Jun	Apr-Jun	May-Jul	May-Oct
		Blueback herring															Mar-Jun	Apr-Jun	May-Oct	Mar-Jun
		Gizzard shad															Mar-Jun	-	Jan-Dec	Jan-Dec
		Striped bass															-	-	Jan-Dec	Jan-Dec
	Estuarine Nursery	Atlantic croaker															-	Dec-Apr	Jan-Dec	May-Nov
		Atlantic menhaden															-	Jan-Dec	Jan-Dec	Jan-Dec
		Black drum															Apr-Jun	-	Apr-Dec	Apr-Dec
		Black sea bass															-	Apr-Nov	Apr-Nov	Apr-Nov
		Bluefish															-	May-Oct	May-Oct	May-Oct
		Kingfishes															-	Apr-Nov	Apr-Nov	Apr-Nov
		Northern puffer															-	Mar-Oct	Mar-Nov	Mar-Nov
		Pigfish															-	Apr-Nov	Apr-Nov	Apr-Nov
		Red drum															-	May-Nov	May-Nov	May-Nov
		Sheepshead															-	Apr-Nov	Apr-Nov	Apr-Nov
		Spot															-	Nov-Apr	Apr-Dec	Apr-Dec
		Spotted seatrout															-	Apr-Nov	Apr-Nov	Apr-Nov
		Summer flounder															-	Feb-May	Apr-Dec	Jan-Dec
		Weakfish															-	May-Aug	Jun-Dec	Apr-Dec
		White perch															-	Jan-Dec	Jan-Dec	Jan-Dec
	Estuarine Resident	Winter flounder															-	-	Dec-Mar	Jan-Dec
		Pipefish															-	Mar-Jul	Mar-Jul	Mar-Nov
		Silversides															-	-	-	Jan-Dec
		Anchovies															-	-	-	Jan-Dec
		Striped burrfish															-	May-Oct	May-Oct	May-Oct
		Atlantic spadefish															-	-	-	Jun-Sep
		Clearnose skate															-	-	-	Jan-Dec

FISH (continued)

Subelement	Species	Monthly Presence											
		J	F	M	A	M	J	J	A	S	O	N	D
	Cownose ray	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Red hake	-	-	-	-	-	-	-	-	-	-	-	Sep-May*
	Silver hake	-	-	-	-	-	-	-	-	-	-	-	Oct-May
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Marine Pelagic	Butterfish	-	-	-	-	-	-	-	-	-	-	-	Jul-Nov
	Cobia	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Lined seahorse	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Spanish mackerel	-	-	-	-	-	-	-	-	-	-	-	May-Sep

REPTILES & AMPHIBIANS

Subelement	Species	Monthly Presence											
		J	F	M	A	M	J	J	A	S	O	N	D
	Turtle	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct
	Diamondback terrapin	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct

INVERTEBRATES

Subelement	Species	Monthly Presence											
		J	F	M	A	M	J	J	A	S	O	N	D
Bivalve	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Hard clam	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Stout razor clam	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Cephalopod	Squid	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Crab	Blue crab	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Horseshoe crab	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Brown shrimp	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Mantis shrimp	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	White shrimp	-	-	-	-	-	-	-	-	-	-	-	May-Oct

MARINE MAMMALS

Subelement	Species	Monthly Presence											
		J	F	M	A	M	J	J	A	S	O	N	D
Dolphin	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-
	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-
	Harbor porpoise	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES

ESI POLYGON HABITAT TYPES		Habitat Classification		Area (Acres)	Area (Sq. Miles)
ESI Rank					
10A		Salt and Brackish Water Marshes		3,987.92	6.23
10B		Freshwater Marshes		5.66	0.01
10C		Swamps		193.03	0.30
10D		Scrub and Shrub Wetlands		75.90	0.12
9A		Sheltered Tidal Flats		7.05	0.01
7		Exposed Tidal Flats		2.54	0.00


ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification		Length (Meters)	% of ESI Shoreline
ESI Rank					
10A		Salt and Brackish Water Marshes		142,150.69	55%
10C		Swamps		381.50	< 1%
10D		Scrub and Shrub Wetlands		4,909.21	2%
9A		Sheltered Tidal Flats		86,039.98	34%
9B		Vegetated Low Banks		490.41	< 1%
8B		Sheltered, Solid Man-Made Structures		1,802.82	1%
8C		Sheltered Riprap		3,253.24	1%
7		Exposed Tidal Flats		3,697.81	1%
6B		Riprap		327.26	< 1%
4		Coarse Grained Sand Beaches		1,184.24	< 1%
3A		Fine to Medium Grained Sand Beaches		11,410.17	4%
3B		Scarps and Steep Slopes (Sand)		297.45	< 1%
1B		Exposed, Solid Man-Made Structures		853.08	< 1%

Total ESI Shoreline: 256,797.87 Total ESI Shoreline: 159.57
 Total Shoreline: 148,905.43 Total Shoreline: 92.53

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





SEE BACK OF MAP
for details about mapped species and
other species that occur in mapped area.
Data Published: September 2016

0 Not for Navigation 1 Miles
0 1 Kilometers
1:50,000

Map 57B Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	Mapping Qualifier			S	F	Concentration	Monthly Presence										
		High Ecological Value	High Ecological Value	High Ecological Value				J	F	M	A	M	J	J	A	S	O	N
SAV	Submerged aquatic veg				-	-	-	-	-	-	-	-	-	-	-	-	-	-
SAV	Submerged aquatic veg				-	-	-	-	-	-	-	-	-	-	-	-	-	-
SAV	Submerged aquatic veg				-	-	-	-	-	-	-	-	-	-	-	-	-	-

BIRDS

Map ID	Subelement	Species	Mapping Qualifier			S	F	Concentration	Monthly Presence										
			J	F	M				J	F	M	A	M	J	J	A	S	O	N
404	Watrfowl	American black duck	Wintering			100S			-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	American wigeon	Wintering			1,000S			-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Blue-winged teal	Migration			100S			-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Brant	Wintering			1,000S			-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Canada goose	Wintering			1,000S			-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Canvasback	Wintering			1,000S			-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Gadwall	Wintering			1,000S			-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Green-winged teal	Wintering			100S			-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Mallard	Wintering			100S			-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Northern pintail	Wintering			100S			-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Redhead	Wintering			1,000S			-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Ruddy duck	Wintering			100S			-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Scaup	Wintering			100S			-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Tundra swan	Wintering			1,000S			-	-	-	-	-	-	-	-	-	-	-
447	Watrfowl	American black duck	Nesting			Moderate			Mar-Aug	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Mallard	Nesting			Moderate-High			Mar-Aug	-	-	-	-	-	-	-	-	-	-
1090	Diving	Common loon	Wintering			100S			-	-	-	-	-	-	-	-	-	-	-
	Diving	D. crested cormorant	Migration			1,000S			-	-	-	-	-	-	-	-	-	-	-
	Diving	Grebes	Wintering			100S			-	-	-	-	-	-	-	-	-	-	-
	Pelagic	Northern gannet	Wintering			100S			-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Scoters	Wintering			10,000S			-	-	-	-	-	-	-	-	-	-	-
1169	Watrfowl	American black duck	Nesting			Moderate-High			Mar-Aug	-	-	-	-	-	-	-	-	-	-

FISH

Map ID	Subelement	Species	Mapping Qualifier			S	F	Concentration	Monthly Presence										
			J	F	M				J	F	M	A	M	J	J	A	S	O	N
1681	Diadromous	Alewife	Nursery Area			-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Diadromous	Blueback herring	Nursery Area			-	-	-	-	-	-	-	-	-	-	-	-	-	-

REPTILES & AMPHIBIANS


Map ID	Subelement	Species	MD/VA			Monthly Presence												Monthly Presence					
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles	Adults
1781	Turtle	Diamondback terrapin	Nesting			Possible Nest													May-Aug	-	-	Apr-Oct	Apr-Oct
1782	Turtle	Diamondback terrapin	Nesting			Probable Nest													May-Aug	-	-	Apr-Oct	Apr-Oct

MARINE MAMMALS

Map ID	Subelement	Species	MD/VA			Monthly Presence												Mating	Calving	Pupping	Molt
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping
1849	Whale	Humpback whale	General Distribution	E/E	E	Very Rare												-	-	-	-

WIDESpread IN MAPPED AREA (> 10 SQUARE KILOMETERS)

BIRDS



BIRDS (continued)

Subelement	Species	MDVA			Monthly Presence																	
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt		
	Snow goose	Wintering		1,000S													-	-	-	-		
	Tundra swan	Wintering		100S													-	-	-	-		
	Wood duck	Nesting		Moderate-High													Mar-Jun	-	Jul-Aug	-		

FISH

Subelement	Species	MDVA			Monthly Presence																	
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults	
	Alewife	Migration		-													-	-	Aug-Nov	Feb-Apr		
	Alewife	Migration		-													-	-	Aug-Nov	Mar-May		
	American shad	Migration		-													-	-	Aug-Nov	Mar-Jun		
	American shad	General Distribution	-IE	E													-	-	Aug-Nov	Feb-Jun		
	Atlantic sturgeon	Migration		-													-	-	Aug-Nov	Jan-Dec		
	Blueback herring	Migration		-													-	-	Aug-Nov	Feb-Jun		
	Blueback herring	Migration		-													-	-	Aug-Nov	Feb-May		
	Hickory shad	Migration		-													-	-	Jun-Oct	Feb-Jun		
	Hickory shad	General Distribution	E/E	E													-	-	Jan-Dec	Jan-Dec		
	Shortnose sturgeon	Migration		-													-	-	Mar-Jun	Feb-Jun		
	Striped bass	Migration		-													-	-	Feb-Jun	Feb-Jun		
	Striped bass	Migration		-													-	-	Feb-Jun	Feb-Jun		

REPTILES & AMPHIBIANS



Subelement	Species	MDVA			Monthly Presence																	
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Interrest	Juveniles	Adults	
	K. ridley sea turtle	General Distribution	E/E	E													-	-	Nov-Dec	-		
	K. ridley sea turtle	General Distribution	E/E	E													-	-	Apr-Oct	-		
	Loggerhead sea turtle	General Distribution	T/T	T													-	-	Apr-Oct	Apr-Oct		
	Loggerhead sea turtle	General Distribution	T/T	T													-	-	Nov-Dec	Nov-Dec		

INVERTEBRATES

Subelement	Species	MDVA			Monthly Presence																	
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults	
	Concentration Area	Highly Abundant															-	-	-	Jun-Mar*		
	Concentration Area	Highly Abundant-Summ															-	-	-	Jun-Oct		

MARINE MAMMALS

Subelement	Species	MDVA			Monthly Presence																	
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping	Molt		
	General Distribution	-IE	E	Rare													-	-	-	-		
	West Indian manatee																-	-	-	-		

BIRDS**FISH**

FISH (continued)

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Red hake	-	-	-	-	-	-	-	-	-	-	-	-	Sep-May*
	Silver hake	-	-	-	-	-	-	-	-	-	-	-	-	Oct-May
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Sep
	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Nov
	Cobia	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Lined seahorse	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Spanish mackerel	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		-	-	-	-	-	-	-	-	-	-	-	-	May-Sep

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Hard clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Stout razor clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Squid	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Horseshoe crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Brown shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Mantis shrimp	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	White shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct

MARINE MAMMALS

Subelement	Species	Monthly Presence												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-
	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-
	Harbor porpoise	-	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov


SHORELINE RESOURCES

ESI POLYGON HABITAT TYPES		Habitat Classification	Area (Acres)	Area (Sq. Miles)
	ESI Rank	Salt and Brackish Water Marshes	3,598.63	5.62
	10A	Scrub and Shrub Wetlands	18.97	0.03
	10D	Exposed Tidal Flats	1.64	0.00
ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification	Length (Meters)	% of ESI Shoreline
	ESI Rank	Salt and Brackish Water Marshes	92,029.18	57.18
	10A	Scrub and Shrub Wetlands	1,358.20	0.84
	10D	Vegetated Low Banks	328.75	0.20
	9B	Sheltered, Solid Man-Made Structures	33.81	< 1%
	8B	Sheltered Riprap	522.83	0.02
	8C	Exposed Tidal Flats	298.17	0.32
	7	Mixed Sand and Gravel Beaches	74.42	0.19
	5	Coarse Grained Sand Beaches	339.96	< 1%
	4			< 1%
	Total ESI Shoreline:		94,985.32	59.02
	Total Shoreline:		92,235.05	Total ESI Shoreline: 57.31

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%.

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 57C Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	Mapping Qualifier	MDVA	S	F	Concentration	Monthly Presence
SAV	Submerged aquatic veg	High Ecological Value				Ephemeral	-
SAV	Submerged aquatic veg	High Ecological Value				High	-
SAV	Submerged aquatic veg	High Ecological Value				Low	-

BIRDS

Map ID	Subelement	Species	Mapping Qualifier	MDVA	S	F	Concentration	Monthly Presence
33	Shorebird	American oystercatcher	Nesting			1 Pair		Apr-Aug
98	Gull/Tern	G. black-backed gull	Nesting			40 Pairs		Apr-Jun
	Gull/Tern	Herring gull	Nesting			220 Pairs		May-Aug
	Wading	Great egret	Nesting			29 Nests		Apr-Aug
	Wading	Snowy egret	Nesting			30 Pairs		Apr-Aug
	Wading	Tricolored heron	Nesting			1 Pair		May-Aug
99	Shorebird	American oystercatcher	Nesting			14 Pairs		Apr-Aug
100	Gull/Tern	Common tern	Nesting			2 Pairs		May-Aug
	Gull/Tern	Forster's tern	Nesting			19 Pairs		May-Jul
136	Gull/Tern	G. black-backed gull	Nesting			38 Pairs		Apr-Jun
	Gull/Tern	Herring gull	Nesting			37 Pairs		May-Aug
	Wading	Great egret	Nesting			3 Nests		Apr-Aug
	Wading	Snowy egret	Nesting			1 Pair		Apr-Aug
142	Wading	Great blue heron	Nesting			44 Nests		Feb-Jun
	Wading	Great egret	Nesting			1 Nest		Apr-Aug
159	Shorebird	American oystercatcher	Nesting			2 Pairs		Apr-Aug
323	Shorebird	American oystercatcher	Nesting			7 Pairs		Apr-Aug
350	Gull/Tern	G. black-backed gull	Nesting			16 Pairs		Apr-Jun
	Gull/Tern	Herring gull	Nesting			6 Pairs		May-Aug
351	Gull/Tern	Forster's tern	Nesting			3 Pairs		May-Jul
798	Raptor	Bald eagle	Nesting			1 Nest		Dec-Jun
821	Watertowl	Bufflehead	Wintering			10S		-
	Watertowl	Canvasback	Wintering			100S		-
	Watertowl	Northern shoveler	Wintering			10S		-
	Watertowl	Ruddy duck	Wintering			100S		-
924	Wading	Black rail	Vulnerable Occurrence			E/E		May-Jul
937	Watertowl	Canvasback	Wintering			100S		-
	Watertowl	Redhead	Wintering			100S		-
	Watertowl	Ring-necked duck	Wintering			10S		-
	Watertowl	Ruddy duck	Wintering			100S		-

BIRDS (continued)

Map ID	Subelement	Species	MD/VA			Monthly Presence															
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
	Waterfowl	Scaup			Wintering													-	-	-	-
939	Waterfowl	Blue-winged teal			Migration													-	Feb-Apr	Aug-Oct	-
	Waterfowl	Canvasback			Wintering													-	-	-	-
	Waterfowl	Green-winged teal			Wintering													-	-	-	-
	Waterfowl	Ruddy duck			Wintering													-	-	-	-
1378	Gull/Tern	Common tern			Nesting												May-Aug	-	-	-	-
	Gull/Tern	Forster's tern			Nesting												May-Jul	-	-	-	-

REPTILES & AMPHIBIANS

Map ID	Subelement	Species	MD/VA			Monthly Presence															
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles
1767	Turtle	Green sea turtle			General Distribution	T/T	T										-	-	-	May-Dec*	-
	Turtle	Green sea turtle			General Distribution	T/T	T										-	-	-	Jul-Oct	-

INVERTEBRATES

Map ID	Subelement	Species	MD/VA			Monthly Presence															
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles
1833	Insect	NE beach tiger beetle			Vulnerable Occurrence	E/T	T	-									Jun-Aug	Jun-Aug	-	-	Jun-Aug

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

BIRDS

Subelement	Species	MD/VA			Monthly Presence																
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt	
• Diving	Common loon			Wintering													-	-	-	-	-
	Grebes			Wintering													-	-	-	-	-
• Pelagic	Red-throated loon			Wintering													-	-	-	-	-
• Waterfowl	Northern gannet			Wintering													Apr-Jul	-	-	-	-
	American black duck			Nesting													-	-	-	-	-
	American black duck			Wintering													-	-	-	-	-
	American wigeon			Wintering													1,000S	-	-	-	-
	Blue-winged teal			Migration													1,000S	-	-	Feb-Apr	Aug-Oct
	Blue-winged teal			Migration													1,000S	-	-	Feb-Apr	Aug-Oct
	Brant			Wintering													10S	-	-	-	-
	Bufflehead			Wintering													1,000S	-	-	-	-
	Canada goose			Nesting													Present	-	-	Mar-Aug	-
	Canada goose			Wintering													1,000S	-	-	-	Jun-Jul
	Canvasback			Wintering													1,000S	-	-	-	-
	Gadwall			Wintering													1,000S	-	-	-	-
	Gadwall			Wintering													1,000S	-	-	-	-
	Green-winged teal			Wintering													100S	-	-	-	-
	Green-winged teal			Wintering													1,000S	-	-	-	-

BIRDS (continued)

Map ID	Subelement	Species	MD/VA		Concentration	Monthly Presence											
			Mapping Qualifier	S F		J F M A M J J A S O N D	Nest	Mig.(S)	Mig.(F)	Molt							
		Long-tailed duck	Wintering		100S	-	-	-	-	-							
	Mallard	Mallard	Wintering	Nesting	Present	1,000S	-	-	-	-							Jul-Aug
	Mallard	Mallard	Wintering	Wintering	1,000S	-	-	-	-	-							-
	Mergansers	Mergansers	Wintering	Wintering	100S	-	-	-	-	-							-
	Northern pintail	Northern shoveler	Wintering	Wintering	10S	-	-	-	-	-							-
	Redhead	Redhead	Wintering	Wintering	Present	1,000S	-	-	-	-							-
	Redhead	Ring-necked duck	Wintering	Wintering	10,000S	-	-	-	-	-							-
	Ruddy duck	Ruddy duck	Wintering	Wintering	Present	10,000S	-	-	-	-							-
	Scaup	Scaup	Wintering	Wintering	1,000S	-	-	-	-	-							-
	Scaup	Scoters	Wintering	Wintering	100S	-	-	-	-	-							-
	Snow goose	Snow goose	Wintering	Wintering	1,000S	-	-	-	-	-							-
	Tundra swan	Tundra swan	Wintering	Wintering	100S	-	-	-	-	-							-

FISH

Subelement	Species	MD/VA		Concentration	Monthly Presence											
		Mapping Qualifier	S F		J F M A M J J A S O N D	Spawn	Eggs	Larvae	Juveniles	Adults						
	Alewife	Migration		-	-	-	-	-	-	-						
	American shad	Migration		-	-	-	-	-	-	-						
	Atlantic sturgeon	General Distribution	-IE	E	-	-	-	-	-	-						
	Blueback herring	Migration		-	-	-	-	-	-	-						
	Hickory shad	Migration		-	-	-	-	-	-	-						
	Shortnose sturgeon	General Distribution	E/E	E	-	-	-	-	-	-						
	Striped bass	Migration		-	-	-	-	-	-	-						

REPTILES & AMPHIBIANS

Subelement	Species	MD/VA		Concentration	Monthly Presence											
		Mapping Qualifier	S F		J F M A M J J A S O N D	Nest	Hatch	Internest	Juveniles	Adults						
	Diamondback terrapin	Concentration Area	High	-	-	-	-	-	-	-						
	K. ridley sea turtle	General Distribution	E/E	E	-	-	-	-	-	-						
	K. ridley sea turtle	General Distribution	E/E	E	Common	-	-	-	-	-						
	Loggerhead sea turtle	General Distribution	T/T	T	Abundant	-	-	-	-	-						
	Loggerhead sea turtle	General Distribution	T/T	T	Common	-	-	-	-	-						

INVERTEBRATES

Subelement	Species	MD/VA		Concentration	Monthly Presence											
		Mapping Qualifier	S F		J F M A M J J A S O N D	Spawn	Eggs	Larvae	Juveniles	Adults						
	Blue crab	Concentration Area	Highly Abundant-Summ	-	-	-	-	-	-	-						
	Blue crab	Concentration Area	Highly Abundant	-	-	-	-	-	-	-						

MARINE MAMMALS

Subelement	Species	Mapping Qualifier	MD/VA												Monthly Presence												
			S	F	Concentration			J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping	Molt				
Manatee	West Indian manatee	General Distribution	-E	E	Rare			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)

BIRDS

Subelement	Species	Monthly Presence											
		J	F	M	A	M	J	J	A	S	O	N	D
Diving	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-
	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-
Gulls	Gulls	-	-	-	-	-	-	-	-	-	-	-	-
Gull/Tern	Terns	-	-	-	-	-	-	-	-	-	-	-	-
	Saltmarsh sharp-tailed sparrow	-	-	-	-	-	-	-	-	-	-	-	-
Passerine	Seaside sparrow	-	-	-	-	-	-	-	-	-	-	-	-
Shorebird	Willet	-	-	-	-	-	-	-	-	-	-	-	-
Wading	Clapper rail	-	-	-	-	-	-	-	-	-	-	-	-
Watertowl	Wood duck	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Subelement	Species	Monthly Presence											
		J	F	M	A	M	J	J	A	S	O	N	D
Diadromous	Alewife	-	-	-	-	-	-	-	-	-	-	-	-
	American eel	-	-	-	-	-	-	-	-	-	-	-	-
	Blueback herring	-	-	-	-	-	-	-	-	-	-	-	-
	Gizzard shad	-	-	-	-	-	-	-	-	-	-	-	-
	Striped bass	-	-	-	-	-	-	-	-	-	-	-	-
Estuarine Nursery	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-
	Atlantic menhaden	-	-	-	-	-	-	-	-	-	-	-	-
	Black drum	-	-	-	-	-	-	-	-	-	-	-	-
	Black sea bass	-	-	-	-	-	-	-	-	-	-	-	-
	Bluefish	-	-	-	-	-	-	-	-	-	-	-	-
	Kingfishes	-	-	-	-	-	-	-	-	-	-	-	-
	Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-
	Pigfish	-	-	-	-	-	-	-	-	-	-	-	-
	Red drum	-	-	-	-	-	-	-	-	-	-	-	-
	Sheepshead	-	-	-	-	-	-	-	-	-	-	-	-
	Spot	-	-	-	-	-	-	-	-	-	-	-	-
	Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-
	Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-
	Weakfish	-	-	-	-	-	-	-	-	-	-	-	-
	Winter flounder	-	-	-	-	-	-	-	-	-	-	-	-
Estuarine Resident	Pipefish	-	-	-	-	-	-	-	-	-	-	-	-
	Silversides	-	-	-	-	-	-	-	-	-	-	-	-
Fish	Anchoovies	-	-	-	-	-	-	-	-	-	-	-	-
	Striped burrfish	-	-	-	-	-	-	-	-	-	-	-	-

FISH (continued)

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Marine Benthic	Atlantic spadefish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Cleartooth skate	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Red hake	-	-	-	-	-	-	-	-	-	-	-	-	Sep-May*
	Silver hake	-	-	-	-	-	-	-	-	-	-	-	-	Oct-May
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Marine Pelagic	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Nov
	Cobia	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Lined seahorse	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Spanish mackerel	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		-	-	-	-	-	-	-	-	-	-	-	-	May-Sep

REPTILES & AMPHIBIANS

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Turtle	Diamondback terrapin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Bivalve	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Stout razor clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Cephalopod	Squid	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
Crab	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Horseshoe crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Shrimp	Brown shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Mantis shrimp	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	White shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct

MARINE MAMMALS

Subelement	Species	Monthly Presence												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
Dolphin	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-
	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-
	Harbor porpoise	-	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES

ESI POLYGON HABITAT TYPES		Habitat Classification	Area (Acres)	Area (Sq. Miles)
ESI Rank				
10A		Salt and Brackish Water Marshes	4,980.40	7.78
10B		Freshwater Marshes	836.44	1.31
10C		Swamps	1,324.44	2.07
10D		Scrub and Shrub Wetlands	463.42	0.72
7		Exposed Tidal Flats	161.26	0.25

ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
ESI Rank					
10A		Salt and Brackish Water Marshes	241,726.33	150.20	50%
10B		Freshwater Marshes	31.89	0.02	< 1%
10C		Swamps	11,134.94	6.92	2%
10D		Scrub and Shrub Wetlands	2,802.32	1.74	1%
9A		Sheltered Tidal Flats	185,960.90	115.55	39%
9B		Vegetated Low Banks	3,542.74	2.20	1%
8B		Sheltered, Solid Man-Made Structures	1,766.26	1.10	< 1%
8C		Sheltered Riprap	134.78	0.08	< 1%
7		Exposed Tidal Flats	13,799.66	8.57	3%
6B		Riprap	313.95	0.20	< 1%
5		Mixed Sand and Gravel Beaches	38.58	0.02	< 1%
3A		Fine to Medium Grained Sand Beaches	18,961.91	11.78	4%
1B		Exposed, Solid Man-Made Structures	57.71	0.04	< 1%
		Total ESI Shoreline:	480,271.98	Total ESI Shoreline:	298.43
		Total Shoreline:	246,796.63	Total Shoreline:	153.35

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov



75° 52' W

75° 50' W

75° 48' W

Map 57B

PRESENT THROUGHOUT
SALT AND BRACKISH
WATER MARSHES

37° 50' N

37° 50' N

POCOMOKE
SOUND

Map 58C

Map 57C

37° 48' N

37° 48' N

37° 46' N

37° 46' N

Map 64B

75° 48' W

Map 64A

75° 52' W

75° 50' W

75° 48' W

Map 57D Central Bay



SEE BACK OF MAP
for details about mapped species
and other species that occur in mapped area.
Data Published: September 2016

0 Not for Navigation 1 Miles

0 1 Kilometers

1:50,000



Map 57D Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	Mapping Qualifier	MDVA						Monthly Presence							
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N
SAV	Submerged aquatic veg	High Ecological Value	-	-	Ephemeral	-	-	-	-	-	-	-	-	-	-	-
SAV	Submerged aquatic veg	High Ecological Value	-	-	High	-	-	-	-	-	-	-	-	-	-	-
SAV	Submerged aquatic veg	High Ecological Value	-	-	Low	-	-	-	-	-	-	-	-	-	-	-

BIRDS

Map ID	Subelement	Species	Mapping Qualifier	MDVA						Monthly Presence							
				S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N
14	Watrfowl	American black duck	Nesting	-	-	Present	-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	American black duck	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Blue-winged teal	Migration	-	-	100S	-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Canada goose	Nesting	-	-	Present	-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Green-winged teal	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Mallard	Nesting	-	-	Present	-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Mallard	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-
33	Shorebird	American oystercatcher	Nesting	-	-	1 Nest	-	-	-	-	-	-	-	-	-	-	-
159	Shorebird	American oystercatcher	Nesting	-	-	1 Pair	-	-	-	-	-	-	-	-	-	-	-
347	Gull/Tern	Forster's tern	Nesting	-	-	2 Pairs	-	-	-	-	-	-	-	-	-	-	-
798	Raptor	Bald eagle	Nesting	-	-	228 Pairs	-	-	-	-	-	-	-	-	-	-	-
939	Watrfowl	Blue-winged teal	Migration	-	-	1 Nest	-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Canvasback	Wintering	-	-	10S	-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Green-winged teal	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Ruddy duck	Wintering	-	-	10S	-	-	-	-	-	-	-	-	-	-	-
1090	Diving	Common loon	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-
	Diving	D. crested cormorant	Migration	-	-	1,000S	-	-	-	-	-	-	-	-	-	-	-
	Diving	Grebes	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-
Pelagic		Northern gannet	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-
	Watrfowl	Scoters	Wintering	-	-	10,000S	-	-	-	-	-	-	-	-	-	-	-

REPTILES & AMPHIBIANS

Map ID	Subelement	Species	Mapping Qualifier	MDVA						Monthly Presence							
				S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N
1762	Turtle	Diamondback terrapin	Concentration Area	-	-	High	-	-	-	-	-	-	-	-	-	-	-

INVERTEBRATES

Map ID	Subelement	Species	Mapping Qualifier	MDVA						Monthly Presence								
				S	F	Concentration	E/T	T	Concentration	J	F	M	A	M	J	O	N	D
1833	Insect	NE beach tiger beetle	Vulnerable Occurrence	-	-	High	-	-	May-Aug	-	-	-	-	-	-	-	-	-

Central Bay: Map 57D

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

BIRDS

Subelement	Species	Mapping Qualifier	S	F	Concentration	Monthly Presence						Molt							
						J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)
Diving	Common loon	Wintering	-	-	Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Grebes	Wintering	-	-	Present	10S	-	-	-	-	-	-	-	-	-	-	-	-	-
	Red-throated loon	Wintering	-	-	Present	100S	-	-	-	-	-	-	-	-	-	-	-	-	-
Pelagic	Northern gannet	Wintering	-	-	Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Waterfowl	American black duck	Wintering	-	-	Present	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-
	American wigeon	Wintering	-	-	Present	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-
	American wigeon	Wintering	-	-	Present	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-
	Blue-winged teal	Migration	-	-	Present	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-
	Brant	Wintering	-	-	Present	10S	-	-	-	-	-	-	-	-	-	-	-	-	-
	Bufflehead	Wintering	-	-	Present	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-
	Canada goose	Wintering	-	-	Present	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-
	Canvasback	Wintering	-	-	Present	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-
	Gadwall	Wintering	-	-	Present	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-
	Gadwall	Wintering	-	-	Present	100S	-	-	-	-	-	-	-	-	-	-	-	-	-
	Green-winged teal	Wintering	-	-	Present	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-
	Long-tailed duck	Wintering	-	-	Present	100S	-	-	-	-	-	-	-	-	-	-	-	-	-
	Mallard	Wintering	-	-	Present	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-
	Mergansers	Wintering	-	-	Present	100S	-	-	-	-	-	-	-	-	-	-	-	-	-
	Northern pintail	Wintering	-	-	Present	10S	-	-	-	-	-	-	-	-	-	-	-	-	-
	Northern shoveler	Wintering	-	-	Present	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-
	Redhead	Wintering	-	-	Present	10,000S	-	-	-	-	-	-	-	-	-	-	-	-	-
	Redhead	Wintering	-	-	Present	10,000S	-	-	-	-	-	-	-	-	-	-	-	-	-
	Ring-necked duck	Wintering	-	-	Present	10,000S	-	-	-	-	-	-	-	-	-	-	-	-	-
	Ruddy duck	Wintering	-	-	Present	10,000S	-	-	-	-	-	-	-	-	-	-	-	-	-
	Scaup	Wintering	-	-	Present	10,000S	-	-	-	-	-	-	-	-	-	-	-	-	-
	Scaup	Wintering	-	-	Present	10,000S	-	-	-	-	-	-	-	-	-	-	-	-	-
	Scoters	Wintering	-	-	Present	10,000S	-	-	-	-	-	-	-	-	-	-	-	-	-
	Snow goose	Wintering	-	-	Present	100S	-	-	-	-	-	-	-	-	-	-	-	-	-
	Tundra swan	Wintering	-	-	Present	100S	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Subelement	Species	Mapping Qualifier	S	F	Concentration	Monthly Presence						Adults								
						J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae
Diadromous	Alewife	Migration	-	-	Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	American shad	Migration	-	-	Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Atlantic sturgeon	General Distribution	-E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Blueback herring	Migration	-	-	Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Hickory shad	Migration	-	-	Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Shortnose sturgeon	General Distribution	E/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Striped bass	Migration	-	-	Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

REPTILES & AMPHIBIANS

		Monthly Presence																				
Subelement	Species	Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles	Adults
Turtle	Green sea turtle	General Distribution	T/T	T	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Oct	-
	Green sea turtle	General Distribution	T/T	T	Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Dec*	-
K. ridley sea turtle	K. ridley sea turtle	General Distribution	E/E	E	Uncommon	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Dec	-
K. ridley sea turtle	General Distribution	E/E	E	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct	-
Loggerhead sea turtle	Loggerhead sea turtle	General Distribution	T/T	T	Abundant	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct	Apr-Oct
	General Distribution	T/T	T	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Dec	Nov-Dec

INVERTEBRATES

		Monthly Presence																				
Subelement	Species	Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
Crab	Blue crab	Concentration Area	-/E	E	Highly Abundant-Summ	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Oct	Jun-Mar*	
	Blue crab	Concentration Area	E/E	E	Highly Abundant	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Mar*	

MARINE MAMMALS

		Monthly Presence																			
Subelement	Species	Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping	Molt
Manatee	West Indian manatee	General Distribution	-/E	E	Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Whale	Humpback whale	General Distribution	E/E	E	Very Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)

BIRDS

		Monthly Presence																			
Subelement	Species	Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
Diving	Brown pelican	-/E	E	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Brown pelican	-/E	E	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	D. crested cormorant	-/E	E	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Gull/Tern	Gulls	-/E	E	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Terns	-/E	E	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Saltmarsh sharp-tailed sparrow	-/E	E	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Seaside sparrow	-/E	E	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Shorebird	Willet	-/E	E	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Clapper rail	-/E	E	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

		Monthly Presence																				
Subelement	Species	Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
Diadromous	Alewife	-/E	E	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Apr	Feb-Apr	Feb-Apr	Feb-Apr	
	American eel	-/E	E	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-May	
	Blueback herring	-/E	E	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct	
	Gizzard shad	-/E	E	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Jun	
	Striped bass	-/E	E	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	
Estuarine Nursery	Atlantic croaker	-/E	E	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov	

FISH (continued)

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Atlantic menhaden	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Black drum	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Black sea bass	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Bluefish	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Kingfishes	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Oct
	Pigfish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Red drum	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Sheepshead	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Spot	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Apr
	Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-	Feb-May
	Weakfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Aug
	Winter flounder	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Dec
	Estuarine Resident	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Fish	-	-	-	-	-	-	-	-	-	-	-	-	Dec-Mar
	Marine Benthic	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Pipefish	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Silversides	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Anchovies	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Striped burrfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Atlantic spadefish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Cleancorse skate	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Red hake	-	-	-	-	-	-	-	-	-	-	-	-	Sep-May*
	Silver hake	-	-	-	-	-	-	-	-	-	-	-	-	Oct-May
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Marine Pelagic	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Nov
	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Cobia	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Lined seahorse	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Spanish mackerel	-	-	-	-	-	-	-	-	-	-	-	-	May-Sep

REPTILES & AMPHIBIANS

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Turtle	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct
	Diamondback terrapin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Bivalve	Eastern oyster	■	■	■	■	■	■	■	■	■	■	■	■	Jan-Dec
	Hard clam	■	■	■	■	■	■	■	■	■	■	■	■	Jan-Dec
	Softshell clam													Jan-Dec
	Stout razor clam													Jan-Dec
Cephalopod	Squid													Jan-Dec
	Blue crab	■	■	■	■	■	■	■	■	■	■	■	■	Apr-Dec
	Horseshoe crab													Jan-Dec
	Brown shrimp													Jan-Dec
	Mantis shrimp													May-Oct
	White shrimp													May-Oct

MARINE MAMMALS

Subelement	Species	Monthly Presence												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
Dolphin	Bottlenose dolphin	■	■	■	■	■	■	■	■	■	■	■	■	-
	Bottlenose dolphin													-
	Harbor porpoise													-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES

ESI Rank	Habitat Classification	Area (Sq. Miles)												% of ESI Shoreline
		J	F	M	A	M	J	J	A	S	O	N	D	
10A	Salt and Brackish Water Marshes	1,500.28												51%
10D	Scrub and Shrub Wetlands													44%
9A	Sheltered, Solid Man-Made Structures													< 1%
7	Exposed Tidal Flats													3%
7	Fine to Medium Grained Sand Beaches													3%

ESI SHORELINE HABITAT TYPES


ESI Rank	Shoreline Habitat Classification	Length (Meters)												% of ESI Shoreline
		J	F	M	A	M	J	J	A	S	O	N	D	
10A	Salt and Brackish Water Marshes	145,091.13												51%
9A	Sheltered Tidal Flats	125,007.18												44%
8B	Sheltered, Solid Man-Made Structures													< 1%
7	Exposed Tidal Flats													3%
3A	Fine to Medium Grained Sand Beaches													3%

Total ESI Shoreline: 285,053.41 Total ESI Shoreline: 177.12
Total Shoreline: 147,010.39 Total Shoreline: 91.35

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 58
Central Bay



Map 58
Central Bay

HUMAN USE RESOURCES

DISPLAYED ON MAP (POINTS)

<i>Map ID</i>	<i>Type</i>	<i>Name</i>	<i>Contact</i>	<i>Phone</i>
1	AIRPORT	TANGIER ISLAND		
2	AQUACULTURE	OYSTER NON-COMM CHES BAY FOUNDATION RF#98-1675	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
3	AQUACULTURE	OYSTER NON-COMM PARKS RF#98-1825	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
4	AQUACULTURE	OYSTER NON-COMM THOMAS RF#99-0915	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
5	ARTIFICIAL REEF	JANE'S ISLAND		
6	ARTIFICIAL REEF	TANGIER SOUND		
7	FERRY	TANGIER FERRY		
8	REPEATED MEASUREMENT SITE	WQ - SHELLFISH HARVEST WATERS, CHESAPEAKE BAY	MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
9	REPEATED MEASUREMENT SITE	WQ - SHELLFISH HARVEST WATERS, LITTLE ANNEMASSEX	MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
10	REPEATED MEASUREMENT SITE	WQ - SHELLFISH HARVEST WATERS, TANGER SOUND	MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
11	REPEATED MEASUREMENT SITE	WQ - STATION CB7.1N	CHESAPEAKE BAY PROGRAM	800-968-7229

DISPLAYED ON MAP (POLYGONS)

<i>Map ID</i>	<i>Type</i>	<i>Name</i>	<i>Contact</i>	<i>Phone</i>
12	MANAGEMENT AREA	CEDAR ISLAND WMA	WILDLIFE AND HERITAGE SERVICE	
13	WILDLIFE REFUGE	MARTIN NATIONAL WILDLIFE REFUGE	UNITED STATES FISH AND WILDLIFE SERVICE	410-228-2692

ALSO PRESENT IN MAPPED AREA (POLYGONS)

<i>Type</i>	<i>Name</i>	<i>Contact</i>	<i>Phone</i>
STATE PROTECTED AREA	BLUE CRAB SANCTUARY	VIRGINIA MARINE RESOURCES COMMISSION	757-247-2200
STATE PROTECTED AREA	TIDAL LANDS	VA MARINE RESOURCES COMMISSION	

JURISDICTIONS

COUNTY: ACCOMACK COUNTY, SOMERSET COUNTY
COAST GUARD: DISTRICT 5, SECTOR BALTIMORE, SECTOR HAMPTON ROADS
USACE: NORTH ATLANTIC DIVISION, BALTIMORE DISTRICT, NORFOLK DISTRICT

FEMA:
REGION I
REGION 3

EPA:
REGION 3

SHORELINE RESOURCES**ESI POLYGON HABITAT TYPES**

ESI Rank	Habitat Classification	Area (Acres)	Area (Sq. Miles)
10A	Salt and Brackish Water Marshes	3,187.33	4.98
10B	Freshwater Marshes	15.57	0.02
10D	Scrub and Shrub Wetlands	25.51	0.04
9A	Sheltered Tidal Flats	3.93	0.01
7	Exposed Tidal Flats	278.14	0.43

ESI SHORELINE HABITAT TYPES


ESI Rank	Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
10A	Salt and Brackish Water Marshes	202,840.79	126.04	59%
10D	Scrub and Shrub Wetlands	7,281.05	4.52	2%
9A	Sheltered Tidal Flats	5,568.01	3.46	2%
9B	Vegetated Low Banks	39.31	0.02	< 1%
8B	Sheltered, Solid Man-Made Structures	2,734.48	1.70	1%
8C	Sheltered Riprap	922.97	0.57	< 1%
7	Exposed Tidal Flats	88,331.47	54.89	26%
6A	Gravel Beaches	191.27	0.12	< 1%
6B	Riprap	2,725.85	1.69	1%
4	Coarse Grained Sand Beaches	5,655.39	3.51	2%
3A	Fine to Medium Grained Sand Beaches	27,231.82	16.92	8%
1B	Exposed, Solid Man-Made Structures	278.93	0.17	< 1%

Total ESI Shoreline: 343,801.34 Total ESI Shoreline: 213.63
 Total Shoreline: 215,616.07 Total Shoreline: 133.98

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%.

All underlying GIS data can be obtained from response.restoration.noaa.gov





SEE BACK OF MAP
for details about mapped species and
other species that occur in mapped area.
Data Published: September 2016

Map 58A Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	Mapping Qualifier			S	F	Concentration	Monthly Presence										
		High Ecological Value	High Ecological Value	High Ecological Value				J	F	M	A	M	J	J	A	S	O	N
SAV	Submerged aquatic veg							-	-	-	-	-	-	-	-	-	-	-
SAV	Submerged aquatic veg							-	-	-	-	-	-	-	-	-	-	-
SAV	Submerged aquatic veg							-	-	-	-	-	-	-	-	-	-	-

BIRDS

Map ID	Subelement	Species	Mapping Qualifier			S	F	Concentration	Monthly Presence										
			J	F	M				J	F	M	A	M	J	J	A	S	O	N
1	Waterfowl	American black duck	Nesting			Present			Present										
	Waterfowl	Canada goose	Nesting			Present			Present										
	Waterfowl	Mallard	Nesting			Present			Present										
2	Shorebird	American oystercatcher	Nesting			2 Pairs			Present										
	Waterfowl	American black duck	Nesting			Present			Present										
	Waterfowl	Canada goose	Nesting			Present			Present										
	Waterfowl	Mallard	Nesting			Present			Present										
85	Gull/Tern	Forster's tern	Nesting			Present			Present										
	Gull/Tern	Laughing gull	Nesting			Present			Present										
209	Shorebird	American oystercatcher	Nesting			Present			Present										
336	Gull/Tern	Black skimmer	Nesting	E/-		70 Pairs			70 Pairs										
	Gull/Tern	Common tern	Nesting			40 Pairs			40 Pairs										
	Gull/Tern	Forster's tern	Nesting			60 Pairs			60 Pairs										
	Gull/Tern	G. black-backed gull	Nesting			13 Pairs			13 Pairs										
	Gull/Tern	Laughing gull	Nesting			87 Pairs			87 Pairs										
	Gull/Tern	Royal tern	Nesting	E/-		40 Pairs			40 Pairs										
401	Waterfowl	Brant	Wintering			100S			100S										
	Waterfowl	Canada goose	Wintering			1,000S			1,000S										
	Waterfowl	Canvasback	Wintering			100S			100S										
	Waterfowl	Redhead	Wintering			10,000S			10,000S										
	Waterfowl	Ruddy duck	Wintering			10S			10S										
	Waterfowl	Scaup	Wintering			1,000S			1,000S										
	Waterfowl	Tundra swan	Wintering			1,000S			1,000S										
406	Waterfowl	American black duck	Wintering			1,000S			1,000S										
	Waterfowl	American wigeon	Wintering			1,000S			1,000S										
	Waterfowl	Blue-winged teal	Migration			100S			100S										
	Waterfowl	Brant	Wintering			100S			100S										
	Waterfowl	Canada goose	Wintering			1,000S			1,000S										
	Waterfowl	Gadwall	Wintering			1,000S			1,000S										
	Waterfowl	Green-winged teal	Wintering			100S			100S										

BIRDS (continued)

Map ID	Subelement	Species	MD\VA			Monthly Presence										Nest		Mig.(S)		Mig.(F)		
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
	Waterfowl	Mallard	Wintering			100S													-	-	-	-
	Waterfowl	Northern pintail	Wintering			100S													-	-	-	-
	Waterfowl	Tundra swan	Wintering			1,000S													-	-	-	-
473	Waterfowl	American black duck	Nesting			Moderate-High													-	Mar-Aug	-	Jul-Aug
	Waterfowl	Canada goose	Nesting			Moderate-High													-	Mar-Jun	-	Jul-Aug
	Waterfowl	Wood duck	Nesting			Moderate-High													-	Mar-Jun	-	Jul-Aug
933	Waterfowl	Canvasback	Wintering			100S													-	-	-	-
	Waterfowl	Redhead	Wintering			1,000S													-	-	-	-
	Waterfowl	Ruddy duck	Wintering			100S													-	-	-	-
	Waterfowl	Scalp	Wintering			100S													-	-	-	-
1160	Diving	Common loon	Wintering			Present													-	-	-	-
	Diving	Grebes	Wintering			10S													-	-	-	-
	Pelagic	Northern gannet	Wintering			Present													-	-	-	-
	Waterfowl	Scoters	Wintering			1,000S													-	-	-	-

FISH

Map ID	Subelement	Species	MD\VA			Monthly Presence										Nest		Mig.(S)		Mig.(F)		
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles
1681	Diadromous	Alewife	Nursery Area			-												-	-	-	Apr-Jun	Apr-Oct
	Diadromous	Blueback herring	Nursery Area			-												-	-	-	May-Jul	May-Oct

INVERTEBRATES

Map ID	Subelement	Species	MD\VA			Monthly Presence										Nest		Mig.(S)		Mig.(F)			
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
1789	Crab	Blue crab	Concentration Area			Highly Abundant											-	-	-	-	Jun-Mar*	Jun-Mar*	
1827	Insect	Federally threatened insect 1	Vulnerable Occurrence	E/T	T	-												-	Jun-Aug	Jun-Jul	Jan-Dec	-	Jun-Aug

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

BIRDS

Map ID	Subelement	Species	MD\VA			Monthly Presence										Nest		Mig.(S)		Mig.(F)		
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
	Diving	Common loon	Wintering			100S												-	-	Mar-May	Aug-Nov	-
		D. crested cormorant	Migration			1,000S												-	-	-	-	-
		Grebies	Wintering			100S												-	-	-	-	-
		Red-throated loon	Wintering			100S												-	-	-	-	-
		Northern gannet	Wintering			1,000S												-	-	-	-	-
		American black duck	Wintering			10S												-	-	-	-	-
		American coot	Wintering			100S												-	-	-	-	-
		American wigeon	Wintering			1,000S												-	-	-	-	-
		Bufflehead	Wintering			1,000S												-	-	-	-	-
		Bufflehead	Wintering			1,000S												-	-	-	-	-
		Canada goose	Wintering			100S												-	-	-	-	-
		Canvasback	Wintering			1,000S												-	-	-	-	-

BIRDS (continued)

Map ID	Subelement	Species	Mapping Qualifier	Monthly Presence											
				M	D	V	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
	Canvasback	Wintering	Wintering	-	-	-	-	-	-	-	-	-	-	-	-
	Common goldeneye	Wintering	Wintering	-	-	-	-	-	-	-	-	-	-	-	-
	Gadwall	Wintering	Wintering	-	-	-	-	-	-	-	-	-	-	-	-
	Green-winged teal	Wintering	Wintering	-	-	-	-	-	-	-	-	-	-	-	-
	Long-tailed duck	Wintering	Wintering	-	-	-	-	-	-	-	-	-	-	-	-
	mallard	Wintering	Wintering	-	-	-	-	-	-	-	-	-	-	-	-
	Mergansers	Wintering	Wintering	-	-	-	-	-	-	-	-	-	-	-	-
	Mergansers	Wintering	Wintering	-	-	-	-	-	-	-	-	-	-	-	-
	Northern pintail	Wintering	Wintering	-	-	-	-	-	-	-	-	-	-	-	-
	Northern shoveler	Wintering	Wintering	-	-	-	-	-	-	-	-	-	-	-	-
	Redhead	Wintering	Wintering	-	-	-	-	-	-	-	-	-	-	-	-
	Redhead	Wintering	Wintering	-	-	-	-	-	-	-	-	-	-	-	-
	Ring-necked duck	Wintering	Wintering	-	-	-	-	-	-	-	-	-	-	-	-
	Ring-necked duck	Wintering	Wintering	-	-	-	-	-	-	-	-	-	-	-	-
	Ruddy duck	Wintering	Wintering	-	-	-	-	-	-	-	-	-	-	-	-
	Ruddy duck	Wintering	Wintering	-	-	-	-	-	-	-	-	-	-	-	-
	Scaup	Wintering	Wintering	-	-	-	-	-	-	-	-	-	-	-	-
	Scaup	Wintering	Wintering	-	-	-	-	-	-	-	-	-	-	-	-
	Scoters	Wintering	Wintering	-	-	-	-	-	-	-	-	-	-	-	-
	Snow goose	Wintering	Wintering	-	-	-	-	-	-	-	-	-	-	-	-
	Tundra swan	Wintering	Wintering	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Subelement	Species	Mapping Qualifier	Monthly Presence												
			M	D	V	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
Diadromous	Alewife	Migration	-	-	-	-	-	-	-	-	-	-	-	Feb-Nov	Feb-Apr
	Alewife	Migration	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Mar-May
	American shad	Migration	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Mar-Jun
	American shad	General Distribution	-E	-E	-	-	-	-	-	-	-	-	-	Aug-Nov	Jan-Dec
	Atlantic sturgeon	Migration	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Mar-Jun
	Blueback herring	Migration	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Feb-May
	Blueback herring	Migration	-	-	-	-	-	-	-	-	-	-	-	Jun-Oct	Feb-Jun
	Hickory shad	Migration	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec
	Hickory shad	General Distribution	-E	-E	-	-	-	-	-	-	-	-	-	Mar-Jun	Feb-Jun
	Shortnose sturgeon	Migration	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Striped bass	Migration	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Striped bass	Migration	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun

REPTILES & AMPHIBIANS

Subelement	Species	Mapping Qualifier	Monthly Presence												Adults
			J	F	M	A	M	J	J	A	S	O	N	D	
Turtle	K. ridley sea turtle	General Distribution	E/E	E	Concentration	-	-	-	-	-	-	-	-	-	Nov-Dec
	K. ridley sea turtle	General Distribution	E/E	E	Common	-	-	-	-	-	-	-	-	-	Apr-Oct
	Loggerhead sea turtle	General Distribution	T/T	T	Common	-	-	-	-	-	-	-	-	-	Nov-Dec
	Loggerhead sea turtle	General Distribution	T/T	T	Abundant	-	-	-	-	-	-	-	-	-	Apr-Oct

INVERTEBRATES

Subelement	Species	Mapping Qualifier	Monthly Presence												Adults
			J	F	M	A	M	J	J	A	S	O	N	D	
Crab	Blue crab	Concentration Area	-	-	Highly Abundant-Summ	-	-	-	-	-	-	-	-	-	Jun-Oct
			-	-	-	-	-	-	-	-	-	-	-	-	-

MARINE MAMMALS

Subelement	Species	Mapping Qualifier	Monthly Presence												Adults
			J	F	M	A	M	J	J	A	S	O	N	D	
Manatee	West Indian manatee	General Distribution	-/E	E	Rare	-	-	-	-	-	-	-	-	-	-
Whale	Humpback whale	General Distribution	E/E	E	Very Rare	-	-	-	-	-	-	-	-	-	-

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)

BIRDS

Subelement	Species	Mapping Qualifier	Monthly Presence												Adults
			J	F	M	A	M	J	J	A	S	O	N	D	
Diving	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Gull/Tern	Gulls	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Terns	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Saltmarsh sharp-tailed sparrow	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Passerine	Seaside sparrow	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Shorebird	Willet	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Clapper rail	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Subelement	Species	Mapping Qualifier	Monthly Presence												Adults
			J	F	M	A	M	J	J	A	S	O	N	D	
Diadromous	Alewife	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct
	American eel	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-May
	Blueback herring	-	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Gizzard shad	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Striped bass	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Estuarine Nursery	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Atlantic menhaden	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Black drum	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Black sea bass	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov

FISH (continued)

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Bluefish	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Kingfishes	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Pigfish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Red drum	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Sheepshead	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Spot	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Weakfish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	White perch	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Winter flounder	-	-	-	-	-	-	-	-	-	-	-	-	Dec-Mar
 Estuarine Resident	Pipefish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
 Fish	Silversides	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Anchoovies	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Striped burrfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
 Marine Benthic	Atlantic spadefish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Cleinnose skate	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Red hake	-	-	-	-	-	-	-	-	-	-	-	-	Sep-May*
	Silver hake	-	-	-	-	-	-	-	-	-	-	-	-	Oct-May
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
 Marine Pelagic	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Sep
	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Cobia	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Lined seahorse	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Spanish mackerel	-	-	-	-	-	-	-	-	-	-	-	-	May-Sep

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
 Bivalve	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Hard clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Stout razor clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
 Cephalopod	Squid	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Horseshoe crab	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
 Shrimp	Brown shrimp	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Mantis shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	White shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct

MARINE MAMMALS

Subelement	Species	Monthly Presence											
		J	F	M	A	M	J	J	A	S	O	N	D
Dolphin	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-
	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-
	Harbor porpoise	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES

ESI POLYGON HABITAT TYPES		Habitat Classification		Area (Acres)		Area (Sq. Miles)	
ESI Rank	10A	Salt and Brackish Water Marshes		1,131.59	1.77		
	9A	Sheltered Tidal Flats		1.15	0.00		
	7	Exposed Tidal Flats		88.89	0.14		


ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification		Length (Meters)		Length (Miles)	
ESI Rank	10A	Salt and Brackish Water Marshes		61,245.89	38.06		
	9A	Sheltered Tidal Flats		3,897.07	2.42		
	7	Exposed Tidal Flats		32,495.58	20.19		
	4	Coarse Grained Sand Beaches		5,655.39	3.51		
	3A	Fine to Medium Grained Sand Beaches		3,214.71	2.00		

Total ESI Shoreline:	106,508.65	Total ESI Shoreline:	66.18
Total Shoreline:	61,669.16	Total Shoreline:	38.32

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 58B Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	Mapping Qualifier	MD/VA	S	F	Concentration	Monthly Presence
SAV	Submerged aquatic veg	High Ecological Value	J F M A M J J A S O N D	-	-	-	-
SAV	Submerged aquatic veg	High Ecological Value					

BIRDS

Map ID	Subelement	Species	Mapping Qualifier	MD/VA	S	F	Concentration	Monthly Presence
			Nesting	J F M A M J J A S O N D	Nest	Mig.(S)	Mig.(F)	Molt
1	Waterfowl	American black duck	Nesting	Present	Present	Present	Present	-
	Waterfowl	Canada goose	Nesting	Present	Present	Present	Present	-
	Waterfowl	Mallard	Nesting	Present	Present	Present	Present	-
33	Shorebird	American oystercatcher	Nesting	1 Pair	22 Pairs	26 Pairs	26 Pairs	-
86	Gull/Tern	G. black-backed gull	Nesting	1 Pair	22 Pairs	26 Pairs	26 Pairs	-
	Gull/Tern	Herring gull	Nesting	1 Pair	22 Pairs	26 Pairs	26 Pairs	-
153	Wading	Great blue heron	Nesting	4 Nests	4 Nests	4 Nests	4 Nests	-
309	Shorebird	American oystercatcher	Nesting	12 Pairs	12 Pairs	12 Pairs	12 Pairs	-
	Shorebird	American oystercatcher	Roosting	3 Pairs	3 Pairs	3 Pairs	3 Pairs	-
337	Gull/Tern	Common tern	Nesting	1 Pair	1 Pair	1 Pair	1 Pair	-
	Gull/Tern	Forster's tern	Nesting	6 Pairs	6 Pairs	6 Pairs	6 Pairs	-
	Gull/Tern	G. black-backed gull	Nesting	27 Pairs	27 Pairs	27 Pairs	27 Pairs	-
	Gull/Tern	Herring gull	Nesting	4 Pairs	4 Pairs	4 Pairs	4 Pairs	-
352	Gull/Tern	G. black-backed gull	Nesting	5 Pairs	5 Pairs	5 Pairs	5 Pairs	-
374	Diving	Brown pelican	Nesting	1128 Pairs	1128 Pairs	1128 Pairs	1128 Pairs	-
	Diving	D. crested cormorant	Nesting	1026 Pairs	1026 Pairs	1026 Pairs	1026 Pairs	-
	Gull/Tern	G. black-backed gull	Nesting	3 Pairs	3 Pairs	3 Pairs	3 Pairs	-
	Gull/Tern	Herring gull	Nesting	7 Pairs	7 Pairs	7 Pairs	7 Pairs	-
379	Diving	Brown pelican	Nesting	729 Pairs	729 Pairs	729 Pairs	729 Pairs	-
	Diving	D. crested cormorant	Nesting	1343 Pairs	1343 Pairs	1343 Pairs	1343 Pairs	-
	Gull/Tern	G. black-backed gull	Nesting	80 Pairs	80 Pairs	80 Pairs	80 Pairs	-
	Gull/Tern	Herring gull	Nesting	15 Pairs	15 Pairs	15 Pairs	15 Pairs	-
418	Raptor	Bald eagle	Nesting	Last Obs. 2004	Very High	Moderate-High	Moderate-High	-
483	Waterfowl	American black duck	Nesting	4 Pairs	4 Pairs	4 Pairs	4 Pairs	-
	Waterfowl	Mallard	Nesting	1,000S	1,000S	1,000S	1,000S	-
493	Wading	Great blue heron	Nesting	100S	100S	100S	100S	-
565	Waterfowl	American black duck	Wintering	100S	100S	100S	100S	-
	Waterfowl	American wigeon	Wintering	100S	100S	100S	100S	-
	Waterfowl	Gadwall	Wintering	100S	100S	100S	100S	-
	Waterfowl	Green-winged teal	Wintering	100S	100S	100S	100S	-
	Waterfowl	Mallard	Wintering	100S	100S	100S	100S	-

BIRDS (continued)

Map ID	Subelement	Species	MD/VA			Monthly Presence										Monthly Presence			Molt		
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
Waterfowl		Northern pintail	Wintering		100S												-	-	-	-	-
1292	Gull/Tern	Herring gull	Nesting		2 Pairs												May-Aug	-	-	-	-
1343	Gull/Tern	Common tern	Nesting		71 Pairs												May-Aug	-	-	-	-
	Gull/Tern	Forster's tern	Nesting		642 Pairs												May-Jul	-	-	-	-
	Gull/Tern	G. black-backed gull	Nesting		19 Pairs												Apr-Jun	-	-	-	-

MARINE MAMMALS

Map ID	Subelement	Species	MD/VA			Monthly Presence										Monthly Presence			Molt		
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
1849	Whale	Humpback whale	General Distribution	E/E	Very Rare												-	-	-	-	-

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

BENTHIC

Subelement	Species	MD/VA			Monthly Presence										Monthly Presence			Molt		
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
SAV	Submerged aquatic veg	High Ecological Value															-	-	-	-

BIRDS

Subelement	Species	MD/VA			Monthly Presence										Monthly Presence			Molt		
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
Diving	Common loon	Wintering		100S													-	-	-	-
	Common loon	Wintering		100S													-	-	-	-
	D. crested cormorant	Migration		1,000S													-	Mar-May	Aug-Nov	-
	D. crested cormorant	Wintering		1,000S													-	Mar-May	Aug-Nov	-
	Grebes	Wintering		100S													-	-	-	-
	Grebes	Wintering		100S													-	-	-	-
	Red-throated loon	Wintering		100S													-	-	-	-
Pelagic	Northern gannet	Wintering		100S													-	-	-	-
Waterfowl	American black duck	Wintering		100S													-	-	-	-
	American wigeon	Wintering		1,000S													-	Feb-Apr	Aug-Oct	-
	Blue-winged teal	Migration		100S													-	-	-	-
	Brant	Wintering		100S													-	-	-	-
	Brant	Wintering		1,000S													-	-	-	-
	Bufflehead	Wintering		1,000S													-	-	-	-
	Bufflehead	Wintering		1,000S													-	-	-	-
	Canada goose	Wintering		1,000S													-	-	-	-
	Canvasback	Wintering		1,000S													-	-	-	-
	Canvasback	Wintering		1,000S													-	-	-	-
	Common goldeneye	Wintering		100S													-	-	-	-
	Gadwall	Wintering		1,000S													-	-	-	-
	Green-winged teal	Wintering		100S													-	-	-	-
	Long-tailed duck	Wintering		100S													-	-	-	-

BIRDS (continued)

Subelement	Species	MD/VA												Monthly Presence											
		Mapping Qualifier			S	F	Concentration			J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
	Mallard	Wintering					100S			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Mergansers	Wintering					100S			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Mergansers	Wintering					100S			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Northern pintail	Wintering					100S			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Redhead	Wintering					1,000S			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Redhead	Wintering					10,000S			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Ring-necked duck	Wintering					10S			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Ring-necked duck	Wintering					Wintering			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Ruddy duck	Wintering					100S			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Ruddy duck	Wintering					10S			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Ruddy duck	Wintering					Wintering			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Scaup	Wintering					Wintering			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Scaup	Wintering					100S			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Scooter's	Wintering					1,000S			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Tundra swan	Wintering					10,000S			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Subelement	Species	MD/VA												Monthly Presence												
		Mapping Qualifier			S	F	Concentration			J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
	Aldiromous	Migration					-			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-May
	Alewife	Migration					Migration			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov
	American shad	Migration					General Distribution	-E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Apr
	Atlantic sturgeon	Migration					Blueback herring			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Blueback herring	Migration					Hickory shad			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Jun
	Blueback herring	Migration					Hickory shad			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Shortnose sturgeon	General Distribution					Shortnose sturgeon	E/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Jun
	Striped bass	Migration					Striped bass			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
	Striped bass	Migration					Striped bass			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Jun

REPTILES & AMPHIBIANS

Subelement	Species	MD/VA												Monthly Presence												
		Mapping Qualifier			S	F	Concentration			J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles	Adults
	K. ridley sea turtle	General Distribution	E/E	E	Uncommon					-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Dec
	K. ridley sea turtle	General Distribution	E/E	E	Common					-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct
	Loggerhead sea turtle	General Distribution	T/T	T	Abundant					-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct
	Loggerhead sea turtle	General Distribution	T/T	T	Common					-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Dec

INVERTEBRATES

Subelement	Species	Mapping Qualifier	MD/V/A	S F	Concentration	J F M A M J J A S O N D	Spawning	Eggs	Larvae	Juveniles	Adults
Crab	Blue crab	Concentration Area			Highly Abundant-Summer				-	Jun-Oct	Jun-Oct
MARINE MAMMALS											
Subelement	Species	Mapping Qualifier	MD/V/A	S F	Concentration	J F M A M J J A S O N D	Mating	Calving	Pupping	Molt	
Manatee	West Indian manatee	General Distribution		-/E	E						-
ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)											

BIRDS

Subelement	Species	Mapping Qualifier	MD/V/A	S F	Concentration	J F M A M J J A S O N D	Nest	Mig.(S)	Mig.(F)	Molt
Diving	Brown pelican									-
	Brown pelican									-
	D. crested cormorant									-
	D. crested cormorant									-
Gull/Tern	Gulls									-
	Terns									-
Pelagic	Storm-petrels									-

FISH

Subelement	Species	Mapping Qualifier	MD/V/A	S F	Concentration	J F M A M J J A S O N D	Spawning	Eggs	Larvae	Juveniles	Adults
Diadromous	American eel								-	Jan-May	Jan-Dec
	Gizzard shad								-	Jan-Dec	Jan-Dec
	Striped bass								-	Jan-Dec	Jan-Dec
Estuarine Nursery	Atlantic croaker								-	Dec-Apr	Jan-Dec
	Atlantic menhaden								-	Jan-Dec	Jan-Dec
	Black drum								-	Apr-Dec	Apr-Dec
	Black sea bass								-	Apr-Nov	Apr-Nov
	Bluefish								-	May-Oct	May-Oct
	Kingfishes								-	Apr-Nov	Apr-Nov
	Northern puffer								-	Mar-Oct	Mar-Nov
	Pigfish								-	Apr-Nov	Apr-Nov
	Red drum								-	May-Nov	May-Nov
	Sheepshead								-	Apr-Nov	Apr-Nov
	Spot								-	Nov-Apr	Apr-Dec
	Spotted seatrout								-	Apr-Nov	Apr-Nov
	Spotted seatrout								-	Aug-Dec	Apr-Dec
	Summer flounder								-	Feb-May	Apr-Dec
	Weakfish								-	May-Aug	Jun-Dec
	White perch								-	Jan-Dec	Jan-Dec
Estuarine Resident	Pipefish								-	Jan-Dec	Jan-Dec

FISH (continued)

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Anchovies	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Striped burrfish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Atlantic spadefish	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Cleinnose skate	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Red hake	-	-	-	-	-	-	-	-	-	-	-	-	Sep-May*
	Silver hake	-	-	-	-	-	-	-	-	-	-	-	-	Oct-May
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Sep
	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Cobia	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Lined seahorse	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Spanish mackerel	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Spanish mackerel	-	-	-	-	-	-	-	-	-	-	-	-	May-Sep

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Hard clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Stout razor clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Squid	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Horseshoe crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Brown shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Mantis shrimp	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	White shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct

MARINE MAMMALS

Subelement	Species	Monthly Presence												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-
	Harbor porpoise	-	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES

ESI POLYGON HABITAT TYPES		Habitat Classification	Area (Acres)	Area (Sq. Miles)
ESI Rank				
10A		Salt and Brackish Water Marshes	14,143.33	2.21
10D		Scrub and Shrub Wetlands	10,244.00	0.02
7		Exposed Tidal Flats	95,320.15	0.15


ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
ESI Rank					
10A		Salt and Brackish Water Marshes	96,871.88	60.19	71%
10D		Scrub and Shrub Wetlands	558.72	0.35	< 1%
9A		Sheltered Tidal Flats	346.65	0.22	< 1%
8B		Sheltered, Solid Man-Made Structures	472.44	0.29	< 1%
8C		Sheltered Riprap	852.07	0.53	1%
7		Exposed Tidal Flats	30,170.27	18.75	22%
6A		Gravel Beaches	191.27	0.12	< 1%
3A		Fine to Medium Grained Sand Beaches	7,545.21	4.69	6%

Total ESI Shoreline: 137,008.51 Total ESI Shoreline: 85.13
 Total Shoreline: 98,703.35 Total Shoreline: 61.33

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%.

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 58C Central Bay



SEE BACK OF MAP
for details about mapped species and
other species that occur in mapped area.
Data Published: September 2016

0 Not for Navigation 1 Miles
0 1 Kilometers 1:50,000



Map 58C Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	MDVA	S	F	Concentration	MDVA	S	F	Concentration	MDVA	S	F	Concentration	MDVA	S	F	Concentration	
SAV	Submerged aquatic veg	High Ecological Value	J	F	Ephemeral	High Ecological Value	M	A	J	High Ecological Value	J	F	M	A	M	J	J	OND
SAV	Submerged aquatic veg	High Ecological Value																

BIRDS

Map ID	Subelement	Species	Mapping Qualifier	S	F	Concentration	Monthly Presence											
							J	F	M	A	M	J	J	A	S	O	N	D
1	Waterfowl	American black duck	Nesting	Present	Present	Present												
	Waterfowl	Canada goose	Nesting															
	Waterfowl	Mallard	Nesting															
176	Shorebird	American oystercatcher	Nesting															
182	Gull/Tern	G. black-backed gull	Nesting															
	Wading	BC night-heron	Nesting															
	Wading	Cattle egret	Nesting															
	Wading	Glossy ibis	Nesting															
	Wading	Great blue heron	Nesting															
	Wading	Great egret	Nesting															
	Wading	Snowy egret	Nesting															
	Wading	Tricolored heron	Nesting															
	Wading	YC night-heron	Nesting															
404	Waterfowl	American black duck	Wintering															
	Waterfowl	American wigeon	Wintering															
	Waterfowl	Blue-winged teal	Migration															
	Waterfowl	Brant	Wintering															
	Waterfowl	Canada goose	Wintering															
	Waterfowl	Canvasback	Wintering															
	Waterfowl	Gadwall	Wintering															
	Waterfowl	Green-winged teal	Wintering															
	Waterfowl	Mallard	Wintering															
	Waterfowl	Northern pintail	Wintering															
	Waterfowl	Redhead	Wintering															
	Waterfowl	Ruddy duck	Wintering															
	Waterfowl	Scaup	Wintering															
	Waterfowl	Tundra swan	Wintering															
405	Waterfowl	American black duck	Wintering															
	Waterfowl	American wigeon	Wintering															
	Waterfowl	Blue-winged teal	Migration															
	Waterfowl	Brant	Wintering															

BIRDS (continued)

Map ID	Subelement	Species	Mapping Qualifier	MD/VA	Monthly Presence												Nest	Mig.(S)	Mig.(F)	Molt		
					S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D			
	Waterfowl	Canada goose	Wintering				1,000S													-	-	-
	Waterfowl	Canvasback	Wintering				100S													-	-	-
	Waterfowl	Gadwall	Wintering				1,000S													-	-	-
	Waterfowl	Green-winged teal	Wintering				100S													-	-	-
	Waterfowl	Mallard	Wintering				100S													-	-	-
	Waterfowl	Northern pintail	Wintering				100S													-	-	-
	Waterfowl	Redhead	Wintering				1,000S													-	-	-
	Waterfowl	Ruddy duck	Wintering				100S													-	-	-
	Waterfowl	Scaup	Wintering				100S													-	-	-
	Waterfowl	Tundra swan	Wintering				1,000S													-	-	-
406	Waterfowl	American black duck	Wintering				100S													-	-	-
	Waterfowl	American wigeon	Wintering				1,000S													-	-	-
	Waterfowl	Blue-winged teal	Migration				100S													-	-	-
	Waterfowl	Brant	Wintering				100S													-	-	-
	Waterfowl	Canada goose	Wintering				1,000S													-	-	-
	Waterfowl	Gadwall	Wintering				1,000S													-	-	-
	Waterfowl	Green-winged teal	Wintering				100S													-	-	-
	Waterfowl	Mallard	Wintering				100S													-	-	-
	Waterfowl	Northern pintail	Wintering				100S													-	-	-
	Waterfowl	Tundra swan	Wintering				1,000S													-	-	-
705	Raptor	Peregrine falcon	Nesting	-T																Feb-Jun	Dec-Jun	
798	Raptor	Bald eagle	Nesting	-																1 Nest		
933	Waterfowl	Canvasback	Wintering																	100S		
	Waterfowl	Redhead	Wintering																	1,000S		
	Waterfowl	Ruddy duck	Wintering																	100S		
	Waterfowl	Scaup	Wintering																	100S		
1160	Diving	Common loon	Wintering																	Present		
	Diving	Grebes	Wintering																	10S		
	Pelagic	Northern gannet	Wintering																	Present		
	Waterfowl	Scoters	Wintering																	1,000S		

REPTILES & AMPHIBIANS

Map ID	Subelement	Species	Mapping Qualifier	MD/VA	Monthly Presence												Nest	Hatch	Internest	Juveniles	Adults
					S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D		
1767	Turtle	Green sea turtle	General Distribution	T/T	T	T	Rare	-	-	-	-	-	-	-	-	-	-	-	May-Dec*	-	-
	Turtle	Green sea turtle	General Distribution	T/T	T	T	Common	-	-	-	-	-	-	-	-	-	-	-	Jul-Oct	-	-

WIDESpread IN MAPPED AREA (> 10 SQUARE KILOMETERS)

BIRDS

Subelement	Species	Mapping Qualifier	MDVA						Monthly Presence											
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mg.(S)	Mg.(F)
Diving	Common loon	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	D. crested cormorant	Migration	-	-	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Grebes	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Red-throated loon	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Pelagic	Northern gannet	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Waterfowl	Bufflehead	Wintering	-	-	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Canvasback	Wintering	-	-	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Long-tailed duck	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Mergansers	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Redhead	Wintering	-	-	10,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Ring-necked duck	Wintering	-	-	Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Ruddy duck	Wintering	-	-	10,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Scaup	Wintering	-	-	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Scooter's	Wintering	-	-	10,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Subelement	Species	Mapping Qualifier	MDVA						Monthly Presence											
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Eggs	Larvae	Juveniles
Diadromous	Alewife	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	American shad	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Atlantic sturgeon	General Distribution	-IE	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Blueback herring	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Hickory shad	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Shortnose sturgeon	General Distribution	E/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Striped bass	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

REPTILES & AMPHIBIANS

Subelement	Species	Mapping Qualifier	MDVA						Monthly Presence												
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles
Turtle	K. ridley sea turtle	General Distribution	E/E	E	Uncommon	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	K. ridley sea turtle	General Distribution	E/E	E	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Loggerhead sea turtle	General Distribution	T/T	T	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Loggerhead sea turtle	General Distribution	T/T	T	Abundant	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

INVERTEBRATES


Subelement	Species	Mapping Qualifier	MDVA						Monthly Presence												
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles
Crab	Blue crab	Concentration Area	Highly Abundant-Summ	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

MARINE MAMMALS


Subelement	Species	Mapping Qualifier	MD/VA												Monthly Presence												
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping	Molt						
Manatee	West Indian manatee	General Distribution	-E	E	Rare													-	-	-	-						
Whale	Humpback whale	General Distribution	E/E	E	Very Rare													-	-	-	-						

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)

BIRDS



FISH



FISH (continued)

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Red hake	-	-	-	-	-	-	-	-	-	-	-	-	Sep-May*
	Silver hake	-	-	-	-	-	-	-	-	-	-	-	-	Oct-May
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Nov
	Cobia	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Lined seahorse	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Spanish mackerel	-	-	-	-	-	-	-	-	-	-	-	-	May-Sep

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Bivalve	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Hard clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Stout razor clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Squid	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Cephalopod	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Crab	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Horseshoe crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Brown shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Mantis shrimp	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	White shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct

MARINE MAMMALS

Subelement	Species	Monthly Presence												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
	Dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-
	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-
	Harbor porpoise	-	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES


ESI POLYGON HABITAT TYPES		Habitat Classification	Area (Acres)	Area (Sq. Miles)
	ESI Rank	Salt and Brackish Water Marshes	19.50	0.03
	10A	Scrub and Shrub Wetlands	13.02	0.02
	10D	Exposed Tidal Flats	23.79	0.04

ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
	ESI Rank	Salt and Brackish Water Marshes	1,831.53	1.14	17%
	10A	Scrub and Shrub Wetlands	1,034.19	0.64	10%
	10D	Exposed Tidal Flats	3,946.67	2.45	37%
	7	Fine to Medium Grained Sand Beaches	3,946.67	2.45	37%
		Total ESI Shoreline:	10,759.06	Total ESI Shoreline: 6.69	
		Total Shoreline:	3,946.67	Total Shoreline: 2.45	

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%.

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 58D
Central Bay



SEE BACK OF MAP
for details about mapped species
and other species that occur in mapped area.
Data Published: September 2016

0 Not for Navigation 1 Miles
0 1 Kilometers
1:50,000



Map 58D Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	Mapping Qualifier						Concentration	Monthly Presence						
		S	F	MD/VNA	J	F	M	A	M	J	J	A	S	O	N
SAV	Submerged aquatic veg	High Ecological Value													
SAV	Submerged aquatic veg	High Ecological Value													
SAV	Submerged aquatic veg	High Ecological Value													

BIRDS

Map ID	Subelement	Species	Mapping Qualifier						Concentration	Monthly Presence					
			S	F	MD/VNA	J	F	M	A	M	J	J	A	S	O
1	Waterfowl	American black duck	Nesting						Present						
	Waterfowl	Canada goose	Nesting						Present						
	Waterfowl	Mallard	Nesting						Present						
287	Gull/Tern	Black skimmer	Nesting						E/-	56 Pairs					
	Gull/Tern	Common tern	Nesting							17 Pairs					
	Gull/Tern	Herring gull	Nesting							9 Pairs					
	Gull/Tern	Royal tern	Nesting							E/-	16 Pairs				
300	Wading	BC night-heron	Nesting								24 Pairs				
	Wading	Glossy ibis	Nesting								26 Pairs				
	Wading	Great egret	Nesting								9 Nests				
	Wading	Little blue heron	Nesting								28 Pairs				
	Wading	Snowy egret	Nesting								34 Pairs				
	Wading	Tricolored heron	Nesting								8 Pairs				
	Wading	YC night-heron	Nesting								7 Pairs				
309	Shorebird	American oystercatcher	Nesting								12 Pairs				
	Shorebird	American oystercatcher	Roosting								3 Pairs				
323	Shorebird	American oystercatcher	Nesting								7 Pairs				
328	Shorebird	American oystercatcher	Nesting								13 Pairs				
	Shorebird	American oystercatcher	Roosting								2 Pairs				
349	Gull/Tern	G. black-backed gull	Nesting								16 Pairs				
353	Gull/Tern	G. black-backed gull	Nesting								5 Pairs				
	Gull/Tern	Herring gull	Nesting								12 Pairs				
933	Waterfowl	Canvasback	Wintering								100S				
	Waterfowl	Redhead	Wintering								1,000S				
	Waterfowl	Ruddy duck	Wintering								100S				
	Waterfowl	Scaup	Wintering								100S				
1160	Diving	Common loon	Wintering								Present				
	Diving	Grebes	Wintering								10S				
	Pelagic	Northern gannet	Wintering								Present				
	Watrfowl	Scoter	Wintering								1,000S				

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

BIRDS

Subelement	Species	Mapping Qualifier	MD\VA						Monthly Presence												
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
Diving	Common loon	Wintering			100S													-	-	-	-
	Common loon	Wintering			100S													-	-	-	-
D. crested cormorant	Migration				1,000S													Mar-May	Aug-Nov	-	-
D. crested cormorant	Wintering				1,000S													Mar-May	Aug-Nov	-	-
Grebes	Wintering				100S													-	-	-	-
Grebes	Wintering				100S													-	-	-	-
Red-throated loon	Wintering				100S													-	-	-	-
Northern gannet	Wintering				100S													-	-	-	-
American black duck	Wintering				100S													-	-	-	-
American wigeon	Wintering				1,000S													-	-	-	-
Blue-winged teal	Migration				100S													Feb-Apr	Aug-Oct	-	-
Brant	Wintering				1,000S													-	-	-	-
Bufflehead	Wintering				1,000S													-	-	-	-
Canada goose	Wintering				1,000S													-	-	-	-
Canvasback	Wintering				1,000S													-	-	-	-
Gadwall	Wintering				1,000S													-	-	-	-
Green-winged teal	Wintering				1,000S													-	-	-	-
Long-tailed duck	Wintering				100S													-	-	-	-
Mallard	Wintering				100S													-	-	-	-
Mergansers	Wintering				100S													-	-	-	-
Northern pintail	Wintering				100S													-	-	-	-
Redhead	Wintering				10,000S													-	-	-	-
Ring-necked duck	Wintering				Present													-	-	-	-
Ruddy duck	Wintering				10,000S													-	-	-	-
Scaup	Wintering				1,000S													-	-	-	-
Scoters	Wintering				10,000S													-	-	-	-
Tundra swan	Wintering				1,000S													-	-	-	-

FISH

Subelement	Species	Mapping Qualifier	MD\VA						Monthly Presence												
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles
Diadromous	Alewife	Migration			-												-	-	-	-	Aug-Nov
	American shad	Migration			-												-	-	-	-	Feb-Jun
Atlantic sturgeon	General Distribution	-IE	IE		-												-	-	-	-	Jan-Dec
Blueback herring	Migration				-												-	-	-	-	Mar-Jun
Hickory shad	General Distribution	E/E	E		-												-	-	-	-	Feb-May
Shortnose sturgeon	Migration				-												-	-	-	-	Jan-Dec
Striped bass																	-	-	-	-	Feb-Jun

REPTILES & AMPHIBIANS

		Monthly Presence																						
Subelement	Species	Mapping Qualifier		S	F	Concentration		J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles	Adults
Turtle	K. ridley sea turtle	General Distribution		E/E	E	Uncommon		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Dec	-
	K. ridley sea turtle	General Distribution		E/E	E	Common		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct	-
	Loggerhead sea turtle	General Distribution		T/T	T	Common		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Dec	Nov-Dec
	Loggerhead sea turtle	General Distribution		T/T	T	Abundant		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct	Apr-Oct

INVERTEBRATES

		Monthly Presence																						
Subelement	Species	Mapping Qualifier		S	F	Concentration		J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
Crab	Blue crab	Concentration Area		Highly Abundant-Summ				-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Oct	Jun-Oct

MARINE MAMMALS

		Monthly Presence																						
Subelement	Species	Mapping Qualifier		S	F	Concentration		J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping	Molt	
Manatee	West Indian manatee	General Distribution		-E	E	Rare		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Whale	Humpback whale	General Distribution		E/E	E	Very Rare		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)

BIRDS

		Monthly Presence																						
Subelement	Species	Mapping Qualifier		S	F	Concentration		J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt	
Diving	Brown pelican			Blue	Blue	Blue	Blue	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	Brown pelican			Blue	Blue	Blue	Blue	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	D. crested cormorant			Blue	Blue	Blue	Blue	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Gull/Tern	Gulls			Blue	Blue	Blue	Blue	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	Terns			Blue	Blue	Blue	Blue	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Pelagic	Storm-petrels			Blue	Blue	Blue	Blue	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

FISH

		Monthly Presence																						
Subelement	Species	Mapping Qualifier		S	F	Concentration		J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
Diadromous	American eel			Blue	Blue	Blue	Blue	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-May	
	Gizzard shad			Blue	Blue	Blue	Blue	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	
	Striped bass			Blue	Blue	Blue	Blue	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	
Estuarine Nursery	Atlantic croaker			Blue	Blue	Blue	Blue	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov	
	Atlantic menhaden			Blue	Blue	Blue	Blue	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	
	Black drum			Blue	Blue	Blue	Blue	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec	
	Black sea bass			Blue	Blue	Blue	Blue	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov	
	Bluefish			Blue	Blue	Blue	Blue	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct	
	Kingfishes			Blue	Blue	Blue	Blue	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov	
	Northern puffer			Blue	Blue	Blue	Blue	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Oct	
	Pigfish			Blue	Blue	Blue	Blue	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov	
	Red drum			Blue	Blue	Blue	Blue	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov	

FISH (continued)

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Sheepshead	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Spot	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Weakfish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Winter flounder	-	-	-	-	-	-	-	-	-	-	-	-	Dec-Mar
 Estuarine Resident	Pipefish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
 Fish	Silversides	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
 Marine Benthic	Anchovies	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Striped burrfish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Atlantic spadefish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Clearnose skate	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Red hake	-	-	-	-	-	-	-	-	-	-	-	-	Sep-May*
	Silver hake	-	-	-	-	-	-	-	-	-	-	-	-	Oct-May
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
 Marine Pelagic	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Nov
	Cobia	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Lined seahorse	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Spanish mackerel	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		-	-	-	-	-	-	-	-	-	-	-	-	May-Sep

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Hard clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Stout razor clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
 Cephalopod	Squid	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
 Crab	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Horseshoe crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
 Shrimp	Brown shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Mantis shrimp	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	White shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct

MARINE MAMMALS

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
 Dolphin	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-
	Harbor porpoise	-	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES

ESI POLYGON HABITAT TYPES		Habitat Classification	Area (Acres)	Area (Sq. Miles)
ESI Rank				
10A		Salt and Brackish Water Marshes	622.42	0.97
10B		Freshwater Marshes	15.57	0.02
10D		Scrub and Shrub Wetlands	2.28	0.00
9A		Sheltered Tidal Flats	2.79	0.00
7		Exposed Tidal Flats	70.14	0.11

ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
ESI Rank					
10A		Salt and Brackish Water Marshes	42,917.97	26.67	48%
10D		Scrub and Shrub Wetlands	5,689.07	3.54	6%
9A		Sheltered Tidal Flats	1,324.28	0.82	1%
9B		Vegetated Low Banks	39.31	0.02	< 1%
8B		Sheltered, Solid Man-Made Structures	2,262.92	1.41	3%
8C		Sheltered Riprap	70.90	0.04	< 1%
7		Exposed Tidal Flats	21,727.24	13.50	24%
6B		Riprap	2,725.85	1.69	3%
3A		Fine to Medium Grained Sand Beaches	12,525.75	7.78	14%
1B		Exposed, Solid Man-Made Structures	278.93	0.17	< 1%
Total ESI Shoreline:		89,562.23	Total ESI Shoreline:	55.65	
Total Shoreline:		51,323.37	Total Shoreline:	31.89	

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%.

All underlying GIS data can be obtained from response.restoration.noaa.gov



Map 51


76° 18'

76° 16'

76° 14'

76° 12'

76° 10'



Map 59 Central Bay



SEE BACK OF MAP
for details about mapped resources and
other resources that occur in mapped area.
Data Published: September 2016

0 Not for Navigation 2 Miles
0 2 Kilometers

1:100,000



Map 59
Central Bay

HUMAN USE RESOURCES

DISPLAYED ON MAP (POINTS)

Map ID	Type	Name	Contact	Phone
1	AIRPORT	FEDERHART-OPHELIA LAND'S END		
2	AIRPORT	REEDVILLE		
3	AIRPORT	REMO PRIVATE		
4	AIRPORT	OYSTER NON-COMM ALLEN RF#98-2025	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
5	AQUACULTURE	OYSTER NON-COMM BRANCH, JAMES RF#98-1619	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
6	AQUACULTURE	OYSTER NON-COMM HILLEN RF#99-2049	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
7	AQUACULTURE	OYSTER NON-COMM KOEHLER, BILL RF#98-1481	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
8	AQUACULTURE	OYSTER NON-COMM LAND RF#99-1985	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
9	AQUACULTURE	OYSTER NON-COMM LAUBER, RAY RF#98-1764	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
10	AQUACULTURE	OYSTER NON-COMM LIVEZEY, JANET RF#2001-1083	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
11	AQUACULTURE	OYSTER NON-COMM OVERCASH, K.E. RF#98-0759	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
12	AQUACULTURE	OYSTER NON-COMM PAYNE, JAMES RF#2000-0721	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
13	AQUACULTURE	OYSTER NON-COMM PERRY, RON RF#00-1826	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
14	AQUACULTURE	OYSTER NON-COMM PISCIOTTA RF#00-0637	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
15	AQUACULTURE	OYSTER NON-COMM PULLY RF#00-0185	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
16	AQUACULTURE	OYSTER NON-COMM SLIEKO, MARK RF#98-1696	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
17	AQUACULTURE	OYSTER NON-COMM WILLIAMS RF#99-0345	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
18	AQUACULTURE	OYSTER NON-COMM WISE RF#98-1772	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
19	AQUACULTURE	OYSTER NON-COMM WISE, LLOYD, E. RF#98-0778	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
20	AQUACULTURE	NORTHERN NECK SHELL	LENE PENNINGTON, VDGIF	804-367-0058
21	ARTIFICIAL REEF	REEDVILLE FERRY	GREG PINIAK, PROJECT MANAGER	301-713-3028
22	BOAT RAMP	MWS - SITE CBIB	MARYLAND DEPARTMENT OF THE ENVIRONMENT	410-537-3608
23	FERRY	WQ - SHELLFISH HARVEST WATERS, POTOMAC RIVER	CHESAPEAKE BAY PROGRAM	800-968-7229
24	REPEATED MEASUREMENT SITE	WQ - STATION CB5.3	CHESAPEAKE BAY PROGRAM	800-968-7229
25	REPEATED MEASUREMENT SITE	WQ - STATION CB5.4	CHESAPEAKE BAY PROGRAM	800-968-7229
26	REPEATED MEASUREMENT SITE	WQ - STATION CB5.4W	VA DEPT OF CONSERVATION AND RECREATION	
27	REPEATED MEASUREMENT SITE			
28	REPEATED MEASUREMENT SITE			

ALSO PRESENT IN MAPPED AREA (POLYGONS)

Type	Name	Contact	Phone
STATE PROTECTED AREA	BLUE CRAB SANCTUARY	VIRGINIA MARINE RESOURCES COMMISSION	757-247-2200
STATE PROTECTED AREA	DAMERON MARSH	VA DEPT OF CONSERVATION AND RECREATION	

JURISDICTIONS

COUNTY: ACCOMACK CO., NORTHUMBERLAND CO., SOMERSET CO., ST. MARY'S CO.
COAST GUARD: DISTRICT 5, SECTOR BALTIMORE, SECTOR HAMPTON ROADS
USACE: NORTH ATLANTIC DIVISION, BALTIMORE DISTRICT, NORFOLK DISTRICT

SHORELINE RESOURCES

FEMA: REGION I
EPA: REGION 3

ESI POLYGON HABITAT TYPES

ESI Rank	Habitat Classification	Area (Acres)	Area (Sq. Miles)
10A	Salt and Brackish Water Marshes	418.11	0.65
10B	Freshwater Marshes	37.73	0.06
10C	Swamps	897.70	1.40
10D	Scrub and Shrub Wetlands	149.31	0.23
7	Exposed Tidal Flats	4.00	0.01

ESI SHORELINE HABITAT TYPES

ESI Rank	Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
10A	Salt and Brackish Water Marshes	72,311.20	44.93	16%
10C	Swamps	2,702.57	1.68	1%
10D	Scrub and Shrub Wetlands	2,229.06	1.39	< 1%
9B	Vegetated Low Banks	187,930.63	116.77	41%
8B	Sheltered, Solid Man-Made Structures	57,196.55	35.54	13%
8C	Sheltered Riprap	33,363.88	20.73	7%
7	Exposed Tidal Flats	36,001.77	22.37	8%
6B	Riprap	21,427.90	13.31	5%
4	Coarse Grained Sand Beaches	33,719.69	20.95	7%
3B	Scars and Steep Slopes (Sand)	1,860.94	1.16	< 1%
1B	Exposed, Solid Man-Made Structures	5,759.58	3.58	1%

Total ESI Shoreline: 454,503.76
Total Shoreline: 371,520.66

Total ESI Shoreline: 282.42
Total Shoreline: 230.85

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%.

All underlying GIS data can be obtained from response.restoration.noaa.gov




Map 51C

76° 18' W

76° 16' W

76° 14' W



Map 59A Central Bay



SEE BACK OF MAP
for details about mapped species and
other species that occur in mapped area.
Data Published: September 2016



Map 59A Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BIRDS

Map ID	Subelement	Species	Monthly Presence											
			J	F	M	A	M	J	J	A	S	O	N	D
			Mapping Qualifier	S	F	Concentration	90 Nests							
197	Wading	Great blue heron	Nesting											
232	Watertowl	Canada goose	Nesting											
	Watertowl	Mallard	Nesting											
669	Watertowl	American black duck	Wintering											
	Watertowl	American coot	Wintering											
	Watertowl	American wigeon	Wintering											
	Watertowl	Bufflehead	Wintering											
	Watertowl	Canada goose	Wintering											
	Watertowl	Canvasback	Wintering											
	Watertowl	Common goldeneye	Wintering											
	Watertowl	Gadwall	Wintering											
	Watertowl	Mallard	Wintering											
	Watertowl	Mergansers	Wintering											
	Watertowl	Ring-necked duck	Wintering											
	Watertowl	Ruddy duck	Wintering											
	Watertowl	Scaup	Wintering											
	Watertowl	Snow goose	Wintering											
	Watertowl	Tundra swan	Wintering											
707	Raptor	Bald eagle	Roosting											
798	Raptor	Bald eagle	Nesting											
809	Watertowl	American black duck	Wintering											
	Watertowl	Bufflehead	Wintering											
	Watertowl	Canada goose	Wintering											
	Watertowl	Mallard	Wintering											
906	Watertowl	American black duck	Wintering											
	Watertowl	Bufflehead	Wintering											
	Watertowl	Canada goose	Wintering											
	Watertowl	Gadwall	Wintering											
	Watertowl	Mallard	Wintering											
	Watertowl	Mergansers	Wintering											
	Watertowl	Ruddy duck	Wintering											
1088	Diving	D. crested cormorant	Migration											
	Diving	Grebes	Wintering											
1124	Diving	Grebes	Wintering											
	Diving	Red-throated loon	Wintering											
	Watertowl	Bufflehead	Wintering											
	Watertowl	Canada goose	Wintering											

BIRDS (continued)

<i>Map ID</i>	<i>Subelement</i>	<i>Species</i>	<i>Mapping Qualifier</i>	<i>MD/VA</i>	<i>S</i>	<i>F</i>	<i>Concentration</i>	<i>J F M A M J J A S O N D</i>	<i>Nest</i>	<i>Mig.(S)</i>	<i>Mig.(F)</i>	<i>Molt</i>
Waterfowl	Long-tailed duck	Wintering			10S			-	-	-	-	-
Waterfowl	Mergansers	Wintering			100S			-	-	-	-	-
Waterfowl	Ruddy duck	Wintering			100S			-	-	-	-	-

FISH

<i>Map ID</i>	<i>Subelement</i>	<i>Species</i>	<i>Mapping Qualifier</i>	<i>MD/VA</i>	<i>S</i>	<i>F</i>	<i>Concentration</i>	<i>J F M A M J J A S O N D</i>	<i>Spawn</i>	<i>Eggs</i>	<i>Larvae</i>	<i>Juveniles</i>	<i>Adults</i>
1623	Diadromous	Alewife	Migration		-			-	-	-	-	Aug-Nov	Feb-Apr
	Diadromous	American shad	Migration		-			-	-	-	-	Aug-Nov	Feb-Jun
	Diadromous	Blueback herring	Migration		-			-	-	-	-	Aug-Nov	Mar-Jun
	Diadromous	Hickory shad	Migration		-			-	-	-	-	Jun-Sep	Feb-May
	Diadromous	Striped bass	Migration		-			-	-	-	-	-	Feb-Jun

<i>Map ID</i>	<i>Subelement</i>	<i>Species</i>	<i>Mapping Qualifier</i>	<i>MD/VA</i>	<i>S</i>	<i>F</i>	<i>Concentration</i>	<i>J F M A M J J A S O N D</i>	<i>Spawn</i>	<i>Eggs</i>	<i>Larvae</i>	<i>Juveniles</i>	<i>Adults</i>
1833	Insect	NE beach tiger beetle	Vulnerable Occurrence		E/T	T	-	-	-	-	-	-	Jun-Aug

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)													
---	--	--	--	--	--	--	--	--	--	--	--	--	--

BIRDS

<i>Subelement</i>	<i>Species</i>	<i>Mapping Qualifier</i>	<i>MD/VA</i>	<i>S</i>	<i>F</i>	<i>Concentration</i>	<i>J F M A M J J A S O N D</i>	<i>Nest</i>	<i>Mig.(S)</i>	<i>Mig.(F)</i>	<i>Molt</i>
Diving	Common loon	Wintering		Present			-	-	-	-	-
	Common loon	Wintering		100S			-	-	-	-	-
	D. crested cormorant	Wintering		1,000S			-	-	-	-	-
	Grebens	Wintering		100S			-	-	-	-	-
	Red-throated loon	Wintering		Present			-	-	-	-	-
	Red-throated loon	Wintering		100S			-	-	-	-	-
Pelagic	Northern gannet	Wintering		100S			-	-	-	-	-
	Northern gannet	Wintering		Present			-	-	-	-	-
	Long-tailed duck	Wintering		100S			-	-	-	-	-
Waterfowl	Scoters	Wintering		10,000S			-	-	-	-	-
	Scoters	Wintering		10S			-	-	-	-	-
	Tundra swan	Wintering					-	-	-	-	-

FISH

Subelement	Species	MD\VA												Monthly Presence													
		S	F	Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults			
Diadromous	Alewife	Migration	-	Moderate Run			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Mar-May		
	Alewife	Migration	-	Strong Run			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov	Mar-May	
	American shad	Migration	-				-																			Aug-Nov	Mar-Jun
	American shad	Migration	-				-																			Aug-Nov	Mar-Jun
	Atlantic sturgeon	General Distribution	-/E	E	Moderate Run			-																		Jan-Dec	
	Blueback herring	Migration	-				-																			Feb-Jun	
	Blueback herring	Migration	-				-																			Feb-Jun	
	Hickory shad	Migration	-				-																			Feb-Jun	
	Hickory shad	Migration	-				-																			Feb-Jun	
	Shortnose sturgeon	General Distribution	E/E	E	Moderate Run			-																		Jan-Dec	
	Shortnose sturgeon	Migration	E/E	E	-			-																		Mar-Jun	
	Striped bass	Migration	-				-																			Mar-Jun	
	Striped bass	Nursery Area	-				-																			-	

REPTILES & AMPHIBIANS



Subelement	Species	MD\VA												Monthly Presence											
		S	F	Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles	Adults	
Turtle	K. ridley sea turtle	General Distribution	E/E	E	Uncommon			-												-					Nov-Dec
	K. ridley sea turtle	General Distribution	E/E	E	Common			-												-					Apr-Oct
	Loggerhead sea turtle	General Distribution	T/T	T	Common			-												-					Nov-Dec
	Loggerhead sea turtle	General Distribution	T/T	T	Abundant			-												-					Apr-Oct

INVERTEBRATES

Subelement	Species	MD\VA												Monthly Presence											
		S	F	Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults	
Crab	Blue crab	Concentration Area	Highly Abundant-Summ												-	-	-	-	-	-	-	-	-	-	Jun-Oct

MARINE MAMMALS

Subelement	Species	MD\VA												Monthly Presence											
		S	F	Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping	Molt		
Manatee	West Indian manatee	General Distribution	-/E	E	Rare			-												-	-	-	-	-	Aug-Nov
Whale	Humpback whale	General Distribution	E/E	E	Very Rare			-												-	-	-	-	-	Mar-May

BIRDS**FISH**

FISH (continued)

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Red hake	-	-	-	-	-	-	-	-	-	-	-	-	Sep-May*
	Silver hake	-	-	-	-	-	-	-	-	-	-	-	-	Oct-May
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Marine Pelagic	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Sep
	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Cobia	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Lined seahorse	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Spanish mackerel	-	-	-	-	-	-	-	-	-	-	-	-	May-Sep

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Stout razor clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Cephalopod	Squid	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
Crab	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Horseshoe crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Brown shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Mantis shrimp	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	White shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct

MARINE MAMMALS

Subelement	Species	Monthly Presence												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-
	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-
	Harbor porpoise	-	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES


ESI POLYGON HABITAT TYPES		Habitat Classification	Area (Acres)	Area (Sq. Miles)
ESI Rank				
10A		Salt and Brackish Water Marshes	139.20	0.22
10B		Freshwater Marshes	34.20	0.05
10C		Swamps	498.37	0.78
10D		Scrub and Shrub Wetlands	64.94	0.10
7		Exposed Tidal Flats	4.00	0.01

ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
ESI Rank					
10A		Salt and Brackish Water Marshes	29,322.94	18.22	14%
10C		Swamps	805.44	0.50	< 1%
10D		Scrub and Shrub Wetlands	816.16	0.51	< 1%
9B		Vegetated Low Banks	93,073.90	57.83	45%
8B		Sheltered, Solid Man-Made Structures	28,694.36	17.83	14%
8C		Sheltered Riprap	14,638.32	9.10	7%
7		Exposed Tidal Flats	13,646.84	8.48	7%
6B		Riprap	5,973.84	3.71	3%
4		Coarse Grained Sand Beaches	17,178.47	10.67	8%
3B		Scars and Steep Slopes (Sand)	511.60	0.32	< 1%
1B		Exposed, Solid Man-Made Structures	2,709.57	1.68	1%
Total ESI Shoreline:		207,371.44	Total ESI Shoreline:	128.85	
Total Shoreline:		173,950.97	Total Shoreline:	108.09	

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 59B
Central Bay



SEE BACK OF MAP
for details about mapped species and
other species that occur in mapped area.
Data Published: September 2016

Map 59B Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	Mapping Qualifier			S	F	Concentration	Monthly Presence										
		High Ecological Value	High Ecological Value	High Ecological Value				J	F	M	A	M	J	J	A	S	O	N
SAV	Submerged aquatic veg							-	-	-	-	-	-	-	-	-	-	-
SAV	Submerged aquatic veg							-	-	-	-	-	-	-	-	-	-	-
SAV	Submerged aquatic veg							-	-	-	-	-	-	-	-	-	-	-

BIRDS

Map ID	Subelement	Species	Mapping Qualifier			S	F	Concentration	Monthly Presence										
			J	F	M				J	F	M	A	M	J	J	A	S	O	N
38	Diving	Grebes	Wintering						-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Bufflehead	Wintering						-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Canada goose	Wintering						-	-	-	-	-	-	-	-	-	-	-
232	Waterfowl	Scoters	Wintering						-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Canada goose	Nesting						-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Mallard	Nesting						-	-	-	-	-	-	-	-	-	-	-
757	Waterfowl	American black duck	Wintering						-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	American wigeon	Wintering						-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Canvasback	Wintering						-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Gadwall	Wintering						-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Mallard	Wintering						-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Redhead	Wintering						-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Scaup	Wintering						-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Tundra swan	Wintering						-	-	-	-	-	-	-	-	-	-	-
798	Raptor	Bald eagle	Nesting						-	-	-	-	-	-	-	-	-	-	-
875	Waterfowl	American black duck	Wintering						-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Blue-winged teal	Migration						-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Canvasback	Wintering						-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Common goldeneye	Wintering						-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Green-winged teal	Wintering						-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Mallard	Wintering						-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Redhead	Wintering						-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Ring-necked duck	Wintering						-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Ruddy duck	Wintering						-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Tundra swan	Wintering						-	-	-	-	-	-	-	-	-	-	-
1189	Diving	D. crested cormorant	Wintering						-	-	-	-	-	-	-	-	-	-	-
	Diving	Grebes	Wintering						-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Scoters	Wintering						-	-	-	-	-	-	-	-	-	-	-

INVERTEBRATES

Map ID	Subelement	Species	Mapping Qualifier			MD/VA			Monthly Presence													
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
1833	Insect	NE beach tiger beetle	Vulnerable Occurrence	-	-	E/T	T	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

WIDESpread IN MAPPED AREA (> 10 SQUARE KILOMETERS)

BIRDS

Subelement	Species	Mapping Qualifier	MD/VA			Monthly Presence																
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt	
Diving	Common loon	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	Common loon	Wintering	-	-	Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	D. crested cormorant	Wintering	-	-	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	Grebes	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	Red-throated loon	Wintering	-	-	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	Red-throated loon	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Pelagic	Northern gannet	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	Waterfowl	American black duck	Wintering	-	Bufflehead	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		Canada goose	Wintering	-	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		Long-tailed duck	Wintering	-	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		Long-tailed duck	Wintering	-	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		Mallard	Wintering	-	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		Mallard	Wintering	-	Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		Mergansers	Wintering	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		Ruddy duck	Wintering	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		Scoters	Wintering	-	10,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		Tundra swan	Wintering	-	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Subelement	Species	Mapping Qualifier	MD/VA			Monthly Presence															
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles
Diadromous	Alewife	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	American shad	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Atlantic sturgeon	General Distribution	-/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Blueback herring	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Hickory shad	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Shortnose sturgeon	General Distribution	E/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Striped bass	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

REPTILES & AMPHIBIANS

Subelement	Species	Mapping Qualifier	MD/VA			Monthly Presence															
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles
Turtle	K. ridley sea turtle	General Distribution	E/E	E	Uncommon	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	K. ridley sea turtle	General Distribution	E/E	E	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Loggerhead sea turtle	General Distribution	T/T	T	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Loggerhead sea turtle	General Distribution	T/T	T	Abundant	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

INVERTEBRATES

Subelement	Species	Mapping Qualifier	Monthly Presence																		
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawning	Eggs	Larvae	Juveniles
Crab	Blue crab	Highly Abundant-Summer	-/E	-/E	Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Oct

MARINE MAMMALS

Subelement	Species	Mapping Qualifier	Monthly Presence																	
			J	F	M	A	M	J	J	A	S	O	N	D	Spawning					
Manatee	West Indian manatee	General Distribution	-/E	-/E	Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Whale	Humpback whale	General Distribution	E/E	E/E	Very Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)

BIRDS

Subelement	Species	Monthly Presence																		
		J	F	M	A	M	J	J	A	S	O	N	D	Spawning						
Diving	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Gull/Tern	Gulls	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Terns	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Pelagic	Storm-petrels	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Wood duck	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Watrfowl		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Subelement	Species	Monthly Presence																		
		J	F	M	A	M	J	J	A	S	O	N	D	Spawning						
Diadromous	Alewife	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	American eel	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Blueback herring	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Gizzard shad	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Estuarine Nursery	Striped bass	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Atlantic menhaden	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Black drum	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Black sea bass	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Bluefish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Kingfishes	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pigfish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Red drum	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Sheepshead	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Spot	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Weakfish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH (continued)

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Winter flounder	-	-	-	-	-	-	-	-	-	-	-	-	Dec-Mar
Estuarine Resident	Pipefish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Silversides	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
Fish	Anchoovies	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Striped burrfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
Marine Benthic	Atlantic spadefish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Red hake	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Silver hake	-	-	-	-	-	-	-	-	-	-	-	-	Sep-May*
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Oct-May
Marine Pelagic	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Cobia	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Lined seahorse	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Spanish mackerel	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		-	-	-	-	-	-	-	-	-	-	-	-	May-Sep

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Bivalve	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Stout razor clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Cephalopod	Squid	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Horseshoe crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Brown shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Mantis shrimp	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	White shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct

MARINE MAMMALS

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Dolphin	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-
	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-
	Harbor porpoise	-	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES


ESI POLYGON HABITAT TYPES		Habitat Classification	Area (Acres)	Area (Sq. Miles)
ESI Rank	ESI/Rank			
	10A	Salt and Brackish Water Marshes	278.91	0.44
	10B	Freshwater Marshes	3.54	0.01
	10C	Swamps	399.33	0.62
	10D	Scrub and Shrub Wetlands	84.37	0.13

ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
ESI Rank	ESI/Rank				
	10A	Salt and Brackish Water Marshes	42,988.25	26.71	17%
	10C	Swamps	1,897.13	1.18	1%
	10D	Scrub and Shrub Wetlands	1,412.90	0.88	1%
	9B	Vegetated Low Banks	94,856.73	58.94	38%
	8B	Sheltered, Solid Man-Made Structures	28,502.19	17.71	12%
	8C	Sheltered Riprap	18,725.56	11.64	8%
	7	Exposed Tidal Flats	22,354.93	13.89	9%
	6B	Riprap	15,454.06	9.60	6%
	4	Coarse Grained Sand Beaches	16,541.22	10.28	7%
	3B	Scars and Steep Slopes (Sand)	1,349.34	0.84	1%
	1B	Exposed, Solid Man-Made Structures	3,050.01	1.90	1%
		Total ESI Shoreline:	247,132.32	Total ESI Shoreline: 153.56	
		Total Shoreline:	197,569.69	Total Shoreline: 122.76	

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 60
Central Bay



SEE BACK OF MAP
for details about mapped resources and
other resources that occur in mapped area.
Data Published: September 2016

Map 60
Central Bay

HUMAN USE RESOURCES

DISPLAYED ON MAP (POINTS)

Map ID	Type	Name	Contact	Phone
1	AIRPORT	JETT AIRPARK		
2	AIRPORT	SERENITY FARM		
3	AQUACULTURE	OYSTER COMMERCIAL MIX, DAVID A. RF#02-0026	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
4	AQUACULTURE	OYSTER NON-COMM CALLEDEN, ROBERT, E. RF#99-1182	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2234
5	AQUACULTURE	OYSTER NON-COMM FOLTS, JAMES RF#98-2026	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2234
6	AQUACULTURE	OYSTER NON-COMM GROMELSKI, STAN RF#99-1248	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
7	AQUACULTURE	OYSTER NON-COMM KAUDERS, ANDREW E. RF#98-1722	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
8	AQUACULTURE	OYSTER NON-COMM LOCKHART, KENT RF#00-0385	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
9	AQUACULTURE	OYSTER NON-COMM MCKINLEY, ROGER RF#01-0357	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
10	AQUACULTURE	OYSTER NON-COMM MIZUNO RF#00-1814	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
11	AQUACULTURE	OYSTER NON-COMM SMITH, HERBERT RF#99-1001	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
12	AQUACULTURE	OYSTER NON-COMM STEG RF#00-2118	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
13	AQUACULTURE	OYSTER NON-COMM TEXTER RF#99-0830	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
14	AQUACULTURE	OYSTER NON-COMM WINKEL, RAYMOND N. RF#99-1177	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2234
15	BOAT RAMP	COOPERS	LENE PENNINGTON, VDGIF	804-367-0058

ALSO PRESENT IN MAPPED AREA (POLYGONS)

Type	Name	Contact	Phone
STATE PROTECTED AREA	BUSH MILL STREAM	VA DEPT OF CONSERVATION AND RECREATION	
STATE PROTECTED AREA	CHILTON WOODS	VA DEPT OF FORESTRY	
STATE PROTECTED AREA	HICKORY HOLLOW	NORTHERN NECK AUDUBON SOCIETY	

JURISDICTIONS

COUNTY: LANCASTER CO., NORTHUMBERLAND CO., RICHMOND CO., ST. MARY'S CO.
 COAST GUARD: DISTRICT 5, SECTOR BALTIMORE, SECTOR HAMPTON ROADS
 USACE: NORTH ATLANTIC DIVISION, BALTIMORE DISTRICT, NORFOLK DISTRICT

SHORELINE RESOURCES

FEMA:
 EPA:
 REGION I
 REGION 3

SHORELINE HABITAT TYPES

<i>ESI Rank</i>	<i>Habitat Classification</i>	<i>Area (Acres)</i>	<i>Area (Sq. Miles)</i>
10A	Salt and Brackish Water Marshes	369.42	0.58
10B	Freshwater Marshes	170.52	0.27
10C	Swamps	3,337.00	5.21
10D	Scrub and Shrub Wetlands	480.37	0.75
7	Exposed Tidal Flats	13.66	0.02

ESI SHORELINE HABITAT TYPES


<i>ESI Rank</i>	<i>Shoreline Habitat Classification</i>	<i>Length (Meters)</i>	<i>Length (Miles)</i>	<i>% of ESI Shoreline</i>
10A	Salt and Brackish Water Marshes	61,992.42	38.52	20%
10B	Freshwater Marshes	12,310.97	7.65	4%
10C	Swamps	1,834.25	1.14	1%
10D	Scrub and Shrub Wetlands	2,166.45	1.35	1%
9B	Vegetated Low Banks	161,918.60	100.61	53%
8B	Sheltered, Solid Man-Made Structures	24,614.37	15.29	8%
8C	Sheltered Riprap	8,105.59	5.04	3%
7	Exposed Tidal Flats	11,184.50	6.95	4%
6B	Riprap	7,288.87	4.53	2%
4	Coarse Grained Sand Beaches	9,238.71	5.74	3%
3B	Scars and Steep Slopes (Sand)	1,330.38	0.83	< 1%
1B	Exposed, Solid Man-Made Structures	4,435.19	2.76	1%

Total ESI Shoreline: 306,420.31 Total ESI Shoreline: 190.40
 Total Shoreline: 280,727.37 Total Shoreline: 174.44

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%.

All underlying GIS data can be obtained from response.restoration.noaa.gov





SEE BACK OF MAP
for details about mapped species and
other species that occur in mapped area.
Data Published: September 2016

0 Not for Navigation 1 Miles
0 1 Kilometers 1:50,000



Map 60A Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BIRDS

Map ID	Subelement	Species	Mapping Qualifier	S	F	Concentration	Monthly Presence														
							J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)
166	Wading	Great blue heron	Nesting			11 Nests	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
198	Wading	Great blue heron	Nesting			46 Nests	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
232	Waterfowl	Canada goose	Nesting			Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Mallard	Nesting			Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
669	Waterfowl	American black duck	Wintering			100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	American coot	Wintering			10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	American wigeon	Wintering			Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Bufflehead	Wintering			1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Canada goose	Wintering			10,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Canvasback	Wintering			11,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Common goldeneye	Wintering			10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Gadwall	Wintering			10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Mallard	Wintering			100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Mergansers	Wintering			100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Ring-necked duck	Wintering			100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Ruddy duck	Wintering			1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Scaup	Wintering			1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Snow goose	Wintering			Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Tundra swan	Wintering			100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
707	Raptor	Bald eagle	Roosting			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
798	Raptor	Bald eagle	Nesting			1 Nest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
808	Waterfowl	American black duck	Wintering			10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Bufflehead	Wintering			10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Canada goose	Wintering			100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Mallard	Wintering			Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Tundra swan	Wintering			10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
905	Waterfowl	American black duck	Wintering			Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Bufflehead	Wintering			100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Canada goose	Wintering			1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Gadwall	Wintering			10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Mallard	Wintering			10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Mergansers	Wintering			10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Ruddy duck	Wintering			100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Tundra swan	Wintering			10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1161	Diving	Common loon	Wintering			Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Diving	Grebes	Wintering			100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Diving	Red-throated loon	Wintering			Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

BIRDS (continued)

Map ID	Subelement	Species	MD/VA			Monthly Presence															
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
Pelagic		Northern gannet			Present												-	-	-	-	-
Watertowl		Long-tailed duck			Wintering												-	-	-	-	-
Watertowl		Scoters			Wintering												-	-	-	-	-

FISH

Map ID	Subelement	Species	MD/VA			Monthly Presence															
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles
1620	Diadromous	Alewife	Migration														-	-	-	Aug-Nov	Feb-Apr
		Blueback herring	Migration														-	-	-	Aug-Nov	Mar-Jun
1625	Diadromous	Alewife	Migration														-	-	-	Aug-Nov	Feb-Apr
		Blueback herring	Migration														-	-	-	Aug-Nov	Mar-Jun
		Striped bass	Migration														-	-	-	Feb-Jun	Feb-Jun
1643	Diadromous	Alewife	Migration														-	-	-	Aug-Nov	Mar-May
		American shad	Migration														-	-	-	Aug-Nov	Mar-Jun
		Blueback herring	Migration														-	-	-	Aug-Nov	Feb-Jun
		Hickory shad	Migration														-	-	-	Jun-Oct	Feb-Jun
		Shortnose sturgeon	Migration														-	-	-	Mar-Jun	Mar-Jun
		Striped bass	Nursery Area														-	-	-	Apr-Jun	Jan-Dec

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

FISH

Subelement	Species	MD/VA			Monthly Presence															
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles
Diadromous	Atlantic sturgeon	General Distribution	-E	E	-											-	-	-	-	Jan-Dec
	Shortnose sturgeon	General Distribution	E/E	E	-											-	-	-	-	Jan-Dec

REPTILES & AMPHIBIANS

Subelement	Species	MD/VA			Monthly Presence															
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles
Turtle	K. ridley sea turtle	General Distribution	E/E	E	Common											-	-	-	-	Apr-Oct
	K. ridley sea turtle	General Distribution	E/E	E	Uncommon											-	-	-	-	Nov-Dec
	Loggerhead sea turtle	General Distribution	T/T	T	Common											-	-	-	-	Nov-Dec
	Loggerhead sea turtle	General Distribution	T/T	T	Abundant											-	-	-	-	Apr-Oct

INVERTEBRATES

Subelement	Species	MD/VA			Monthly Presence															
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles
Crab	Blue crab	Concentration Area	Highly Abundant-Summ													-	-	-	-	Jun-Oct

MARINE MAMMALS

Subelement	Species	MD/VA			Monthly Presence															
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping	Molt
Manatee	West Indian manatee	General Distribution	-E	E	Rare											-	-	-	-	Jun-Oct

BIRDS

Sublement	Species	<i>Monthly Presence</i>												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
Diving	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-	-
Gull/Tern	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-	-
	Gulls	-	-	-	-	-	-	-	-	-	-	-	-	-
	Terns	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Sublement	Species	<i>Monthly Presence</i>												Adults			
		J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles
Diadromous	Alewife	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Apr	Feb-Apr	Feb-Apr	Feb-Apr
	American eel	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Blueback herring	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Gizzard shad	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Striped bass	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Estuarine Nursery	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Atlantic menhaden	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Black drum	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Black sea bass	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Bluefish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Kingfishes	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pigfish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Red drum	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Sheepshead	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Spot	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Weakfish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	White perch	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Winter flounder	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Estuarine Resident	Pipefish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Silversides	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Anchovies	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Striped burrfish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Atlantic spadefish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Cleawnose skate	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Red hake	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Silver hake	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Marine Pelagic	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH (continued)

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Sep
	Cobia	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Lined seahorse	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Spanish mackerel	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Squid	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Horseshoe crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Brown shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Mantis shrimp	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	White shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct

MARINE MAMMALS

Subelement	Species	Monthly Presence												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES

ESI POLYGON HABITAT TYPES		Habitat Classification		Area (Acres)	Area (Sq. Miles)
ESI Rank					
10A		Salt and Brackish Water Marshes		124.04	0.19
10B		Freshwater Marshes		17.71	0.03
10C		Swamps		482.30	0.75
10D		Scrub and Shrub Wetlands		89.83	0.14
7		Exposed Tidal Flats		13.66	0.02

ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification		Length (Meters)	Length (Miles)	% of ESI Shoreline
ESI Rank						
10A		Salt and Brackish Water Marshes		25,993.10	16.15	17%
10B		Freshwater Marshes		2,953.27	1.84	2%
10C		Swamps		144.85	0.09	< 1%
10D		Scrub and Shrub Wetlands		1,115.75	0.69	1%
9B		Vegetated Low Banks		86,819.81	53.95	56%
8B		Sheltered, Solid Man-Made Structures		10,148.76	6.31	7%
8C		Sheltered Riprap		2,458.19	1.53	2%
7		Exposed Tidal Flats		8,929.42	5.55	6%
6B		Riprap		5,266.07	3.27	3%
4		Coarse Grained Sand Beaches		6,890.27	4.28	4%
3B		Scars and Steep Slopes (Sand)		1,032.88	0.64	1%
1B		Exposed, Solid Man-Made Structures		2,372.07	1.47	2%
		Total ESI Shoreline:		154,124.45	Total ESI Shoreline:	95.77
		Total Shoreline:		135,389.38	Total Shoreline:	84.13

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov




Map 52C

76° 30' W

76° 28' W

Map 51D

76° 26' W



Map 60B
Central Bay

Map 60B Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BIRDS

Map ID	Subelement	Species	Mapping Qualifier	S	F	Concentration	Monthly Presence						
							J	F	M	A	M	J	J
232	Waterfowl	Canada goose	Nesting			Present	-	-	-	-	-	-	-
	Waterfowl	Mallard	Nesting			Present	-	-	-	-	-	-	-
669	Waterfowl	American black duck	Wintering			100S	-	-	-	-	-	-	-
	Waterfowl	American coot	Wintering			10S	-	-	-	-	-	-	-
	Waterfowl	American wigeon	Wintering			Present	-	-	-	-	-	-	-
	Waterfowl	Bufflehead	Wintering			1,000S	-	-	-	-	-	-	-
	Waterfowl	Canada goose	Wintering			10,000S	-	-	-	-	-	-	-
	Waterfowl	Canvasback	Wintering			1,000S	-	-	-	-	-	-	-
	Waterfowl	Common goldeneye	Wintering			10S	-	-	-	-	-	-	-
	Waterfowl	Gadwall	Wintering			10S	-	-	-	-	-	-	-
	Waterfowl	Mallard	Wintering			100S	-	-	-	-	-	-	-
	Waterfowl	Mergansers	Wintering			100S	-	-	-	-	-	-	-
	Waterfowl	Ring-necked duck	Wintering			100S	-	-	-	-	-	-	-
	Waterfowl	Ruddy duck	Wintering			1,000S	-	-	-	-	-	-	-
	Waterfowl	Scaup	Wintering			1,000S	-	-	-	-	-	-	-
	Waterfowl	Snow goose	Wintering			Present	-	-	-	-	-	-	-
	Waterfowl	Tundra swan	Wintering			100S	-	-	-	-	-	-	-
798	Raptor	Bald eagle	Nesting			1 Nest	-	-	-	-	-	-	-

FISH

Map ID	Subelement	Species	Mapping Qualifier	S	F	Concentration	Monthly Presence						
							J	F	M	A	M	J	J
1561	Diadromous	Alewife	Migration			-	-	-	-	-	-	-	-
	Diadromous	Atlantic sturgeon	General Distribution	-E	E	-	-	-	-	-	-	-	-
	Diadromous	Blueback herring	Migration			-	-	-	-	-	-	-	-
	Diadromous	Shortnose sturgeon	General Distribution	E/E	E	-	-	-	-	-	-	-	-
	Diadromous	Striped bass	Migration			-	-	-	-	-	-	-	-
1563	Diadromous	Alewife	Migration			-	-	-	-	-	-	-	-
	Diadromous	Atlantic sturgeon	General Distribution	-E	E	-	-	-	-	-	-	-	-
	Diadromous	Blueback herring	Migration			-	-	-	-	-	-	-	-
	Diadromous	Shortnose sturgeon	General Distribution	E/E	E	-	-	-	-	-	-	-	-

REPTILES & AMPHIBIANS

<i>Map ID</i>	<i>Subelement</i>	<i>Species</i>	<i>Mapping Qualifier</i>	<i>MD/VA</i>	<i>S</i>	<i>F</i>	<i>Concentration</i>	<i>J F M A M J J A S O N D</i>	<i>Nest</i>	<i>Hatch</i>	<i>Internest</i>	<i>Juveniles</i>	<i>Adults</i>
1775	Turtle	K. ridley sea turtle	General Distribution	E/E	E	E	Uncommon	-	-	-	-	Nov-Dec	-
	Turtle	K. ridley sea turtle	General Distribution	E/E	E	E	Common	-	-	-	-	Apr-Oct	-
	Turtle	Loggerhead sea turtle	General Distribution	T/T	T	T	Abundant	-	-	-	-	Apr-Oct	Apr-Oct
	Turtle	Loggerhead sea turtle	General Distribution	T/T	T	T	Common	-	-	-	-	Nov-Dec	Nov-Dec

INVERTEBRATES

<i>Map ID</i>	<i>Subelement</i>	<i>Species</i>	<i>Mapping Qualifier</i>	<i>MD/VA</i>	<i>S</i>	<i>F</i>	<i>Concentration</i>	<i>J F M A M J J A S O N D</i>	<i>Spawn</i>	<i>Eggs</i>	<i>Larvae</i>	<i>Juveniles</i>	<i>Adults</i>
1788	Crab	Blue crab	Concentration Area	-	-	-	Highly Abundant-Summ	-	-	-	-	Jun-Oct	Jun-Oct

MARINE MAMMALS

<i>Map ID</i>	<i>Subelement</i>	<i>Species</i>	<i>Mapping Qualifier</i>	<i>MD/VA</i>	<i>S</i>	<i>F</i>	<i>Concentration</i>	<i>J F M A M J J A S O N D</i>	<i>Mating</i>	<i>Calving</i>	<i>Pupping</i>	<i>Molt</i>	<i>-</i>
1844	Manatee	West Indian manatee	General Distribution	-/E	E	-	Rare	-	-	-	-	-	-

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)

FISH

<i>Subelement</i>	<i>Species</i>	<i>Monthly Presence</i>											
		<i>J</i>	<i>F</i>	<i>M</i>	<i>A</i>	<i>M</i>	<i>J</i>	<i>J</i>	<i>A</i>	<i>S</i>	<i>O</i>	<i>N</i>	<i>D</i>
Diadromous	Alewife	-	-	-	-	-	-	-	-	-	-	-	-
	American eel	-	-	-	-	-	-	-	-	-	-	-	-
	Blueback herring	-	-	-	-	-	-	-	-	-	-	-	-
	Gizzard shad	-	-	-	-	-	-	-	-	-	-	-	-
	Striped bass	-	-	-	-	-	-	-	-	-	-	-	-
Estuarine Nursery	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-
	Atlantic menhaden	-	-	-	-	-	-	-	-	-	-	-	-
	Black drum	-	-	-	-	-	-	-	-	-	-	-	-
	Black sea bass	-	-	-	-	-	-	-	-	-	-	-	-
	Bluefish	-	-	-	-	-	-	-	-	-	-	-	-
	Kingfishes	-	-	-	-	-	-	-	-	-	-	-	-
	Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-
	Pigfish	-	-	-	-	-	-	-	-	-	-	-	-
	Red drum	-	-	-	-	-	-	-	-	-	-	-	-
	Sheepshead	-	-	-	-	-	-	-	-	-	-	-	-
	Spot	-	-	-	-	-	-	-	-	-	-	-	-
	Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-
	Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-
	Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-
	Weakfish	-	-	-	-	-	-	-	-	-	-	-	-
	White perch	-	-	-	-	-	-	-	-	-	-	-	-
Estuarine Resident	Winter flounder	-	-	-	-	-	-	-	-	-	-	-	-
	Pipefish	-	-	-	-	-	-	-	-	-	-	-	-
	Silversides	-	-	-	-	-	-	-	-	-	-	-	-

Central Bay: Map 60B

FISH (continued)

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Fish	Anchovies	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Striped burrfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
Marine Benthic	Atlantic spadefish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Clearnose skate	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Red hake	-	-	-	-	-	-	-	-	-	-	-	-	Sep-May*
	Silver hake	-	-	-	-	-	-	-	-	-	-	-	-	Oct-May
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Marine Pelagic	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Sep
	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Cobia	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Lined seahorse	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Spanish mackerel	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		-	-	-	-	-	-	-	-	-	-	-	-	May-Sep

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Bivalve	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Cephalopod	Squid	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
Crab	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Brown shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
Shrimp	Mantis shrimp	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	White shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct

MARINE MAMMALS

Subelement	Species	Monthly Presence												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
Dolphin	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES

ESI POLYGON HABITAT TYPES	
	Habitat Classification
10A	Salt and Brackish Water Marshes
10B	Freshwater Marshes
10C	Swamps
10D	Scrub and Shrub Wetlands

ESI SHORELINE HABITAT TYPES	
	Shoreline Habitat Classification
10A	Salt and Brackish Water Marshes
10B	Freshwater Marshes
10C	Swamps
9B	Vegetated Low Banks
8B	Sheltered, Solid Man-Made Structures
8C	Sheltered Riprap

10A




Total ESI Shoreline: 43,133.71
Total Shoreline: 43,114.35

Total ESI Shoreline: 43,133.71
Total Shoreline: 43,114.35

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%.

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 60C
Central Bay



SEE BACK OF MAP
for details about mapped species and
other species that occur in mapped area.
Data Published: September 2016

Map 60C Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

	Species	Mapping Qualifier	MDVA	Concentration	Monthly Presence
Subelement	Submerged aquatic veg	High Ecological Value	J F M A M J J A S O N D	Ephemeral	-
SAV	Submerged aquatic veg	High Ecological Value		High	-
SAV	Submerged aquatic veg	High Ecological Value		Low	-

BIRDS

	Species	Mapping Qualifier	MDVA	Concentration	Monthly Presence
Map ID Subelement	Nesting	J F M A M J J A S O N D	1 Nest	Molt	-
798 Raptor	Bald eagle		10S	-	-
1133 Diving	Red-throated loon		100S	-	-
Pelagic	Northern gannet		10S	-	-
Waterfowl	American black duck		100S	-	-
Waterfowl	Bufflehead		100S	-	-
Waterfowl	Canada goose		100S	-	-
Waterfowl	Mallard		10S	-	-
Waterfowl	Mergansers		10S	-	-
Waterfowl	Ruddy duck		10S	-	-
Waterfowl	Tundra swan		10S	-	-
1134 Diving	Red-throated loon	Wintering	10S	-	-
Pelagic	Northern gannet	Wintering	100S	-	-
Waterfowl	American black duck	Wintering	Present	Feb-Apr	Aug-Oct
Waterfowl	Blue-winged teal	Migration	10S	-	-
Waterfowl	Bufflehead	Wintering	100S	-	-
Waterfowl	Canada goose	Wintering	1,000S	-	-
Waterfowl	Canvasback	Wintering	10S	-	-
Waterfowl	Common goldeneye	Wintering	Present	-	-
Waterfowl	Green-winged teal	Wintering	10S	-	-
Waterfowl	Mallard	Wintering	10S	-	-
Waterfowl	Mergansers	Wintering	100S	-	-
Waterfowl	Redhead	Wintering	100S	-	-
Waterfowl	Ring-necked duck	Wintering	10S	-	-
Waterfowl	Ruddy duck	Wintering	1,000S	-	-
Waterfowl	Tundra swan	Wintering	Present	-	-
1135 Diving	Red-throated loon	Wintering	10S	-	-
Pelagic	Northern gannet	Wintering	100S	-	-
Waterfowl	American black duck	Wintering	10S	-	-
Waterfowl	Bufflehead	Wintering	100S	-	-
Waterfowl	Canada goose	Wintering	1,000S	-	-

BIRDS (continued)

Map ID	Subelement	Species	MD/VA												Monthly Presence											
			Mapping Qualifier			S	F	Concentration			J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
Waterfowl	Mallard	Wintering						Present			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Waterfowl	Mergansers	Wintering						100S			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Waterfowl	Ruddy duck	Wintering						100S			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Waterfowl	Tundra swan	Wintering						10S			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Map ID	Subelement	Species	MD/VA												Monthly Presence												
			Mapping Qualifier			S	F	Concentration			J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
1552	Diadromous	Alewife	Migration			-																		-	-	-	Feb-Apr
	Diadromous	American shad	Migration			-																		-	-	-	Aug-Nov
	Diadromous	Atlantic sturgeon	General Distribution	-E	E	-																		-	-	-	Feb-Jun
	Diadromous	Blueback herring	Migration			-																		-	-	-	Mar-Jun
	Diadromous	Hickory shad	Migration			-																		-	-	-	Feb-May
	Diadromous	Shortnose sturgeon	General Distribution	E/E	E	-																		-	-	-	Jan-Dec
	Diadromous	Striped bass	Migration			-																		-	-	-	Feb-Jun
	Estuarine Nursery	Atl. sharpnose shark	Nursery Area			-																		-	-	-	Jun-Sep
	Marine Pelagic	Sandbar shark	Nursery Area			-																		-	-	-	May-Sep
1561	Diadromous	Alewife	Migration			-																		-	-	-	Aug-Nov
	Diadromous	Atlantic sturgeon	General Distribution	-E	E	-																	-	-	-	Jan-Dec	
	Diadromous	Blueback herring	Migration			-																	-	-	-	Aug-Nov	
	Diadromous	Shortnose sturgeon	General Distribution	E/E	E	-																	-	-	-	Jan-Dec	
	Diadromous	Striped bass	Migration			-																	-	-	-	Feb-Jun	

REPTILES & AMPHIBIANS

Map ID	Subelement	Species	MD/VA												Monthly Presence											
			Mapping Qualifier			S	F	Concentration			J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles
1775	Turtle	K. ridley sea turtle	General Distribution	E/E	E	Uncommon					-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Turtle	K. ridley sea turtle	General Distribution	E/E	E	Common					-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Turtle	Loggerhead sea turtle	General Distribution	T/T	T	Common					-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Turtle	Loggerhead sea turtle	General Distribution	T/T	T	Abundant					-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

INVERTEBRATES

Map ID	Subelement	Species	MD/VA												Monthly Presence											
			Mapping Qualifier			S	F	Concentration			J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles
1788	Crab	Blue crab	Concentration Area																							
1814	Bivalve	Eastern elliptio	Vulnerable Occurrence				-																			

MARINE MAMMALS

Map ID	Subelement	Species	MD/VA												Monthly Presence												
			Mapping Qualifier			S	F	Concentration			J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatching	Calving	Pupping	Molt
1844	Manatee	West Indian manatee	General Distribution	-E	E	Rare																					

BIRDS

	Subelement	Species	<i>Monthly Presence</i>												Molt
			J	F	M	A	M	J	J	A	S	O	N	D	
	Diving	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-	-
	Gull/Tern	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-	-
		Gulls	-	-	-	-	-	-	-	-	-	-	-	-	-
			<i>Monthly Presence</i>												
			J	F	M	A	M	J	J	A	S	O	N	D	
	Diacromous	Alewife	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Apr
		American eel	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Oct
		Blueback herring	-	-	-	-	-	-	-	-	-	-	-	-	Jan-May
		Gizzard shad	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Oct
		Striped bass	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Estuarine Nursery	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		Atlantic menhaden	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		Black drum	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		Black sea bass	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
		Bluefish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
		Kingfishes	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
		Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
		Pigfish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
		Red drum	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
		Sheepshead	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
		Spot	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
		Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
		Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
		Weakfish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
		Winter flounder	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Estuarine Resident	Pipefish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
		Silversides	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
		Anchoovies	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
		Striped burrfish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Marine Benthic	Atlantic spadefish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
		Cleinnose skate	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
		Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
		Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
		Red hake	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
		Silver hake	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Marine Pelagic	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
		Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
		Cobia	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
		Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
		Lined seahorse	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
		Spanish mackerel	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Bivalve	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Stout razor clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Cephalopod	Squid	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
Crab	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Horseshoe crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Shrimp	Brown shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Mantis shrimp	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	White shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct

MARINE MAMMALS

Subelement	Species	Monthly Presence												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
Dolphin	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-
	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-
	Harbor porpoise	-	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES


ESI POLYGON HABITAT TYPES		Habitat Classification	Area (Acres)	Area (Sq. Miles)
ESI/Rank				
10A		Salt and Brackish Water Marshes	75.97	0.12
10B		Freshwater Marshes	39.26	0.06
10C		Swamps	1,151.93	1.80
10D		Scrub and Shrub Wetlands	87.18	0.14

ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
ESI/Rank					
10A		Salt and Brackish Water Marshes	19,643.33	12.21	19%
10B		Freshwater Marshes	295.76	0.18	< 1%
10C		Swamps	715.86	0.44	1%
10D		Scrub and Shrub Wetlands	1,050.70	0.65	1%
9B		Vegetated Low Banks	53,821.08	33.44	53%
8B		Sheltered, Solid Man-Made Structures	11,353.52	7.05	11%
8C		Sheltered Riprap	5,444.40	3.38	5%
7		Exposed Tidal Flats	2,255.08	1.40	2%
6B		Riprap	2,022.80	1.26	2%
4		Coarse Grained Sand Beaches	2,348.44	1.46	2%
3B		Scarps and Steep Slopes (Sand)	297.50	0.18	< 1%
1B		Exposed, Solid Man-Made Structures	2,063.12	1.28	2%
Total ESI Shoreline:		101,311.59	Total ESI Shoreline:	62.95	
Total Shoreline:		94,744.33	Total Shoreline:	58.87	

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 61
Central Bay

HUMAN USE RESOURCES

DISPLAYED ON MAP (POINTS)			Name	Contact	Phone
Map ID	Type				
1	AQUACULTURE		OYSTER COMMERCIAL ELDERT RF#01-1093	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
2	AQUACULTURE		OYSTER NON-COMM CHES BAY GOVERNOR'S RF#2000-1731	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
3	AQUACULTURE		OYSTER NON-COMM GRIM RF#98-1625	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
4	AQUACULTURE		OYSTER NON-COMM PARK, GEORGE M. RF#98-1937	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
5	AQUACULTURE		OYSTER NON-COMM RIGTERINK, RAY RF#98-0758	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
6	AQUACULTURE		OYSTER NON-COMM STEELE RF#98-2021	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
7	BOAT RAMP		SIMONSON LANDING	LENE PENNINGTON, VDGIF	804-367-0058
8	BOAT RAMP		TOTUSKEY	LENE PENNINGTON, VDGIF	804-367-0058
9	EPA FACILITY		HAYNESVILLE, VA 326	MANAGER	804-333-4234
10	REPEATED MEASUREMENT SITE		WQ - STATION LE3.1	CHESAPEAKE BAY PROGRAM	800-968-7229
11	REPEATED MEASUREMENT SITE		WQ - STATION RET3.2	CHESAPEAKE BAY PROGRAM	800-968-7229
DISPLAYED ON MAP (POLYGONS)			Name	Contact	Phone
Map ID	Type				
12	PARK		BELLE ISLE STATE PARK	VA DEPT OF CONSERVATION AND RECREATION	
13	WILDLIFE REFUGE		RAPPAHANNOCK RIVER VALLEY NWR	UNITED STATES FISH AND WILDLIFE SERVICE	804-333-1470
ALSO PRESENT IN MAPPED AREA (POLYGONS)			Name	Contact	Phone
Type					
STATE PROTECTED AREA			CHILTON WOODS	VA DEPT OF FORESTRY	

JURISDICTIONS
COUNTY: ESSEX CO., LANCASTER CO., MIDDLESEX CO., NORTHUMBERL. CO., RICHMOND CO.
COAST GUARD: DISTRICT 5, SECTOR HAMPTON ROADS
USACE: NORTH ATLANTIC DIVISION, BALTIMORE DISTRICT, NORFOLK DISTRICT

FEMA: REGION I
EPA: REGION 3

SHORELINE RESOURCES

ESI POLYGON HABITAT TYPES	
ESI Rank	Habitat Classification
10A	Salt and Brackish Water Marshes
10B	Freshwater Marshes
10C	Swamps
10D	Scrub and Shrub Wetlands
9A	Sheltered Tidal Flats
7	Exposed Tidal Flats



ESI SHORELINE HABITAT TYPES

ESI Shoreline Habitat Classification	
ESI Rank	Shoreline Habitat Classification
10A	Salt and Brackish Water Marshes
10B	Freshwater Marshes
10C	Swamps
10D	Scrub and Shrub Wetlands
9A	Sheltered Tidal Flats
9B	Vegetated Low Banks
8B	Sheltered, Solid Man-Made Structures
8C	Sheltered Riprap
7	Exposed Tidal Flats
6B	Riprap
4	Coarse Grained Sand Beaches
3B	Scarps and Steep Slopes (Sand)
1B	Exposed, Solid Man-Made Structures



Total ESI Shoreline:	Length (Meters)	% of ESI Shoreline:
238,722.51	127,499.09	53%
Total Shoreline:	79.22	
218,596.89		


Total ESI Shoreline:	Area (Acres)	% of ESI Shoreline:
238,722.51	1,385.81	53%
Total Shoreline:	180.56	
218,596.89		

Total ESI Shoreline:	Area (Sq. Miles)	% of ESI Shoreline:
238,722.51	2.17	53%
Total Shoreline:	0.28	
218,596.89		

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%.

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 61A
Central Bay



SEE BACK OF MAP
for details about mapped species and
other species that occur in mapped area.
Data Published: September 2016

0 Not for Navigation 1 Miles
0 1 Kilometers 1:50,000



Map 61A Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BIRDS

Map ID	Subelement	Species	Mapping Qualifier	S	F	Concentration	Monthly Presence													
							J	F	M	A	M	J	J	A	S	O	N	D	Nest	
																				Mig.(S)
232	Waterfowl	Canada goose	Nesting			Present														Mar-Aug
	Waterfowl	Mallard	Nesting			Present														Mar-Aug
267	Waterfowl	American black duck	Wintering																	
	Waterfowl	Canada goose	Nesting			Present														Mar-Aug
	Waterfowl	Canada goose	Wintering																	
	Waterfowl	Mallard	Nesting			Present														Mar-Aug
	Waterfowl	Mallard	Wintering																	
	Waterfowl	Snow goose	Wintering																	
798	Raptor	Bald eagle	Nesting																	Dec-Jun
806	Waterfowl	Bufflehead	Wintering																	
	Waterfowl	Canada goose	Wintering																	
	Waterfowl	Mallard	Wintering																	
	Waterfowl	Tundra swan	Wintering																	
807	Waterfowl	American black duck	Wintering																	
	Waterfowl	Bufflehead	Wintering																	
	Waterfowl	Canada goose	Wintering																	
	Waterfowl	Mallard	Wintering																	
	Waterfowl	Tundra swan	Wintering																	
871	Waterfowl	Bufflehead	Wintering																	
	Waterfowl	Canvasback	Wintering																	
	Waterfowl	Mergansers	Wintering																	
	Waterfowl	Ruddy duck	Wintering																	
892	Waterfowl	American black duck	Wintering																	
	Waterfowl	Canada goose	Wintering																	
	Waterfowl	Mallard	Wintering																	
	Waterfowl	Scaup	Wintering																	
	Waterfowl	Canada goose	Wintering																	
	Waterfowl	Mallard	Wintering																	
893	Waterfowl	Snow goose	Wintering																	
	Waterfowl	Common loon	Wintering																	
	Diving	Grebes	Wintering																	
1020	Diving	Red-throated loon	Wintering																	
	Waterfowl	American black duck	Wintering																	
	Waterfowl	Brant	Wintering																	
	Waterfowl	Bufflehead	Wintering																	
	Waterfowl	Canada goose	Wintering																	

BIRDS (continued)

Map ID	Subelement	Species	MD\VA			Monthly Presence										Nest		Mig.(S)		Mig.(F)	
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)
Waterfowl	Canvasback	Wintering	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Waterfowl	Long-tailed duck	Wintering	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Waterfowl	Mallard	Wintering	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Waterfowl	Mergansers	Wintering	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Waterfowl	Ruddy duck	Wintering	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Waterfowl	Scaup	Wintering	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Waterfowl	Scoters	Wintering	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Waterfowl	Tundra swan	Wintering	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1072	Waterfowl	American black duck	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Map ID	Subelement	Species	MD\VA			Monthly Presence										Nest		Eggs		Larvae		Juveniles		Adults	
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Eggs	Larvae	Juveniles	Adults	Adults	
1556	Diadromous	Atlantic sturgeon	General Distribution	-IE	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	Diadromous	Shortnose sturgeon	General Distribution	E/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	Diadromous	Striped bass	Nursery Area	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

REPTILES & AMPHIBIANS

Map ID	Subelement	Species	MD\VA			Monthly Presence										Nest		Hatch		Internest		Juveniles		Adults	
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles	Adults	Adults	
1767	Turtle	Green sea turtle	General Distribution	T/T	T	Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	Turtle	Green sea turtle	General Distribution	T/T	T	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

INVERTEBRATES

Map ID	Subelement	Species	MD\VA			Monthly Presence										Nest		Eggs		Larvae		Juveniles		Adults	
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Eggs	Larvae	Juveniles	Adults	Adults	
1788	Crab	Blue crab	Concentration Area	-	-	Highly Abundant-Summ	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

MARINE MAMMALS

Map ID	Subelement	Species	MD\VA			Monthly Presence										Nest		Mating		Calving		Pupping		Molt	
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mating	Calving	Pupping	Molt	Molt	
1844	Manatee	West Indian manatee	General Distribution	-IE	E	Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		

BIRDS

Subelement	Species	Monthly Presence									
		J	F	M	A	M	J	J	A	S	O
Diving	Brown pelican	-	-	-	-	-	-	-	-	-	-
	D. crested cormorant	-	-	-	-	-	-	-	-	-	-
Gull/Tern	Gulls	-	-	-	-	-	-	-	-	-	-
	Terns	-	-	-	-	-	-	-	-	-	-
Waterfowl	Wood duck	-	-	-	-	-	-	-	-	-	-
		Apr-Jul	-	-	-	-	-	-	-	-	-

FISH

	Subelement	Species	Monthly Presence												Adults
			J	F	M	A	M	J	J	A	S	O	N	D	
	Diadromous	Alewife	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Apr
		American eel	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct
		Blueback herring	-	-	-	-	-	-	-	-	-	-	-	-	Jan-May
		Striped bass	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Jun
	Estuarine Nursery	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		Atlantic herring	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
		Atlantic menhaden	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Jun
		Bluefish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		Kingfishes	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
		Red drum	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
		Spot	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
		Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Apr
		Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
		Weakfish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Estuarine Resident	White perch	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		Pipefish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		Silversides	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Fish	Anchoovies	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Freshwater	Blue catfish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		Bullhead catfish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		Channel catfish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		Longnose gar	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		Spottail shiner	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		Sunfish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		Tasseled darter	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		Yellow perch	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Marine Benthic	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Sep
		Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Dec-Jun
	Marine Pelagic	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Nov
		Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
			-	-	-	-	-	-	-	-	-	-	-	-	May-Nov

INVERTEBRATES

	Subelement	Species	Monthly Presence												Adults
			J	F	M	A	M	J	J	A	S	O	N	D	
	Bivalve	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Crab	Stout razor clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Shrimp	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		Brown shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
		White shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct

For additional information about species locations and extent, reference the underlying GIS data available from restoration.noaa.gov

SHORELINE RESOURCES

ESI POLYGON HABITAT TYPES		Habitat Classification	Area (Acres)	Area (Sq. Miles)
ESI Rank				
10A		Salt and Brackish Water Marshes	482.80	0.75
10B		Freshwater Marshes	49.78	0.08
10C		Swamps	808.36	1.26
10D		Scrub and Shrub Wetlands	212.00	0.33
9A		Sheltered Tidal Flats	1.94	0.00


ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
ESI Rank					
10A		Salt and Brackish Water Marshes	46,398.18	28.83	73%
10B		Freshwater Marshes	3,317.17	2.06	5%
10C		Swamps	670.80	0.42	1%
10D		Scrub and Shrub Wetlands	1,884.50	1.17	3%
9A		Sheltered Tidal Flats	452.04	0.28	1%
9B		Vegetated Low Banks	9,568.60	5.95	15%
8B		Sheltered, Solid Man-Made Structures	577.23	0.36	1%
8C		Sheltered Riprap	272.78	0.17	<1%

Total ESI Shoreline: 63,141.29 Total ESI Shoreline: 39.23
 Total Shoreline: 62,141.64 Total Shoreline: 38.61

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%.

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 61B
Central Bay



SEE BACK OF MAP
for details about mapped species and
other species that occur in mapped area.
Data Published: September 2016

Map 61B Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BIRDS

Map ID	Subelement	Species	Mapping Qualifier	MD/VA	S	F	Concentration	Monthly Presence															
								J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
34	Waterfowl	Scoters	Wintering																	34 Nests	Feb-Jun	-	-
106	Wading	Great blue heron	Nesting																	16 Nests	Feb-Jun	-	-
196	Wading	Great blue heron	Nesting																	Present	Mar-Aug	Jun-Jul	-
232	Waterfowl	Canada goose	Nesting																	Present	Mar-Aug	Jul-Aug	-
678	Waterfowl	Mallard	Wintering																	10S	-	-	-
	Waterfowl	American black duck	Wintering																	10S	-	Feb-Apr	Aug-Oct
	Waterfowl	Blue-winged teal	Migration																	100S	-	-	-
	Waterfowl	Bufflehead	Wintering																	10,000S	-	-	-
	Waterfowl	Canada goose	Wintering																	100S	-	-	-
	Waterfowl	Canvasback	Wintering																	10S	-	-	-
	Waterfowl	Gadwall	Wintering																	10S	-	-	-
	Waterfowl	Green-winged teal	Wintering																	10S	-	-	-
	Waterfowl	Mallard	Wintering																	100S	-	-	-
	Waterfowl	Mergansers	Wintering																	10S	-	-	-
	Waterfowl	Ruddy duck	Wintering																	1,000S	-	-	-
	Waterfowl	Scaup	Wintering																	100S	Present	Dec-Jun	Feb-Apr
	Waterfowl	Snow goose	Wintering																	Present	100S	-	-
	Waterfowl	Tundra swan	Wintering																	10S	-	-	-
680	Waterfowl	American black duck	Wintering																	10,000S	-	-	-
	Waterfowl	Canada goose	Wintering																	100S	-	-	-
	Waterfowl	Mallard	Wintering																	100S	-	-	-
	Waterfowl	Ruddy duck	Wintering																	1,000S	-	-	-
	Waterfowl	Tundra swan	Wintering																	100S	-	-	-
	Raptor	Bald eagle	Nesting																1 Nest	-	Dec-Jun	Feb-Apr	
910	Waterfowl	Blue-winged teal	Migration																	10S	-	Aug-Oct	-
	Waterfowl	Bufflehead	Wintering																	100S	-	-	-
	Waterfowl	Canvasback	Wintering																	100S	-	-	-
	Waterfowl	Gadwall	Wintering																	10S	-	-	-
	Waterfowl	Green-winged teal	Wintering																	10S	-	-	-
	Waterfowl	Mergansers	Wintering																	10S	-	-	-
	Waterfowl	Scaup	Wintering																	100S	Present	10S	-
	Waterfowl	Snow goose	Wintering																	100S	-	-	-
955	Waterfowl	Brant	Wintering																	1,000S	-	-	-
	Waterfowl	Bufflehead	Wintering																	1,000S	-	-	-
	Waterfowl	Canvasback	Wintering																	1,000S	-	-	-
	Waterfowl	Mergansers	Wintering																	1,000S	-	-	-
	Waterfowl	Scaup	Wintering																	1,000S	-	-	-

BIRDS (continued)

Map ID	Subelement	Species	MD\VA						Monthly Presence												
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
1020	Diving	Common loon	Wintering															-	-	-	-
	Diving	Grebes	Wintering															-	-	-	-
	Diving	Red-throated loon	Wintering															-	-	-	-
	Watrfowl	American black duck	Wintering															-	-	-	-
	Watrfowl	Brant	Wintering															-	-	-	-
	Watrfowl	Bufflehead	Wintering															-	-	-	-
	Watrfowl	Canada goose	Wintering															-	-	-	-
	Watrfowl	Canvasback	Wintering															-	-	-	-
	Watrfowl	Long-tailed duck	Wintering															-	-	-	-
	Watrfowl	Mallard	Wintering															-	-	-	-
	Watrfowl	Mergansers	Wintering															-	-	-	-
	Watrfowl	Ruddy duck	Wintering															-	-	-	-
	Watrfowl	Scaup	Wintering															-	-	-	-
	Watrfowl	Scoters	Wintering															-	-	-	-
	Watrfowl	Tundra swan	Wintering															-	-	-	-
1022	Diving	Common loon	Wintering															-	-	-	-
	Diving	Grebes	Wintering															-	-	-	-
	Diving	Red-throated loon	Wintering															-	-	-	-
	Watrfowl	Long-tailed duck	Wintering															-	-	-	-
1194	Diving	D. crested cormorant	Wintering															-	-	-	-
	Watrfowl	Scoters	Wintering															-	-	-	-

FISH

Map ID	Subelement	Species	MD\VA						Monthly Presence												
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Eggs	Larvae	Juveniles	Adults
1599	Diadromous	Alewife	Migration															-	-	-	-
	Diadromous	American shad	Migration															-	-	-	-
	Diadromous	Atlantic sturgeon	Migration															-	-	-	-
	Diadromous	Blueback herring	Migration															-	-	-	-
	Diadromous	Hickory shad	Migration															-	-	-	-
1620	Diadromous	Alewife	Migration															-	-	-	-
	Diadromous	Blueback herring	Migration															-	-	-	-

INVERTEBRATES

Map ID	Subelement	Species	MD\VA						Monthly Presence												
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Eggs	Larvae	Juveniles	Adults
1788	Crab	Blue crab	Highly Abundant-Summ															-	-	-	-
1789	Crab	Blue crab	Highly Abundant															-	-	-	-
1813	Bivalve	Eastern elliptio	Vulnerable Occurrence															-	-	-	-
1824	Bivalve	Eastern floater	Vulnerable Occurrence															-	-	-	-
1834	Crab	Blue crab	Concentration Area															-	-	-	-

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

FISH

Subelement	Species	MD\VA												Monthly Presence											
		Mapping Qualifier			S F			Concentration			J F M A M J J A S O N D			Eggs			Larvae			Juveniles			Adults		
Diadromous	Atlantic sturgeon	General Distribution	-E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec
	Shortnose sturgeon	General Distribution	E/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec
	Striped bass	Nursery Area	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

REPTILES & AMPHIBIANS

Subelement	Species	MD\VA												Monthly Presence												
		Mapping Qualifier			S F			Concentration			J F M A M J J A S O N D			Nest			Hatch			Internest			Juveniles			Adults
	Green sea turtle	General Distribution	T/T	T	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Green sea turtle	General Distribution	T/T	T	Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Dec*

MARINE MAMMALS

Subelement	Species	MD\VA												Monthly Presence											
		Mapping Qualifier			S F			Concentration			J F M A M J J A S O N D			Mating			Calving			Pupping			Molt		
	West Indian manatee	General Distribution	-E	E	Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)


BIRDS

Subelement	Species	MD\VA												Monthly Presence											
		Mapping Qualifier			S F			Concentration			J F M A M J J A S O N D			Nest			Mig.(S)			Mig.(F)			Molt		
Diving	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Gull/Tern	Gulls	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Terns	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Waterfowl	Wood duck	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Subelement	Species	MD\VA												Monthly Presence												
		Mapping Qualifier			S F			Concentration			J F M A M J J A S O N D			Spawn			Eggs			Larvae			Juveniles			Adults
Diadromous	Alewife	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Apr	Feb-Apr
	American eel	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-May	Jan-Dec
	Blueback herring	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct	Mar-Jun
	Striped bass	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec
Estuarine Nursery	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct	Jan-Jun
	Atlantic herring	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Jun	Jan-Jun
	Atlantic menhaden	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec	Jan-Dec
	Black drum	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec	Apr-Dec
	Bluefish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct	May-Oct
	Kingfishes	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov	Apr-Nov
	Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Oct	Mar-Oct
	Pigfish	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov	Apr-Nov

FISH (continued)

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Red drum	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Spot	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Weakfish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	White perch	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Pipefish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Silversides	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Silversides	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Dec
	Anchovies	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Freshwater	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Blue catfish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Bullhead catfish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Spottail shiner	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Sunfish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Sep
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Nov
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Lined seahorse	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Spanish mackerel	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Stout razor clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Brown shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Mantis shrimp	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	White shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct

MARINE MAMMALS

Subelement	Species	Monthly Presence												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov


ESI POLYGON HABITAT TYPES		Habitat Classification	Area (Acres)	Area (Sq. Miles)
ESI Rank				
10A		Salt and Brackish Water Marshes	437.47	0.68
10B		Freshwater Marshes	74.02	0.12
10C		Swamps	1,097.12	1.71
10D		Scrub and Shrub Wetlands	612.40	0.96

ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
ESI Rank					
10A		Salt and Brackish Water Marshes	40,964.04	25.45	52%
10C		Swamps	1,322.21	0.82	2%
10D		Scrub and Shrub Wetlands	114.01	0.07	< 1%
9B		Vegetated Low Banks	27,218.10	16.91	35%
8B		Sheltered, Solid Man-Made Structures	2,734.39	1.70	4%
8C		Sheltered Riprap	1,387.01	0.86	2%
6B		Riprap	1,928.81	1.20	2%
4		Coarse Grained Sand Beaches	1,039.74	0.65	1%
3B		Scars and Steep Slopes (Sand)	938.50	0.58	1%
1B		Exposed, Solid Man-Made Structures	391.46	0.24	1%
Total ESI Shoreline:			78,038.27	Total ESI Shoreline:	48.49
Total Shoreline:			73,314.13	Total Shoreline:	45.56

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 61C
Central Bay



SEE BACK OF MAP
for details about mapped species and
other species that occur in mapped area.
Data Published: September 2016

Map 61C Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BIRDS

Map ID	Subelement	Species	Mapping Qualifier	S	F	Concentration	Monthly Presence											
							J	F	M	A	M	J	J	A	S	O	N	D
126	Wading	Great blue heron	Nesting			3 Nests	-	-	-	-	-	-	-	-	-	-	-	-
188	Wading	Great blue heron	Nesting			24 Nests	-	-	-	-	-	-	-	-	-	-	-	-
232	Watertowl	Canada goose	Nesting			Present	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Mallard	Nesting			Present	-	-	-	-	-	-	-	-	-	-	-	-
798	Raptor	Bald eagle	Nesting			1 Nest	-	-	-	-	-	-	-	-	-	-	-	-
910	Watertowl	Blue-winged teal	Migration			10S	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Bufflehead	Wintering			100S	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Canvasback	Wintering			100S	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Gadwall	Wintering			10S	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Green-winged teal	Wintering			10S	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Mergansers	Wintering			100S	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Scaup	Wintering			100S	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Snow goose	Wintering			Present	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Canvasback	Wintering			100S	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Redhead	Wintering			10S	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Ruddy duck	Wintering			100S	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Scaup	Wintering			10S	-	-	-	-	-	-	-	-	-	-	-	-
923	Watertowl	Canada goose	Wintering			1,000S	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Tundra swan	Wintering			Present	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Map ID	Subelement	Species	Mapping Qualifier	S	F	Concentration	Monthly Presence											
							J	F	M	A	M	J	J	A	S	O	N	D
	Nursery Area	Alewife	-			-	-	-	-	-	-	-	-	-	-	-	-	-
	Nursery Area	American shad	-			-	-	-	-	-	-	-	-	-	-	-	-	-
	Nursery Area	Blueback herring	-			-	-	-	-	-	-	-	-	-	-	-	-	-

INVERTEBRATES

Map ID	Subelement	Species	Mapping Qualifier	S	F	Concentration	Monthly Presence											
							J	F	M	A	M	J	J	A	S	O	N	D
1614	Diadromous	Alewife	Highly Abundant-Wint	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Diadromous	American shad	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Diadromous	Blueback herring	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

BIRDS

Subelement	Species	Mapping Qualifier	MD/VA						Monthly Presence											
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)
Diving	Common loon	Wintering	-	-	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Grebes	Wintering	-	-	Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Red-throated loon	Wintering	-	-	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Waterfowl	American black duck	Wintering	-	-	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Brant	Wintering	-	-	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Bufflehead	Wintering	-	-	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Canada goose	Wintering	-	-	10,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Canvasback	Wintering	-	-	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Long-tailed duck	Wintering	-	-	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Mallard	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Mergansers	Wintering	-	-	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Ruddy duck	Wintering	-	-	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Scaup	Wintering	-	-	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Scooter's	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Tundra swan	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Subelement	Species	Mapping Qualifier	MD/VA						Monthly Presence											
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Eggs	Larvae	Juveniles
Diadromous	Alewife	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	American shad	Migration	-	-	General Distribution	-E	E	-	-	-	-	-	-	-	-	-	-	-	-	-
	Atlantic sturgeon	Migration	-E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Atlantic sturgeon	Migration	-E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Blueback herring	Migration	-	-	General Distribution	E/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-
	Hickory shad	Migration	-	-	Nursery Area	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Shortnose sturgeon	Migration	-	-	Striped bass	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

REPTILES & AMPHIBIANS

Subelement	Species	Mapping Qualifier	MD/VA						Monthly Presence											
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Eggs	Larvae	Juveniles
Turtle	Green sea turtle	General Distribution	T/T	T	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Green sea turtle	General Distribution	T/T	T	Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

INVERTEBRATES

Subelement	Species	Mapping Qualifier	MD/VA						Monthly Presence											
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Eggs	Larvae	Juveniles
Crab	Blue crab	Highly Abundant-Summ	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Blue crab	Highly Abundant	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

MARINE MAMMALS

Subelement	Species	Mapping Qualifier	MD/VA	S F	Concentration	J F M A M J J A S O N D	Mating	Calving	Pupping	Molt
Manatee	West Indian manatee	General Distribution	-IE	E	Rare					-

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)

BIRDS

Subelement	Species	Mapping Qualifier	MD/VA	S F	Concentration	J F M A M J J A S O N D	Nest	Mig.(S)	Mig.(F)	Molt
Diving	Brown pelican	General Distribution	-IE	E	Rare					-
	D. crested cormorant									-
Gull/Tern	Gulls									-
	Terns									-
Watertowl	Wood duck						Apr-Jul		Jul-Aug	-

FISH

Subelement	Species	Mapping Qualifier	MD/VA	S F	Concentration	J F M A M J J A S O N D	Spawn	Eggs	Larvae	Juveniles	Adults
Diacromous	Alewife	General Distribution	-IE	E	Rare		Feb-Apr	Feb-Apr	Feb-Apr	Apr-Oct	Feb-Apr
	American eel									Jan-May	Jan-Dec
	Blueback herring						Mar-Jun	Mar-Jun	Apr-Jun	May-Oct	Mar-Jun
	Striped bass									Jan-Dec	Jan-Dec
Estuarine Nursery	Atlantic croaker	General Distribution	-IE	E	Rare					Apr-Dec	May-Oct
	Atlantic herring									-	Jan-Jun
	Atlantic menhaden									Jan-Dec	Jan-Dec
	Bluefish									May-Oct	May-Oct
	Kingfishes									-	Apr-Nov
	Red drum									-	May-Nov
	Spot									Nov-Apr	Apr-Dec
	Spotted seatrout									Apr-Nov	Apr-Nov
	Summer flounder									Apr-Oct	Jan-Dec
	Weakfish									Jun-Dec	Apr-Dec
	White perch									Jan-Dec	Jan-Dec
Estuarine Resident	Pipefish	General Distribution	-IE	E	Rare					Jan-Dec	Jan-Dec
	Silversides									-	Jan-Dec
	Anchoovies									-	Jan-Dec
Fish	Blue catfish	General Distribution	-IE	E	Rare		Apr-Sep	-	-	Jan-Dec	Jan-Dec
Freshwater	Bullhead catfish						May-Jun	May-Jun	May-Jun	Jan-Dec	Jan-Dec
	Spottail shiner						Apr-Jun			-	Jan-Dec
	Sunfish						May-Aug			-	Jan-Dec
	Northern searobin						Mar-Jul			-	Jan-Dec
Marine Benthic	Spotted hake	General Distribution	-IE	E	Rare					Dec-Jun	Dec-Jun
	Butterfish									Jul-Nov	Mar-Nov
	Harvestfish									-	May-Nov

REPTILES & AMPHIBIANS

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Turtle	Diamondback terrapin	■	■	■	■	■	■	■	■	■	■	■	■	Apr-Oct

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Bivalve	Eastern oyster	■	■	■	■	■	■	■	■	■	■	■	■	Jan-Dec
	Softshell clam	■	■	■	■	■	■	■	■	■	■	■	■	Jan-Dec
	Stout razor clam	■	■	■	■	■	■	■	■	■	■	■	■	Jan-Dec
Crab	Blue crab	■	■	■	■	■	■	■	■	■	■	■	■	Jan-Dec
	Brown shrimp	■	■	■	■	■	■	■	■	■	■	■	■	May-Oct
	White shrimp	■	■	■	■	■	■	■	■	■	■	■	■	May-Oct

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES

ESI POLYGON HABITAT TYPES		Habitat Classification	Area (Acres)	Area (Sq. Miles)
ESI Rank				
10A		Salt and Brackish Water Marshes	465.54	0.73
10B		Freshwater Marshes	24.88	0.04
10C		Swamps	853.68	1.33
10D		Scrub and Shrub Wetlands	356.46	0.56
7		Exposed Tidal Flats	6.76	0.01


ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
ESI Rank					
10A		Salt and Brackish Water Marshes	40,136.87	24.94	41%
10B		Freshwater Marshes	142.03	0.09	< 1%
10C		Swamps	768.87	0.48	1%
10D		Scrub and Shrub Wetlands	486.36	0.30	< 1%
9A		Sheltered Tidal Flats	1,817.24	1.13	2%
9B		Vegetated Low Banks	23,850.57	14.82	24%
8B		Sheltered, Solid Man-Made Structures	5,545.13	3.45	6%
8C		Sheltered Riprap	1,521.09	0.95	2%
7		Exposed Tidal Flats	121.16	0.08	< 1%
6B		Riprap	7,873.07	4.89	8%
4		Coarse Grained Sand Beaches	6,465.56	4.02	7%
3B		Scarps and Steep Slopes (Sand)	1,462.87	0.91	1%
1B		Exposed, Solid Man-Made Structures	7,352.11	4.57	8%

Total ESI Shoreline: 97,542.94 Total ESI Shoreline: 60.61
 Total Shoreline: 83,141.12 Total Shoreline: 51.66

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 66
Central Bay



SEE BACK OF MAP
for details about mapped resources and
other resources that occur in mapped area.
Data Published: September 2016

0 Not for Navigation 2 Miles
0 2 Kilometers

1:100,000



Map 66
Central Bay

HUMAN USE RESOURCES

DISPLAYED ON MAP (POINTS)

Map ID	Type	Name	Contact	Phone
1	AIRPORT	HUMMEL FIELD		
2	AIRPORT	LONGBRANCH		
3	AQUACULTURE	OYSTER COMMERCIAL SPECK RF#001950	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
4	AQUACULTURE	OYSTER NON-COMM BAILEY RF#001852	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
5	AQUACULTURE	OYSTER NON-COMM BATTISTA RF#00-1849	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
6	AQUACULTURE	OYSTER NON-COMM BERRY, FREDERIC A. RF#99-0692	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
7	AQUACULTURE	OYSTER NON-COMM BOOKER RF#99-1528	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
8	AQUACULTURE	OYSTER NON-COMM BRAUER, JOHN L. RF#98-0802	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
9	AQUACULTURE	OYSTER NON-COMM BUTLER, ROBERT W. RF#98-0777	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
10	AQUACULTURE	OYSTER NON-COMM CALKINS RF#001978	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
11	AQUACULTURE	OYSTER NON-COMM CARDWELL, JOHN RF#99-1922	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
12	AQUACULTURE	OYSTER NON-COMM CHAMPLAIN, O.S. RF#98-1706	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
13	AQUACULTURE	OYSTER NON-COMM CHES BAY GOVERNOR'S RF#99-2002	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
14	AQUACULTURE	OYSTER NON-COMM CONSTANTINE RF#98-2186	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
15	AQUACULTURE	OYSTER NON-COMM CORNWELL, JERRY A. JR. RF#98-1191	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
16	AQUACULTURE	OYSTER NON-COMM CRICHTON, GEORGE F. RF#98-1705	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
17	AQUACULTURE	OYSTER NON-COMM DEAVER RF#02-0215	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
18	AQUACULTURE	OYSTER NON-COMM DOUMAR RF#01-1152	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
19	AQUACULTURE	OYSTER NON-COMM DUFFER RF#990426	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
20	AQUACULTURE	OYSTER NON-COMM FUNKHouser, RAYMOND RF#98-1499	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
21	AQUACULTURE	OYSTER NON-COMM GARY, ROBERT L. RF#98-1081	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
22	AQUACULTURE	OYSTER NON-COMM HAMILTON, RUTH P. RF#98-2041	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
23	AQUACULTURE	OYSTER NON-COMM HAYES RF#002117	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
24	AQUACULTURE	OYSTER NON-COMM HERNDON, DAVID RF#99-1819	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
25	AQUACULTURE	OYSTER NON-COMM HUNTER RF#99-1414	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
26	AQUACULTURE	OYSTER NON-COMM HUNTER, LINDA M. RF#99-1383	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
27	AQUACULTURE	OYSTER NON-COMM KUEHLHORN, BERT O. RF#99-0983	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
28	AQUACULTURE	OYSTER NON-COMM LOCKERMAN RF#00-0722	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
29	AQUACULTURE	OYSTER NON-COMM LYNCH RF#002121	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
30	AQUACULTURE	OYSTER NON-COMM MAPES RF#01-0676	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
31	AQUACULTURE	OYSTER NON-COMM MCLEAN, DONALD RF#99-2098	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
32	AQUACULTURE	OYSTER NON-COMM MORRIS RF#99-1743	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
33	AQUACULTURE	OYSTER NON-COMM NOTHWANGER RF#001666	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
34	AQUACULTURE	OYSTER NON-COMM ORR, MARSHALL RF#98-1792	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
35	AQUACULTURE	OYSTER NON-COMM PORTER RF#00-2093	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
36	AQUACULTURE	OYSTER NON-COMM RHODE RF#991440	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
37	AQUACULTURE	OYSTER NON-COMM SCOTT, NORMAN RF#01-0152	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
38	AQUACULTURE	OYSTER NON-COMM SHUMADINE, WILL RF#00-0340	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
39	AQUACULTURE	OYSTER NON-COMM SMILEY RF#001896	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
40	AQUACULTURE	OYSTER NON-COMM SMITH, DOUGLAS A. RF#98-1624	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254

DISPLAYED ON MAP (POINTS) (continued)

Map ID	Type	Name	Contact	Phone
41	AQUACULTURE	OYSTER NON-COMM STEBBINS RF#00-1732	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
42	AQUACULTURE	OYSTER NON-COMM STEIN RF#001644	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
43	AQUACULTURE	OYSTER NON-COMM SUTPHEN RF#001880	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
44	AQUACULTURE	OYSTER NON-COMM TEEPLES, THOMAS C. RF#98-0828	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
45	AQUACULTURE	OYSTER NON-COMM TOWNES, JOHN M., III RF#99-0974	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
46	AQUACULTURE	OYSTER NON-COMM VIMS RF#99-1654	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
47	AQUACULTURE	OYSTER NON-COMM WALKER, DOUGLAS RF#99-1763	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
48	AQUACULTURE	OYSTER NON-COMM WALKER, HOWARD P. RF#01-1764	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
49	AQUACULTURE	OYSTER NON-COMM WICK, JOHN H., III RF#98-1809	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
50	AQUACULTURE	OYSTER NON-COMM WILLETT RF#99-2097	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
51	AQUACULTURE	OYSTER NON-COMM WILSON, DAVID RF#98-0895	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
52	AQUACULTURE	OYSTER NON-COMM WINELAND RF#99-1731	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
53	AQUACULTURE	OYSTER NON-COMM WINTER RF#99-0693	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
54	BOAT RAMP	MILL CREEK	LENE PENNINGTON, VDGIF	804-367-0058
55	HELIPORT	NICHOLLS		
56	HELIPORT	RAPPAHANNOCK GENERAL HOSPITAL		
57	HELIPORT	WHITE STONE		
58	HELIPORT	WHITE STONE FAMILY PRACTICE		
59	REPEATED MEASUREMENT SITE	DB - NOS, 8636580 - WINDMILL POINT, VA	NATIONAL COASTAL DATA DEVELOPMENT CENTER PORTS PROGRAM	866-732-2382 301-713-4392
60	REPEATED MEASUREMENT SITE	PORTS - WINDMILL POINT, RAPPAHANNOCK RIVER	CHESAPEAKE BAY PROGRAM	800-968-7229
61	REPEATED MEASUREMENT SITE	WQ - STATION LE3.4	CHESAPEAKE BAY PROGRAM	800-968-7229
62	REPEATED MEASUREMENT SITE	WQ - STATION LE3.6	CHESAPEAKE BAY PROGRAM	800-968-7229
63	REPEATED MEASUREMENT SITE	WQ - STATION LE3.6N	CHESAPEAKE BAY PROGRAM	800-968-7229
64	REPEATED MEASUREMENT SITE	WQ - STATION LE3.6S		

ALSO PRESENT IN MAPPED AREA (POLYGONS)

Type	Name	Contact	Phone
STATE PROTECTED AREA	BLUE CRAB SANCTUARY	VIRGINIA MARINE RESOURCES COMMISSION	757-247-2200
STATE PROTECTED AREA	HUGHLETT POINT	V/A DEPT OF CONSERVATION AND RECREATION	

JURISDICTIONS

COUNTY: LANCASTER COUNTY, MIDDLESEX COUNTY, NORTHUMBERLAND COUNTY
COAST GUARD: DISTRICT 5, SECTOR HAMPTON ROADS
USACE: NORTH ATLANTIC DIVISION, NORFOLK DISTRICT

SHORELINE RESOURCES**ESI POLYGON HABITAT TYPES**

<i>ESI Rank</i>	<i>Habitat Classification</i>	<i>Area (Acres)</i>	<i>Area (Sq. Miles)</i>
10A	Salt and Brackish Water Marshes	811.73	1.27
10B	Freshwater Marshes	74.06	0.12
10C	Swamps	1,731.73	2.71
10D	Scrub and Shrub Wetlands	255.72	0.40
7	Exposed Tidal Flats	8.75	0.01

ESI SHORELINE HABITAT TYPES


<i>ESI Rank</i>	<i>Shoreline Habitat Classification</i>	<i>Length (Meters)</i>	<i>Length (Miles)</i>	<i>% of ESI Shoreline</i>
10A	Salt and Brackish Water Marshes	148,433.48	92.23	22%
10C	Swamps	7,374.06	4.58	1%
10D	Scrub and Shrub Wetlands	2,815.21	1.75	< 1%
9A	Sheltered Tidal Flats	3,178.02	1.97	< 1%
9B	Vegetated Low Banks	226,201.07	140.56	33%
8B	Sheltered, Solid Man-Made Structures	83,476.46	51.87	12%
8C	Sheltered Ripprap	61,287.72	38.08	9%
7	Exposed Tidal Flats	47,591.95	29.57	7%
6B	Ripprap	34,383.00	21.36	5%
4	Coarse Grained Sand Beaches	35,907.56	22.31	5%
3A	Fine to Medium Grained Sand Beaches	7,006.48	4.35	1%
3B	Scarps and Steep Slopes (Sand)	2,758.45	1.71	< 1%
1B	Exposed, Solid Man-Made Structures	15,692.62	9.75	2%

Total ESI Shoreline: 676,106.10 Total ESI Shoreline: 420.11
 Total Shoreline: 529,114.80 Total Shoreline: 328.78

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%.

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 66A Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	Mapping Qualifier	MDVA	S	F	Concentration	Monthly Presence
SAV	Submerged aquatic veg	High Ecological Value				Ephemeral	
SAV	Submerged aquatic veg	High Ecological Value				High	
SAV	Submerged aquatic veg	High Ecological Value				Low	

BIRDS

Map ID	Subelement	Species	Mapping Qualifier	MDVA	S	F	Concentration	Monthly Presence			
								J F M A M J J A S O N D	Nest	Mig.(S)	Mig.(F)
126	Wading	Great blue heron	Nesting				3 Nests	Feb-Jun	-	-	-
150	Wading	Great blue heron	Nesting				10 Nests	Feb-Jun	-	-	-
232	Waterfowl	Canada goose	Nesting				Present	Mar-Aug	-	-	-
	Waterfowl	Mallard	Nesting				Present	Mar-Aug	-	-	-
798	Raptor	Bald eagle	Nesting				1 Nest	Dec-Jun	-	-	-
862	Waterfowl	American black duck	Wintering				10S	-	-	-	-
	Waterfowl	Canada goose	Wintering				100S	-	-	-	-
	Waterfowl	Ruddy duck	Wintering				10S	-	-	-	-
	Waterfowl	Tundra swan	Wintering				10S	-	-	-	-
896	Waterfowl	American wigeon	Wintering				10S	-	-	-	-
	Waterfowl	Canvasback	Wintering				100S	-	-	-	-
	Waterfowl	Gadwall	Wintering				100S	-	-	-	-
	Waterfowl	Mallard	Wintering				100S	-	-	-	-
	Waterfowl	Mergansers	Wintering				100S	-	-	-	-
	Waterfowl	Redhead	Wintering				Present	-	-	-	-
1064	Waterfowl	Bufflehead	Wintering				100S	-	-	-	-
1072	Waterfowl	American black duck	Wintering				10S	-	-	-	-
1189	Diving	D. crested cormorant	Wintering				100S	-	-	Mar-May	Aug-Nov
	Diving	Grebes	Wintering				10S	-	-	-	-
	Waterfowl	Scoters	Wintering				10S	-	-	Mar-May	Aug-Nov
1190	Diving	D. crested cormorant	Wintering				100S	-	-	-	-
	Diving	Grebes	Wintering				10S	-	-	-	-
	Waterfowl	American wigeon	Wintering				10S	-	-	-	-
	Waterfowl	Bufflehead	Wintering				100S	-	-	-	-
	Waterfowl	Canvasback	Wintering				100S	-	-	-	-
	Waterfowl	Gadwall	Wintering				100S	-	-	-	-
	Waterfowl	Mallard	Wintering				100S	-	-	-	-
	Waterfowl	Mergansers	Wintering				100S	-	-	-	-
	Waterfowl	Redhead	Wintering				Present	-	-	-	-
	Waterfowl	Scoters	Wintering				10S	-	-	-	-

INVERTEBRATES

Map ID	Subelement	Species	Mapping Qualifier			MD/VA			Monthly Presence																
			Vulnerable Occurrence	E/T	T	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
1833	Insect	NE beach tiger beetle	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

BIRDS

Subelement	Species	Mapping Qualifier	MD/VA			Monthly Presence															
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
Diving	Common loon	Wintering	Present	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Common loon	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	D. crested cormorant	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Grebes	Grebes	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Grebes	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Red-throated loon	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Pelagic	Red-throated loon	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Northern gannet	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Waterfowl	American black duck	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Bufflehead	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Canada goose	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Canvasback	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Long-tailed duck	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Mallard	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Mergansers	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Ring-necked duck	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Ruddy duck	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Scaup	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Scoters	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Scoters	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Tundra swan	Wintering	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Subelement	Species	Mapping Qualifier	MD/VA			Monthly Presence															
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles
Diadromous	Alewife	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	American shad	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Atlantic sturgeon	General Distribution	-E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Blueback herring	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Hickory shad	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Shortnose sturgeon	General Distribution	E/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Striped bass	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Estuarine Nursery	Atl. sharpnose shark	Nursery Area	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Marine Pelagic	Sandbar shark	Nursery Area	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

REPTILES & AMPHIBIANS

		Monthly Presence																						
Subelement	Species	Mapping Qualifier		S	F	Concentration		J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles	Adults
Turtle	K. ridley sea turtle	General Distribution	E/E	E	Uncommon			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Dec	-
	K. ridley sea turtle	General Distribution	E/E	E	Common			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct	-
	Loggerhead sea turtle	General Distribution	T/T	T	Common			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Dec	Nov-Dec
	Loggerhead sea turtle	General Distribution	T/T	T	Abundant			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct	Apr-Oct

INVERTEBRATES

		Monthly Presence																						
Subelement	Species	Mapping Qualifier		S	F	Concentration		J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
Crab	Blue crab	Concentration Area				Highly Abundant-Summ													-	-	-	-	Jun-Oct	Jun-Oct

MARINE MAMMALS

		Monthly Presence																						
Subelement	Species	Mapping Qualifier		S	F	Concentration		J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping	Molt	
Manatee	West Indian manatee	General Distribution	-E	E	Rare														-	-	-	-	-	-
Whale	Humpback whale	General Distribution	E/E	E	Very Rare														-	-	-	-	-	-

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)

BIRDS

		Monthly Presence																						
Subelement	Species	Mapping Qualifier		S	F	Concentration		J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt	
Diving	Brown pelican																		-	-	-	-	-	-
	Brown pelican																		-	-	-	-	-	-
	D. crested cormorant																		-	-	-	-	-	-
Gull/Tern	Gulls																		-	-	-	-	-	-
	Terns																		-	-	-	-	-	-
Pelagic	Storm-petrels																		-	-	-	-	-	-
	Wood duck																		Apr-Jul	-	-	-	-	-
																			-	-	-	-	-	-

FISH

		Monthly Presence																						
Subelement	Species	Mapping Qualifier		S	F	Concentration		J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
Diadromous	Alewife																		-	-	-	-	-	-
	American eel																		Feb-Apr	Feb-Apr	Feb-Apr	Feb-Apr	Feb-Apr	Feb-Apr
	Blueback herring																		Mar-Jun	Mar-Jun	Mar-Jun	Mar-Jun	Mar-Jun	Mar-Jun
	Gizzard shad																		-	-	-	-	-	-
	Striped bass																		-	-	-	-	-	-
Estuarine Nursery	Atlantic croaker																		-	-	-	-	-	-
	Atlantic menhaden																		Apr-Jun	Apr-Jun	Apr-Jun	Apr-Jun	Apr-Jun	Apr-Jun
	Black drum																		-	-	-	-	-	-
	Black sea bass																		-	-	-	-	-	-
	Bluefish																		-	-	-	-	-	-
	Kingfishes																		-	-	-	-	-	-

FISH (continued)

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Pigfish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Red drum	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Sheepshead	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Spot	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Weakfish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Winter flounder	-	-	-	-	-	-	-	-	-	-	-	-	Dec-Mar
	Pipefish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Silversides	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Anchovies	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Striped burrfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Atlantic spadefish	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Clearnose skate	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Red hake	-	-	-	-	-	-	-	-	-	-	-	-	Sep-May*
	Silver hake	-	-	-	-	-	-	-	-	-	-	-	-	Oct-May
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Nov
	Cobia	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Lined seahorse	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Spanish mackerel	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Stout razor clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Squid	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Horseshoe crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Brown shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Mantis shrimp	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	White shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct

MARINE MAMMALS

Subelement	Species	Monthly Presence												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-
	Harbor porpoise	-	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES


ESI POLYGON HABITAT TYPES		Habitat Classification	Area (Acres)	Area (Sq. Miles)
ESI Rank				
10A		Salt and Brackish Water Marshes	273.39	0.43
10B		Freshwater Marshes	2.90	0.00
10C		Swamps	431.10	0.67
10D		Scrub and Shrub Wetlands	50.99	0.08

ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
ESI Rank					
10A		Salt and Brackish Water Marshes	35,230.88	21.89	17%
10C		Swamps	1,045.65	0.65	1%
10D		Scrub and Shrub Wetlands	660.41	0.41	< 1%
9A		Sheltered Tidal Flats	1,856.42	1.15	1%
9B		Vegetated Low Banks	79,424.99	49.35	38%
8B		Sheltered, Solid Man-Made Structures	19,361.20	12.03	9%
8C		Sheltered Riprap	25,195.13	15.66	12%
7		Exposed Tidal Flats	18,652.84	11.59	9%
6B		Riprap	8,709.23	5.41	4%
4		Coarse Grained Sand Beaches	13,594.38	8.45	7%
3A		Fine to Medium Grained Sand Beaches	25.02	0.02	< 1%
3B		Scars and Steep Slopes (Sand)	363.75	0.23	< 1%
1B		Exposed, Solid Man-Made Structures	2,530.82	1.57	1%
		Total ESI Shoreline:	206,650.73	Total ESI Shoreline:	128.41
		Total Shoreline:	160,556.85	Total Shoreline:	99.77

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 66B
Central Bay



SEE BACK OF MAP
for details about mapped species and
other species that occur in mapped area.
Data Published: September 2016

Map 66B Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	Mapping Qualifier	MD/VA	S	F	Concentration	Monthly Presence
SAV	Submerged aquatic veg	High Ecological Value	J F M A M J J A S O N D	J F M A M J J A S O N D	J F M A M J J A S O N D	Ephemeral	J F M A M J J A S O N D

BIRDS

Map ID	Subelement	Species	Mapping Qualifier	MD/VA	S	F	Concentration	Monthly Presence	
				J F M A M J J A S O N D	J F M A M J J A S O N D	Nest	Mig.(S)	Mig.(F)	Molt
143	Wading	Great blue heron	Nesting	7 Nests	Feb-Jun	-	-	-	-
180	Wading	Great blue heron	Nesting	12 Nests	Feb-Jun	-	-	-	-
232	Waterfowl	Canada goose	Nesting	Present	Mar-Aug	-	-	-	Jun-Jul
	Waterfowl	Mallard	Nesting	Present	Mar-Aug	-	-	-	Jul-Aug
798	Raptor	Bald eagle	Nesting	1 Nest	Dec-Jun	-	-	-	-
858	Waterfowl	Canada goose	Wintering	100S	-	-	-	-	-
	Waterfowl	Ruddy duck	Wintering	10S	-	-	-	-	-
	Waterfowl	Scaup	Wintering	10S	-	-	-	-	-
943	Waterfowl	American black duck	Wintering	10S	-	-	-	-	-
	Waterfowl	Bufflehead	Wintering	100S	-	-	-	-	-
	Waterfowl	Canada goose	Wintering	1,000S	-	-	-	-	-
	Waterfowl	Gadwall	Wintering	Present	Present	-	-	-	-
	Waterfowl	Mallard	Wintering	10S	-	-	-	-	-
	Waterfowl	Mergansers	Wintering	100S	-	-	-	-	-
	Waterfowl	Ruddy duck	Wintering	1,000S	-	-	-	-	-
	Waterfowl	Scaup	Wintering	100S	-	-	-	-	-
	Waterfowl	Tundra swan	Wintering	10S	-	-	-	-	-
986	Waterfowl	Bufflehead	Wintering	100S	-	-	-	-	-
	Waterfowl	Mallard	Wintering	10S	-	-	-	-	-
1127	Diving	Common loon	Wintering	Present	Present	-	-	-	-
	Diving	Grebes	Wintering	Present	Present	-	-	-	-
	Diving	Red-throated loon	Wintering	10S	-	-	-	-	-
	Pelagic	Northern gannet	Wintering	100S	-	-	-	-	-
	Waterfowl	American black duck	Wintering	10S	-	-	-	-	-
	Waterfowl	Bufflehead	Wintering	1,000S	-	-	-	-	-
	Waterfowl	Canada goose	Wintering	1,000S	-	-	-	-	-
	Waterfowl	Canvasback	Wintering	10S	-	-	-	-	-
	Waterfowl	Long-tailed duck	Wintering	10S	-	-	-	-	-
	Waterfowl	Mallard	Wintering	10S	-	-	-	-	-
	Waterfowl	Mergansers	Wintering	10S	-	-	-	-	-
	Waterfowl	Ring-necked duck	Wintering	10S	-	-	-	-	-
	Waterfowl	Ruddy duck	Wintering	100S	-	-	-	-	-

BIRDS (continued)

Map ID	Subelement	Species	MD/VA			Monthly Presence												Nest		Mig.(S)	Mig.(F)	Molt	
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest		Mig.(S)	Mig.(F)	Molt
Waterfowl	Waterfowl	Scaup	Wintering																-	-	-	-	-
Waterfowl	Waterfowl	Scoters	Wintering																-	-	-	-	-
Waterfowl	Waterfowl	Tundra swan	Wintering																-	-	-	-	-

FISH

Map ID	Subelement	Species	MD/VA			Monthly Presence												Nest		Mig.(S)	Mig.(F)	Molt	
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
1552	Diadromous	Alewife	Migration																-	-	-	Aug-Nov	Feb-Apr
	Diadromous	American shad	Migration																-	-	-	Aug-Nov	Feb-Jun
	Diadromous	Atlantic sturgeon	General Distribution	-E	E	-													-	-	-	Aug-Nov	Mar-Jun
	Diadromous	Blueback herring	Migration																-	-	-	Jun-Sep	Feb-May
	Diadromous	Hickory shad	General Distribution																-	-	-	Jan-Dec	Jan-Dec
	Diadromous	Shortnose sturgeon	General Distribution	E/E	E	-													-	-	-	Feb-Jun	Jan-Dec
	Diadromous	Striped bass	Migration																-	-	-	Jun-Sep	Jun-Sep
	Estuarine Nursery	Atl. sharpnose shark	Nursery Area																-	-	-	May-Sep	May-Sep
	Marine Pelagic	Sandbar shark	Nursery Area																-	-	-	Aug-Nov	Feb-Apr
1555	Diadromous	Alewife	Migration																-	-	-	Aug-Nov	Jan-Dec
	Diadromous	Atlantic sturgeon	General Distribution	-E	E	-													-	-	-	Aug-Nov	Mar-Jun
	Diadromous	Blueback herring	Migration																-	-	-	Jan-Dec	Jan-Dec
	Diadromous	Shortnose sturgeon	General Distribution	E/E	E	-													-	-	-	Mar-May	Jan-Dec
	Diadromous	Striped bass	Nursery Area																-	-	-	Jan-Dec	Jan-Dec
1556	Diadromous	Atlantic sturgeon	General Distribution	-E	E	-													-	-	-	Mar-May	Jan-Dec
	Diadromous	Shortnose sturgeon	General Distribution	E/E	E	-													-	-	-	Jan-Dec	Jan-Dec
	Diadromous	Striped bass	Nursery Area																-	-	-	Mar-May	Jan-Dec

REPTILES & AMPHIBIANS

Map ID	Subelement	Species	MD/VA			Monthly Presence												Nest		Hatch	Internest	Juveniles	Adults
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
1767	Turtle	Green sea turtle	General Distribution	T/T	T	Rare												-	-	-	May-Dec*	-	-
	Turtle	Green sea turtle	General Distribution	T/T	T	Common												-	-	-	Jul-Oct	-	-
1775	Turtle	K. ridley sea turtle	General Distribution	E/E	E	Common												-	-	-	Apr-Oct	-	-
	Turtle	K. ridley sea turtle	General Distribution	E/E	E	Uncommon												-	-	-	Nov-Dec	-	-
	Turtle	Loggerhead sea turtle	General Distribution	T/T	T	Common												-	-	-	Nov-Dec	Nov-Dec	-
	Turtle	Loggerhead sea turtle	General Distribution	T/T	T	Abundant												-	-	-	Apr-Oct	Apr-Oct	-

INVERTEBRATES

Map ID	Subelement	Species	MD/VA			Monthly Presence												Nest		Hatch	Internest	Juveniles	Adults
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
1788	Crab	Blue crab	Concentration Area			Highly Abundant-Summ												-	-	-	Jun-Oct	Jun-Oct	-
1792	Bivalve	Eastern elliptio	Vulnerable Occurrence															-	-	-	Jan-Dec	Jan-Dec	-
1813	Bivalve	Variable spike	Vulnerable Occurrence															-	-	-	Jan-Dec	Jan-Dec	-
	Bivalve	Eastern elliptio	Vulnerable Occurrence															-	-	-	Jan-Dec	Jan-Dec	-
	Bivalve	Eastern floater	Vulnerable Occurrence															-	-	-	Jan-Dec	Jan-Dec	-

MARINE MAMMALS

Map ID	Subelement	Species	Mapping Qualifier	MD/VA	S F	Concentration	J F M A M J J A S O N D	Mating	Calving	Pupping	Molt
1844	Manatee	West Indian manatee	General Distribution	-IE	E	Rare		-	-	-	-

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)

BIRDS

Subelement	Species	Monthly Presence											
		J	F	M	A	M	J	J	A	S	O	N	D
Diving	Brown pelican												
	D. crested cormorant												
Gull/Tern	Gulls												
	Terns												
Waterfowl	Wood duck												

FISH

Subelement	Species	Monthly Presence											
		J	F	M	A	M	J	J	A	S	O	N	D
Diacromous	Alewife												
	American eel												
	Blueback herring												
	Gizzard shad												
	Striped bass												
Estuarine Nursery	Atlantic croaker												
	Atlantic croaker												
	Atlantic herring												
	Atlantic menhaden												
	Black drum												
	Black sea bass												
	Bluefish												
	Kingfishes												
	Northern puffer												
	Northern puffer												
	Pigfish												
	Red drum												
	Sheepshead												
	Spot												
	Spotted seatrout												
	Summer flounder												
	Weakfish												
	White perch												
	Winter flounder												
Estuarine Resident	Pipefish												
	Silversides												
Fish	Anchovies												
	Striped burrfish												

FISH (continued)

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Marine Benthic	Atlantic spadefish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Clearmose skate	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Sep
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Red hake	-	-	-	-	-	-	-	-	-	-	-	-	Sep-May*
	Silver hake	-	-	-	-	-	-	-	-	-	-	-	-	Oct-May
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Marine Pelagic	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Cobia	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Lined seahorse	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Spanish mackerel	-	-	-	-	-	-	-	-	-	-	-	-	May-Sep

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Bivalve	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Stout razor clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Squid	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Cephalopod	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
Crab	Horseshoe crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Brown shrimp	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Shrimp	Mantis shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	White shrimp	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec

MARINE MAMMALS

Subelement	Species	Monthly Presence												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
Dolphin	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-
	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-
	Harbor porpoise	-	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES

ESI POLYGON HABITAT TYPES		Habitat Classification	Area (Acres)	Area (Sq. Miles)
ESI Rank				
10A		Salt and Brackish Water Marshes	118.64	0.19
10B		Freshwater Marshes	45.73	0.07
10C		Swamps	831.65	1.30
10D		Scrub and Shrub Wetlands	143.21	0.22


ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
ESI Rank					
10A		Salt and Brackish Water Marshes	17,448.46	10.84	15%
10C		Swamps	1,584.43	0.98	1%
10D		Scrub and Shrub Wetlands	248.57	0.15	< 1%
9B		Vegetated Low Banks	54,574.12	33.91	48%
8B		Sheltered, Solid Man-Made Structures	22,651.55	14.08	20%
8C		Riprap	16,499.28	10.25	14%
6B		Coarse Grained Sand Beaches	322.19	0.20	< 1%
4		Scarp and Steep Slopes (Sand)	541.29	0.34	< 1%
3B		Scarp and Steep Slopes (Sand)	474.18	0.29	< 1%

Total ESI Shoreline: 114,344.07 Total ESI Shoreline: 71.05
 Total Shoreline: 108,158.04 Total Shoreline: 67.21

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%.

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 66C Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	Mapping Qualifier						Concentration	Monthly Presence							
		S	F	MDVA	J	F	M	A	M	J	J	A	S	O	N	D
SAV	Submerged aquatic veg	High Ecological Value														
SAV	Submerged aquatic veg	High Ecological Value														
SAV	Submerged aquatic veg	High Ecological Value														

BIRDS

Map ID	Subelement	Species	Mapping Qualifier						Concentration	Monthly Presence							
			S	F	MDVA	J	F	M	A	M	J	J	A	S	O	N	D
179	Wading	Great blue heron	Nesting						26 Nests	Feb-Jun							
193	Wading	Great blue heron	Nesting						47 Nests	Feb-Jun							
232	Waterfowl	Canada goose	Nesting						Present	Mar-Aug							
	Waterfowl	Mallard	Nesting						Present	Mar-Aug							
710	Waterfowl	American black duck	Wintering						100S								
798	Raptor	Bald eagle	Nesting						1 Nest	Dec-Jun							
861	Waterfowl	American black duck	Wintering						Present								
	Waterfowl	Bufflehead	Wintering						100S								
	Waterfowl	Canada goose	Wintering						100S								
	Waterfowl	Canvasback	Wintering						100S								
	Waterfowl	Common goldeneye	Wintering						Present								
	Waterfowl	Tundra swan	Wintering						10S								
898	Waterfowl	American black duck	Wintering						10S								
	Waterfowl	Bufflehead	Wintering						100S								
	Waterfowl	Canvasback	Wintering						100S								
	Waterfowl	Gadwall	Wintering						100S								
	Waterfowl	Mallard	Wintering						1,000S								
	Waterfowl	Ruddy duck	Wintering						1,000S								
	Waterfowl	Tundra swan	Wintering						Present								
1072	Waterfowl	American black duck	Wintering						100S								
1122	Diving	Red-throated loon	Wintering						10S								
	Waterfowl	Bufflehead	Wintering						1,000S								
	Waterfowl	Canvasback	Wintering						10S								
	Waterfowl	Long-tailed duck	Wintering						10S								
	Waterfowl	Ring-necked duck	Wintering						10S								
	Waterfowl	Scoters	Wintering						1,000S								
	Waterfowl	Tundra swan	Wintering						100S								
1123	Diving	Red-throated loon	Wintering						10S								
	Waterfowl	American black duck	Wintering						10S								
	Waterfowl	Bufflehead	Wintering						1,000S								

BIRDS (continued)

<i>Map ID</i>	<i>Subelement</i>	<i>Species</i>	<i>Mapping Qualifier</i>	<i>MD/VA</i>	<i>S</i>	<i>F</i>	<i>Concentration</i>	<i>Monthly Presence</i>											
				J	F	M	A	M	J	J	A	S	O	N	D	<i>Nest</i>	<i>Mig.(S)</i>	<i>Mig.(F)</i>	<i>Molt</i>
	Waterfowl	Canvasback	Wintering				10S									-	-	-	-
	Waterfowl	Long-tailed duck	Wintering				10S									-	-	-	-
	Waterfowl	Ring-necked duck	Wintering				1,000S									-	-	-	-
	Waterfowl	Scoters	Wintering				100S									-	-	-	-
	Waterfowl	Tundra swan	Wintering				100S									-	-	-	-

INVERTEBRATES

<i>Map ID</i>	<i>Subelement</i>	<i>Species</i>	<i>Mapping Qualifier</i>	<i>MD/VA</i>	<i>S</i>	<i>F</i>	<i>Concentration</i>	<i>Monthly Presence</i>											
			Vulnerable Occurrence	E/T	T	-		J	F	M	A	M	J	J	A	S	O	D	<i>Spawn</i>
1833	Insect	NE beach tiger beetle																	

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

BIRDS

<i>Subelement</i>	<i>Species</i>	<i>Mapping Qualifier</i>	<i>MD/VA</i>	<i>S</i>	<i>F</i>	<i>Concentration</i>	<i>Monthly Presence</i>											
				J	F	M	A	M	J	J	A	S	O	N	<i>Nest</i>	<i>Mig.(S)</i>	<i>Mig.(F)</i>	<i>Molt</i>
Diving	Common loon	Wintering				100S									-	-	-	-
	Common loon	Wintering				Present									-	-	-	-
	D. crested cormorant	Wintering				1,000S									-	Mar-May	Aug-Nov	-
	D. crested cormorant	Wintering				100S									-	Mar-May	Aug-Nov	-
	Grebes	Wintering				100S									-	-	-	-
	Grebes	Wintering				10S									-	-	-	-
	Red-throated loon	Wintering				Present									-	-	-	-
	Red-throated loon	Wintering				100S									-	-	-	-
Pelagic	Northern gannet	Wintering				1,000S									-	-	-	-
	Canada goose	Wintering				100S									-	-	-	-
	Long-tailed duck	Wintering				100S									-	-	-	-
	Mallard	Wintering				10S									-	-	-	-
	Mergansers	Wintering				10S									-	-	-	-
	Ruddy duck	Wintering				100S									-	-	-	-
	Scaup	Wintering				100S									-	-	-	-
	Scoters	Wintering				10,000S									-	-	-	-
	Scoters	Wintering													-	-	-	-

FISH

<i>Subelement</i>	<i>Species</i>	<i>Mapping Qualifier</i>	<i>MD/VA</i>	<i>S</i>	<i>F</i>	<i>Concentration</i>	<i>Monthly Presence</i>												
				J	F	M	A	M	J	J	A	S	O	N	<i>Spawn</i>	<i>Eggs</i>	<i>Larvae</i>	<i>Juveniles</i>	<i>Adults</i>
Diadromous	Alewife	Migration													-	-	Aug-Nov	Feb-Apr	
	American shad	Migration													-	-	Aug-Nov	Feb-Jun	
	Atlantic sturgeon	General Distribution	-E	E											-	-	Jan-Dec	-	
	Atlantic sturgeon	Migration	-E	E											-	-	Jun-Nov	Mar-Jun	
	Blueback herring	Migration													-	-	Aug-Nov	Feb-May	
	Hickory shad	Migration													-	-	Jun-Sep	Feb-May	

FISH (continued)

Subelement	Species	MD/VA						Monthly Presence						Adults						
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles
Shortnose sturgeon	General Distribution	E/E	E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Striped bass	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Jun
Striped bass	Nursery Area	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
Estuarine Nursery	Atl. sharpnose shark	Nursery Area	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Sep
Marine Pelagic	Sandbar shark	Nursery Area	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Sep

REPTILES & AMPHIBIANS

Subelement	Species	MD/VA						Monthly Presence						Adults						
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles
Turtle	Green sea turtle	T/T	T	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Oct
	Green sea turtle	T/T	T	Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Dec*
	K. ridley sea turtle	E/E	E	Uncommon	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Dec
	K. ridley sea turtle	E/E	E	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct
	Loggerhead sea turtle	T/T	T	Common	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Dec
	Loggerhead sea turtle	T/T	T	Abundant	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct

INVERTEBRATES

Subelement	Species	MD/VA						Monthly Presence						Adults						
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles
Crab	Blue crab	Concentration Area						Highly Abundant-Summ						-	-	-	-	-	-	Jun-Oct

MARINE MAMMALS

Subelement	Species	MD/VA						Monthly Presence						Adults						
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping	Molt
Manatee	West Indian manatee	-/E	E	Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Whale	Humpback whale	E/E	E	Very Rare	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)

BIRDS

Subelement	Species	MD/VA						Monthly Presence						Adults					
		J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt		
Diving	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Gull/Tern	Gulls	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Terns	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Pelagic	Storm-petrels	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
 Diadromous	Alewife	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Apr
	American eel	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct
	Blueback herring	-	-	-	-	-	-	-	-	-	-	-	-	Jan-May
	Gizzard shad	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Jun
	Striped bass	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
 Estuarine Nursery	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Atlantic herring	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Jun
	Atlantic menhaden	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Black drum	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Black sea bass	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Bluefish	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Kingfishes	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Oct
	Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Oct
	Pigfish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Red drum	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Sheepshead	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Spot	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Weakfish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	White perch	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Winter flounder	-	-	-	-	-	-	-	-	-	-	-	-	Dec-Mar
 Estuarine Resident	Pipefish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Silversides	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
 Fish	Anchoovies	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Striped burrfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
 Marine Benthic	Atlantic spadefish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Cleinnose skate	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Sep
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Red hake	-	-	-	-	-	-	-	-	-	-	-	-	Sep-May*
	Silver hake	-	-	-	-	-	-	-	-	-	-	-	-	Oct-May
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Nov
 Marine Pelagic	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Cobia	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Lined seahorse	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Spanish mackerel	-	-	-	-	-	-	-	-	-	-	-	-	May-Sep

REPTILES & AMPHIBIANS

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Turtle	Diamondback terrapin											May-Aug	-	Apr-Oct

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Bivalve	Eastern oyster											Jun-Sep	Jun-Sep	Jan-Dec
	Hard clam											Jun-Sep	Jun-Sep	Jan-Dec
	Softshell clam											May-Dec*	May-Dec*	Jan-Dec
	Stout razor clam											Jun-Nov	-	Jan-Dec
	Squid											-	Apr-Dec	Apr-Dec
Cephalopod	Blue crab											-	-	Jan-Dec
	Horseshoe crab											-	-	Jan-Dec
	Brown shrimp											-	-	May-Oct
Shrimp	Mantis shrimp											-	-	Jan-Dec
	White shrimp											-	-	May-Oct

MARINE MAMMALS

Subelement	Species	Monthly Presence												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
Dolphin	Bottlenose dolphin											May-Aug	May-Aug	-
	Harbor porpoise											-	-	-
Pinniped	Harbor seal											-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES

ESI POLYGON HABITAT TYPES		Habitat Classification	Area (Acres)	Area (Sq. Miles)
ESI Rank				
10A		Salt and Brackish Water Marshes	246.40	0.39
10B		Freshwater Marshes	6.20	0.01
10C		Swamps	111.83	0.17
10D		Scrub and Shrub Wetlands	25.41	0.04
7		Exposed Tidal Flats	6.00	0.01


ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
ESI Rank					
10A		Salt and Brackish Water Marshes	46,846.12	29.11	26%
10C		Swamps	528.10	0.33	< 1%
10D		Scrub and Shrub Wetlands	934.60	0.58	1%
9A		Sheltered Tidal Flats	215.32	0.13	< 1%
9B		Vegetated Low Banks	41,655.91	25.88	23%
8B		Sheltered, Solid Man-Made Structures	21,638.66	13.45	12%
8C		Sheltered Riprap	7,427.97	4.62	4%
7		Exposed Tidal Flats	23,850.88	14.82	13%
6B		Riprap	15,351.52	9.54	9%
4		Coarse Grained Sand Beaches	14,007.55	8.70	8%
3A		Fine to Medium Grained Sand Beaches	1,485.24	0.92	1%
1B		Exposed, Solid Man-Made Structures	5,385.63	3.35	3%

Total ESI Shoreline: 179,327.51 Total ESI Shoreline: 111.43
 Total Shoreline: 126,512.86 Total Shoreline: 78.61

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 66D Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	Mapping Qualifier	MDVA	S	F	Concentration	Monthly Presence
SAV	Submerged aquatic veg	High Ecological Value				Ephemeral	
SAV	Submerged aquatic veg	High Ecological Value				High	
SAV	Submerged aquatic veg	High Ecological Value				Low	

BIRDS

Map ID	Subelement	Species	Mapping Qualifier	MDVA	S	F	Concentration	Monthly Presence												
								J	F	M	A	M	J	J	A	S	O	N	D	Nest
52	Diving	D. crested cormorant	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Mar-May
	Diving	Grebes	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	Aug-Nov
108	Wading	Great blue heron	Nesting					-	-	-	-	-	-	-	-	-	-	-	-	-
153	Wading	Great blue heron	Nesting					-	-	-	-	-	-	-	-	-	-	-	-	-
169	Wading	Great blue heron	Nesting					-	-	-	-	-	-	-	-	-	-	-	-	-
232	Watertowl	Canada goose	Nesting					-	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Mallard	Nesting					-	-	-	-	-	-	-	-	-	-	-	-	-
681	Watertowl	Brant	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Bufflehead	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Canada goose	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Canvasback	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Mergansers	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Scaup	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Tundra swan	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	-
705	Raptor	Peregrine falcon	Nesting					-	-	-	-	-	-	-	-	-	-	-	-	-
798	Raptor	Bald eagle	Nesting					-	-	-	-	-	-	-	-	-	-	-	-	-
853	Watertowl	Canada goose	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Mallard	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Ruddy duck	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Scaup	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	-
1237	Diving	Common loon	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	-
	Diving	Red-throated loon	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	-
	Pelagic	Northern gannet	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Long-tailed duck	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Scoters	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	-
1241	Diving	Common loon	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	-
	Pelagic	Northern gannet	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Bufflehead	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Canvasback	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	-
	Watertowl	Mallard	Wintering					-	-	-	-	-	-	-	-	-	-	-	-	-

BIRDS (continued)

Map ID	Subelement	Species	MD\VA			Monthly Presence																
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
	Watertowl	Ring-necked duck	Wintering			10S													-	-	-	-
	Watertowl	Ruddy duck	Wintering			100S													-	-	-	-
	Watertowl	Scoters	Wintering			1,000S													-	-	-	-
	Watertowl	Tundra swan	Wintering			100S													-	-	-	-
1369	Diving	D. crested cormorant	Wintering			1,000S													Mar-May	Aug-Nov	-	-
	Watertowl	American black duck	Wintering			Present													-	-	-	-
	Watertowl	Canada goose	Wintering			100S													-	-	-	-
	Watertowl	Common goldeneye	Wintering			Present													-	-	-	-
	Watertowl	Mallard	Wintering			10S													-	-	-	-
	Watertowl	Ruddy duck	Wintering			100S													-	-	-	-

FISH

Map ID	Subelement	Species	MD\VA			Monthly Presence																	
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
1606	Diadromous	Striped bass	Migration			-													-	-	-	-	Feb-Jun
	Estuarine Nursery	Atl. sharpnose shark	Nursery Area			-													-	-	-	-	Jun-Sep
	Marine Pelagic	Sandbar shark	Nursery Area			-													-	-	-	-	May-Sep

REPTILES & AMPHIBIANS

Map ID	Subelement	Species	MD\VA			Monthly Presence																
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles
1762	Turtle	Diamondback terrapin	Concentration Area			High													-	-	Apr-Oct	Apr-Oct

INVERTEBRATES

Map ID	Subelement	Species	MD\VA			Monthly Presence																	
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
1788	Crab	Blue crab	Concentration Area			Highly Abundant-Summ												-	-	-	-	Jun-Oct	
1833	Insect	NE beach tiger beetle	Vulnerable Occurrence	E/T	T	-													Jun-Aug	Jun-Aug	Jan-Dec	-	Jun-Aug

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

Map ID	Subelement	Species	MD\VA			Monthly Presence																
			Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
1788	Diving	Common loon	Wintering			10S												-	Mar-May	Aug-Nov	-	-
		D. crested cormorant	Wintering			100S												-	-	-	-	-
		Grebes	Wintering			Present												-	-	-	-	-
		Red-throated loon	Wintering			10S												-	-	-	-	-
		American black duck	Wintering			100S												-	-	-	-	-
		Bufflehead	Wintering			1,000S												-	-	-	-	-
		Canada goose	Wintering			100S												-	-	-	-	-
		Canvasback	Wintering			Present												-	-	-	-	-
		Gadwall	Wintering			10S												-	-	-	-	-
		Long-tailed duck	Wintering			Present												-	-	-	-	-

BIRDS (continued)

Map ID	Subelement	Species	MD/VA			Monthly Presence															
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
		Mallard			Wintering													-	-	-	-
		Mergansers			Wintering													-	-	-	-
		Ruddy duck			Wintering													-	-	-	-
		Scaup			Wintering													-	-	-	-
		Scooter's			Wintering													-	-	-	-
		Tundra swan			Wintering													-	-	-	-

FISH

Subelement	Species	MD/VA			Monthly Presence																
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
	Alewife			Migration													-	-	-	Aug-Nov	Feb-Apr
	American shad			Migration													-	-	-	Aug-Nov	Feb-Jun
	Atlantic sturgeon			General Distribution	-E	E											-	-	-	-	Jan-Dec
	Atlantic sturgeon			Migration	-E	E											-	-	-	-	Jun-Nov
	Blueback herring			Migration	-												-	-	-	Aug-Nov	Mar-Jun
	Hickory shad			Migration	-												-	-	-	Jun-Sep	Feb-May
	Shortnose sturgeon			General Distribution	E/E	E											-	-	-	Jan-Dec	Jan-Dec
	Striped bass			Nursery Area	-												-	-	Mar-May	Jan-Dec	-

REPTILES & AMPHIBIANS

Subelement	Species	MD/VA			Monthly Presence																
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles	Adults
	Green sea turtle	General Distribution	T/T	T	Rare											-	-	-	-	May-Dec*	-
	Green sea turtle	General Distribution	T/T	T	Common											-	-	-	-	Jul-Oct	-
	K. ridley sea turtle	General Distribution	E/E	E	Uncommon											-	-	-	-	Nov-Dec	-
	K. ridley sea turtle	General Distribution	E/E	E	Common											-	-	-	-	Apr-Oct	-
	Loggerhead sea turtle	General Distribution	T/T	T	Common											-	-	-	-	Nov-Dec	Nov-Dec
	Loggerhead sea turtle	General Distribution	T/T	T	Abundant											-	-	-	-	Apr-Oct	Apr-Oct

INVERTEBRATES

Subelement	Species	MD/VA			Monthly Presence																
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
	Concentration Area	General Distribution	-/E	E	Highly Abundant											-	-	-	-	Jun-Mar*	Jun-Mar*
	Concentration Area	Concentration Area	Concentration Area	Concentration Area	Highly Abundant-Wint											-	-	-	-	Dec-Mar	Dec-Mar

MARINE MAMMALS

Subelement	Species	MD/VA			Monthly Presence															
		S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping	Molt
	West Indian manatee	General Distribution	-/E	E	Rare											-	-	-	-	-

BIRDS

Sublement	Species	Monthly Presence												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
Diving	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-	-
	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-	-
	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-	-
Gull/Tern	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-	-
	Gulls	-	-	-	-	-	-	-	-	-	-	-	-	-
	Terns	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Sublement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Diadromous	Alewife	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Apr
	American eel	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct
	Blueback herring	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Apr
	Gizzard shad	-	-	-	-	-	-	-	-	-	-	-	-	Jan-May
	Striped bass	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Oct
Estuarine Nursery	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Atlantic herring	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Atlantic menhaden	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Jun
	Black drum	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Black sea bass	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Bluefish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Kingfishes	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Oct
	Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Oct
	Pigfish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Red drum	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Sheepshead	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Spot	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Apr
	Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Weakfish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Aug
	White perch	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Winter flounder	-	-	-	-	-	-	-	-	-	-	-	-	Dec-Mar
Estuarine Resident	Pipefish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Silversides	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Anchovies	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Striped burrfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
Marine Benthic	Atlantic spadefish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Clearamose skate	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec

FISH (continued)

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Sep
	Red hake	-	-	-	-	-	-	-	-	-	-	-	-	Sep-May*
	Silver hake	-	-	-	-	-	-	-	-	-	-	-	-	Oct-May
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Marine Pelagic	Cobia	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Lined seahorse	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Spanish mackerel	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		-	-	-	-	-	-	-	-	-	-	-	-	May-Sep

REPTILES & AMPHIBIANS

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Diamondback terrapin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Bivalve	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Stout razor clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Cephalopod	Squid	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Crab	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Horseshoe crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Shrimp	Brown shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Mantis shrimp	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	White shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct

MARINE MAMMALS

Subelement	Species	Monthly Presence												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
Dolphin	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-
	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-
	Harbor porpoise	-	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES


ESI POLYGON HABITAT TYPES		Habitat Classification	Area (Acres)	Area (Sq. Miles)
ESI Rank				
10A		Salt and Brackish Water Marshes	173.29	0.27
10B		Freshwater Marshes	19.22	0.03
10C		Swamps	357.16	0.56
10D		Scrub and Shrub Wetlands	36.12	0.06
7		Exposed Tidal Flats	2.75	0.00

ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
ESI Rank					
10A		Salt and Brackish Water Marshes	48,908.01	30.39	28%
10C		Swamps	4,215.88	2.62	2%
10D		Scrub and Shrub Wetlands	971.63	0.60	1%
9A		Sheltered Tidal Flats	1,106.28	0.69	1%
9B		Vegetated Low Banks	50,546.06	31.41	29%
8B		Sheltered, Solid Man-Made Structures	19,825.05	12.32	11%
8C		Sheltered Riprap	12,165.34	7.56	7%
7		Exposed Tidal Flats	5,088.23	3.16	3%
6B		Riprap	10,000.06	6.21	6%
4		Coarse Grained Sand Beaches	7,764.34	4.82	4%
3A		Fine to Medium Grained Sand Beaches	5,496.21	3.42	3%
3B		Scarps and Steep Slopes (Sand)	1,920.52	1.19	1%
1B		Exposed, Solid Man-Made Structures	7,776.18	4.83	4%
		Total ESI Shoreline:	175,783.79	Total ESI Shoreline:	109.23
		Total Shoreline:	133,887.04	Total Shoreline:	83.19

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 67
Central Bay



Map 67
Central Bay

HUMAN USE RESOURCES

DISPLAYED ON MAP (POINTS)

Map ID	Type	Name	Contact	Phone
1	AIRPORT	ARROWHEAD POINT		
2	AIRPORT	ROSEGILL FARM AIRSTRIP		
3	AQUACULTURE	CRAB SHED SHORES AND RUARK SEAFOOD RF#98-0545	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
4	AQUACULTURE	OYSTER COMMERCIAL MARTIN RF#98-1768	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
5	AQUACULTURE	OYSTER NON-COMM ANDERSON, DAREL RF#2001-1692	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
6	AQUACULTURE	OYSTER NON-COMM BEARD RF#98-0891	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
7	AQUACULTURE	OYSTER NON-COMM BIGFORD RF#002176	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
8	AQUACULTURE	OYSTER NON-COMM CATHEY RF#99-0543	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
9	AQUACULTURE	OYSTER NON-COMM CHES BAY GOV'S SCHOOL RF#99-2001	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
10	AQUACULTURE	OYSTER NON-COMM CHES BAY GOVERNOR'S RF#20020210	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
11	AQUACULTURE	OYSTER NON-COMM CHRISTCHURCH SCHOOL RF#99-1181	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
12	AQUACULTURE	OYSTER NON-COMM DAVIS, JERRY B. RF#98-1147	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
13	AQUACULTURE	OYSTER NON-COMM FERRITTER, NICHOLAS RF#00-0756	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
14	AQUACULTURE	OYSTER NON-COMM FOLKES RF#99-0605	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
15	AQUACULTURE	OYSTER NON-COMM GREGORY, THOMAS RF#98-1694	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
16	AQUACULTURE	OYSTER NON-COMM HEARNS, CHARLES B. RF#98-0832	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
17	AQUACULTURE	OYSTER NON-COMM HENDERSON, GEORGE RF#98-0907	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
18	AQUACULTURE	OYSTER NON-COMM HOFFMAN, GREG RF#00-0638	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
19	AQUACULTURE	OYSTER NON-COMM HOLTON RF#99-1765	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
20	AQUACULTURE	OYSTER NON-COMM HOUCK RF#99-2031	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
21	AQUACULTURE	OYSTER NON-COMM JACOBI RF#99-1896	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
22	AQUACULTURE	OYSTER NON-COMM JOHNSON, FREDERICK RF#01-1965	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
23	AQUACULTURE	OYSTER NON-COMM MATTHEWS, ED RF#99-1968	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
24	AQUACULTURE	OYSTER NON-COMM PFEIFFER, W. A. RF#99-1202	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
25	AQUACULTURE	OYSTER NON-COMM RACKEL, E. J. RF#98-0774	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
26	AQUACULTURE	OYSTER NON-COMM RICHARDSON RF#991247	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
27	AQUACULTURE	OYSTER NON-COMM ROBERTSON RF#99-1633	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
28	AQUACULTURE	OYSTER NON-COMM SAVINO RF#001977	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
29	AQUACULTURE	OYSTER NON-COMM SANDY RF#98-1876	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
30	AQUACULTURE	OYSTER NON-COMM SIMON, R. W. RF#98-1923	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
31	AQUACULTURE	OYSTER NON-COMM THOMPSON, GEORGE E. RF#99-1945	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
32	AQUACULTURE	OYSTER NON-COMM VETTE, JAMES RF#01-2014	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
33	AQUACULTURE	OYSTER NON-COMM WEEKLY, ROBERT RF#98-1023	CHIP NEIKIRK, VIRGINIA MARINE RESOURCES COMM.	757-247-2254
34	BOAT RAMP	GREENVALE CREEK	LENE PENNINGTON, VDGIF	804-367-0058
35	BOAT RAMP	MILL STONE	LENE PENNINGTON, VDGIF	804-367-0058
36	BOAT RAMP	SALUDA	LENE PENNINGTON, VDGIF	804-367-0058
37	REPEATED MEASUREMENT SITE	W/Q - STATION LE3.2	CHESAPEAKE BAY PROGRAM	800-968-7229
38	REPEATED MEASUREMENT SITE	W/Q - STATION LE3.2N	CHESAPEAKE BAY PROGRAM	800-968-7229
39	REPEATED MEASUREMENT SITE	W/Q - STATION LE3.2S	CHESAPEAKE BAY PROGRAM	800-968-7229
40	REPEATED MEASUREMENT SITE	W/Q - STATION LE3.3	CHESAPEAKE BAY PROGRAM	800-968-7229


DISPLAYED ON MAP (POINTS) (continued)

Map ID	Type	Name	Contact	Phone
41	REPEATED MEASUREMENT SITE	WQ - STATION LE3.3A	CHESAPEAKE BAY PROGRAM	800-968-7229
Map ID	Type	Name	Contact	Phone
42	NATURE CONSERVANCY	DRAGON RUN	THE NATURE CONSERVANCY	703-841-5300
JURISDICTIONS				
COUNTY:	Gloucester County, Lancaster County, Middlesex County	FEMA:	REGION I	
COAST GUARD:	District 5, Sector Hampton Roads	EPA:	REGION 3	
USACE:	North Atlantic Division, Norfolk District			
SHORELINE RESOURCES				
ESI POLYGON HABITAT TYPES				
ESI Rank	Habitat Classification	Area (Acres)	Area (Sq. Miles)	% of ESI Shoreline
10A	Salt and Brackish Water Marshes	1,117.64	1.75	30%
10B	Freshwater Marshes	372.15	0.58	3%
10C	Swamps	2,542.70	3.97	3%
10D	Scrub and Shrub Wetlands	666.72	1.04	< 1%
9A	Sheltered Tidal Flats	5.81	0.01	1%
7	Exposed Tidal Flats	12.57	0.02	34%
ESI SHORELINE HABITAT TYPES				
ESI Rank	Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
10A	Salt and Brackish Water Marshes	155,565.83	96.66	34%
10B	Freshwater Marshes	14,729.51	9.15	5%
10C	Swamps	16,796.31	10.44	2%
10D	Scrub and Shrub Wetlands	2,076.76	1.29	1%
9A	Sheltered Tidal Flats	3,380.97	2.10	1%
9B	Vegetated Low Banks	177,510.78	110.30	9%
8B	Sheltered, Solid Man-Made Structures	47,058.68	29.24	5%
8C	Sheltered Riprap	22,251.64	13.83	< 1%
7	Exposed Tidal Flats	3,851.76	2.39	2%
6B	Riprap	27,264.58	16.94	2%
5	Mixed Sand and Gravel Beaches	25.60	0.02	2%
4	Coarse Grained Sand Beaches	27,264.29	16.94	5%
3B	Scars and Steep Slopes (Sand)	9,456.83	5.88	2%
1B	Exposed, Solid Man-Made Structures	13,019.88	8.09	3%
Total ESI Shoreline:	520,253.42	Total ESI Shoreline:	323.27	
Total Shoreline:	401,300.35	Total Shoreline:	249.36	

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%.

All underlying GIS data can be obtained from response.noaa.gov





Map 67A Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	Mapping Qualifier	MD/VA												Monthly Presence											
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt					
SAV	Submerged aquatic veg	High Ecological Value	-	-	Ephemeral	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
SAV	Submerged aquatic veg	High Ecological Value	-	-	High	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
SAV	Submerged aquatic veg	High Ecological Value	-	-	Low	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

BIRDS

Map ID	Subelement	Species	Mapping Qualifier	MD/VA												Monthly Presence											
				S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt					
108	Wading	Great blue heron	Nesting	-	-	6 Nests	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
195	Wading	Great blue heron	Nesting	-	-	48 Nests	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
232	Watertowl	Canada goose	Nesting	-	-	Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
798	Raptor	Mallard	Nesting	-	-	Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
944	Watertowl	Bald eagle	Nesting	-	-	1 Nest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
		Bufflehead	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
		Canada goose	Wintering	-	-	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
		Gadwall	Wintering	-	-	Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
		Mallard	Wintering	-	-	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
		Mergansers	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
		Scaup	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
		Tundra swan	Wintering	-	-	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

INVERTEBRATES

Map ID	Subelement	Species	Mapping Qualifier	MD/VA												Monthly Presence											
				S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt					
1789	Crab	Blue crab	Concentration Area	-	-	Highly Abundant	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

BIRDS

Subelement	Species	Mapping Qualifier	MD/VA												Monthly Presence											
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt					
Diving	Common loon	Wintering	-	-	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	D. crested cormorant	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Grebes	Wintering	-	-	Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Red-throated loon	Wintering	-	-	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	American black duck	Wintering	-	-	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Brant	Wintering	-	-	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Bufflehead	Wintering	-	-	1,000S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

BIRDS (continued)

Subelement	Species	MD\VA		Monthly Presence																			
		S	F	Mapping Qualifier		Concentration		J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
	Canada goose	Wintering				10,000S														-	-	-	-
	Canvasback	Wintering				1,000S														-	-	-	-
	Long-tailed duck	Wintering				10S														-	-	-	-
	Mallard	Wintering				100S														-	-	-	-
	Mergansers	Wintering				1,000S														-	-	-	-
	Ruddy duck	Wintering				1,000S														-	-	-	-
	Scaup	Wintering				1,000S														-	-	-	-
	Scoters	Wintering				10S														-	-	-	-
	Tundra swan	Wintering				100S														-	-	-	-

FISH

Subelement	Species	MD\VA		Monthly Presence																				
		S	F	Mapping Qualifier		Concentration		J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
	Alewife	Migration				-														-	-	-	Aug-Nov	Feb-Apr
	American shad	Migration				General Distribution	-E	E	-											-	-	-	Aug-Nov	Feb-Jun
	Atlantic sturgeon	Migration				General Distribution	-E	E	-											-	-	-	Jan-Dec	Jan-Dec
	Atlantic sturgeon	Migration				Hickory shad	-E	E	-											-	-	-	Jun-Nov	Jun-Nov
	Blueback herring	Migration				Shortnose sturgeon	-													-	-	-	Aug-Nov	Mar-Jun
	Hickory shad	Migration				Shortnose sturgeon	-													-	-	-	Jun-Sep	Feb-May
	Striped bass	Nursery Area				Striped bass	E/E	E	-											-	-	-	Jan-Dec	Jan-Dec

REPTILES & AMPHIBIANS

Subelement	Species	MD\VA		Monthly Presence																				
		S	F	Mapping Qualifier		Concentration		J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles	Adults
	Green sea turtle	General Distribution	T/T	T	Rare															-	-	-	May-Dec*	-
	Green sea turtle	General Distribution	T/T	T	Common															-	-	-	Jul-Oct	-

INVERTEBRATES

Subelement	Species	MD\VA		Monthly Presence																				
		S	F	Mapping Qualifier		Concentration		J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
	Blue crab	Concentration Area	Highly Abundant-Summ																	-	-	-	Jun-Oct	Jun-Oct

MARINE MAMMALS

Subelement	Species	MD\VA		Monthly Presence																			
		S	F	Mapping Qualifier		Concentration		J	F	M	A	M	J	J	A	S	O	N	D	Mating	Calving	Pupping	Molt
	Crab	General Distribution	-E	E	Rare															-	-	-	-
	Manatee	West Indian manatee																		-	-	-	-

BIRDS

Sublement	Species	<i>Monthly Presence</i>												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
Diving	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-	-
Gull/Tern	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-	-
Gulls	Gulls	-	-	-	-	-	-	-	-	-	-	-	-	-
Terns	Terns	-	-	-	-	-	-	-	-	-	-	-	-	-
Waterfowl	Wood duck	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Sublement	Species	<i>Monthly Presence</i>												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Diadromous	Alewife	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Apr
	American eel	-	-	-	-	-	-	-	-	-	-	-	-	Jan-May
	Blueback herring	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Jun
	Striped bass	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Estuarine Nursery	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Atlantic herring	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Jun
	Atlantic menhaden	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Black drum	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Black sea bass	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Bluefish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Kingfishes	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Oct
	Pigfish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Red drum	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Sheepshead	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Spot	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Apr
	Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Weakfish	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	White perch	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Estuarine Resident	Pipefish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Silversides	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Silversides	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Anchoovies	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Striped burrfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
Freshwater	Blue catfish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Marine Benthic	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Sep
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Marine Pelagic	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Lined seahorse	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Spanish mackerel	-	-	-	-	-	-	-	-	-	-	-	-	May-Sep

INVERTEBRATES

Subelement	Species	Monthly Presence											
		J	F	M	A	M	J	J	A	S	O	N	D
Bivalve	Eastern oyster												
	Softshell clam												
	Stout razor clam												
Cephalopod	Squid												
Crab	Blue crab												
Shrimp	Brown shrimp												
	Mantis shrimp												
	White shrimp												

MARINE MAMMALS

Subelement	Species	Monthly Presence											
		J	F	M	A	M	J	J	A	S	O	N	D
Dolphin	Bottlenose dolphin												

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov


ESI POLYGON HABITAT TYPES		Habitat Classification	Area (Acres)	Area (Sq. Miles)
ESI Rank				
10A		Salt and Brackish Water Marshes	180.55	0.28
10B		Freshwater Marshes	19.91	0.03
10C		Swamps	450.50	0.70
10D		Scrub and Shrub Wetlands	298.27	0.47

ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
ESI Rank					
10A		Salt and Brackish Water Marshes	27,441.37	17.05	15%
10C		Swamps	226.50	0.14	< 1%
10D		Scrub and Shrub Wetlands	537.56	0.33	< 1%
9B		Vegetated Low Banks	90,991.03	56.54	49%
8B		Sheltered, Solid Man-Made Structures	20,164.02	12.53	11%
8C		Sheltered Riprap	15,998.83	9.94	9%
6B		Riprap	10,763.13	6.69	6%
4		Coarse Grained Sand Beaches	8,669.02	5.39	5%
3B		Scars and Steep Slopes (Sand)	6,141.24	3.82	3%
1B		Exposed, Solid Man-Made Structures	5,137.87	3.19	3%
Total ESI Shoreline:		186,070.56	Total ESI Shoreline:	115.62	
Total Shoreline:		168,368.15	Total Shoreline:	104.62	

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





SEE BACK OF MAP
for details about mapped species and
other species that occur in mapped area.
Data Published: September 2016

Map 67B Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	Mapping Qualifier	MDVA												Monthly Presence														
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	J	F	M	A	M	J	J	A	S	O	N	D
SAV	Submerged aquatic veg	High Ecological Value	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
SAV	Submerged aquatic veg	High Ecological Value	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

BIRDS

Map ID	Subelement	Species	Mapping Qualifier	MDVA												Monthly Presence														
				S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	J	F	M	A	M	J	J	A	S	O	N	D
34	Waterfowl	Scoter	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
150	Wading	Great blue heron	Nesting	-	-	10 Nests	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
232	Waterfowl	Canada goose	Nesting	-	-	Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Mallard	Nesting	-	-	Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
798	Raptor	Bald eagle	Nesting	-	-	1 Nest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
849	Waterfowl	Bufflehead	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Canada goose	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Canvasback	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Mallard	Wintering	-	-	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Mergansers	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Waterfowl	Scaup	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

REPTILES & AMPHIBIANS

Map ID	Subelement	Species	Mapping Qualifier	MDVA												Monthly Presence														
				S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	J	F	M	A	M	J	J	A	S	O	N	D
1762	Turtle	Diamondback terrapin	Concentration Area	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

INVERTEBRATES

Map ID	Subelement	Species	Mapping Qualifier	MDVA												Monthly Presence														
				S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	J	F	M	A	M	J	J	A	S	O	N	D
1834	Crab	Blue crab	Highly Abundant-Wint	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

BIRDS

Subelement	Species	MDVA			Monthly Presence																
		Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mg.(S)	Mg.(F)	Molt
Diving	Common loon	Wintering			10S													-	-	-	-
	D. crested cormorant	Wintering			100S													Mar-May	Aug-Nov	-	-
Grebens		Wintering			Present													-	-	-	-
Red-throated loon		Wintering			10S													-	-	-	-
Waterfowl	American black duck	Wintering			10S													-	-	-	-
	Brant	Wintering			10S													-	-	-	-
Bufflehead		Wintering			1,000S												-	-	-	-	
Canada goose		Wintering			10,000S												-	-	-	-	
Canvasback		Wintering			1,000S												-	-	-	-	
Long-tailed duck		Wintering			10S												-	-	-	-	
Mallard		Wintering			100S												-	-	-	-	
Mergansers		Wintering			1,000S												-	-	-	-	
Ruddy duck		Wintering			1,000S												-	-	-	-	
Scaup		Wintering			1,000S												-	-	-	-	
Scoters		Wintering			10S												-	-	-	-	
Tundra swan		Wintering			100S												-	-	-	-	

FISH

Subelement	Species	MDVA			Monthly Presence																	
		Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
Diadromous	Alewife	Migration			-												-	-	-	-	Aug-Nov	Feb-Apr
	American shad	Migration			-												-	-	-	-	Aug-Nov	Feb-Jun
Atlantic sturgeon	General Distribution	-E	E	-													-	-	-	-	Jan-Dec	-
Atlantic sturgeon	Migration	-E	E	-													-	-	-	-	Jun-Nov	-
Blueback herring	Migration	-	-														-	-	-	-	Mar-Jun	-
Hickory shad	Migration	-	-														-	-	-	-	Jun-Sep	Feb-May
Shortnose sturgeon	General Distribution	E/E	E	-													-	-	-	-	Jan-Dec	Jan-Dec
Striped bass	Nursery Area																-	-	-	-	Mar-May	Jan-Dec

REPTILES & AMPHIBIANS

Subelement	Species	MDVA			Monthly Presence																	
		Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles	Adults
Turtle	Green sea turtle	General Distribution	T/T	T	Rare												-	-	-	-	May-Dec*	-
	Green sea turtle	General Distribution	T/T	T	Common												-	-	-	-	Jul-Oct	-

INVERTEBRATES

Subelement	Species	MDVA			Monthly Presence																	
		Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
Crab	Concentration Area	Highly Abundant															-	-	-	-	Jun-Mar*	Jun-Mar*
	Concentration Area	Highly Abundant-Summ															-	-	-	-	Jun-Oct	Jun-Oct

MARINE MAMMALS

Subelement	Species	Mapping Qualifier	MD/VA						Monthly Presence								
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D
Manatee	West Indian manatee	General Distribution	-E	E	Rare	-	-	-	-	-	-	-	-	-	-	-	-

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)

BIRDS

Subelement	Species	Mapping Qualifier	MD/VA						Monthly Presence								
			J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)
Diving	Brown pelican	General Distribution	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	D. crested cormorant		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Gull/Tern	Gulls	General Distribution	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Terns		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Subelement	Species	Mapping Qualifier	MD/VA						Monthly Presence									
			J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles
Diadromous	Alewife	General Distribution	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Apr	Feb-Apr	Feb-Apr	Feb-Oct
	American eel		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Feb-Apr
	Blueback herring		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Striped bass		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Jun
Estuarine Nursery	Atlantic croaker	General Distribution	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Atlantic herring		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Jun
	Atlantic menhaden		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Black drum		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Bluefish		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Kingfishes		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Northern puffer		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Oct
	Pigfish		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Nov
	Red drum		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Spot		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Nov-Apr
	Spotted seatrout		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Summer flounder		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Weakfish		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	White perch		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Estuarine Resident	Pipefish	General Distribution	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Silversides		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Silversides		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Anchoovies		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Blue catfish		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Bullhead catfish		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Spottail shiner		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Sunfish		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Northern searobin		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Sep
Marine Benthic			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Sep

FISH (continued)

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Dec-Jun
Marine Pelagic	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Nov
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Lined seahorse	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Spanish mackerel	-	-	-	-	-	-	-	-	-	-	-	-	May-Sep

REPTILES & AMPHIBIANS

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Turtle	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct
	Diamondback terrapin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Bivalve	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Stout razor clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Crab	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Brown shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Mantis shrimp	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	White shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct

MARINE MAMMALS

Subelement	Species	Monthly Presence												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
Dolphin	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES


ESI POLYGON HABITAT TYPES		Habitat Classification	Area (Acres)	Area (Sq. Miles)
ESI Rank				
10A		Salt and Brackish Water Marshes	581.39	0.91
10B		Freshwater Marshes	56.28	0.09
10C		Swamps	217.48	0.34
10D		Scrub and Shrub Wetlands	138.46	0.22
7		Exposed Tidal Flats	4.12	0.01

ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification	Length (Meters)	Length (Miles)	% of ESI Shoreline
ESI Rank					
10A		Salt and Brackish Water Marshes	66,826.05	41.52	47%
10C		Swamps	341.63	0.21	< 1%
9A		Sheltered Tidal Flats	2,554.59	1.59	2%
9B		Vegetated Low Banks	37,484.22	23.29	26%
8B		Sheltered, Solid Man-Made Structures	10,616.24	6.60	7%
8C		Sheltered Riprap	2,166.71	1.35	2%
7		Exposed Tidal Flats	1,917.64	1.19	1%
6B		Riprap	7,223.00	4.49	5%
4		Coarse Grained Sand Beaches	7,892.59	4.90	6%
3B		Scars and Steep Slopes (Sand)	756.39	0.47	1%
1B		Exposed, Solid Man-Made Structures	4,284.38	2.66	3%
Total ESI Shoreline:		142,063.44	Total ESI Shoreline:	88.27	
Total Shoreline:		97,937.30	Total Shoreline:	60.86	

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 67C Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	Mapping Qualifier	MDVA												Monthly Presence											
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Ephemeral	High	Low						
SAV	Submerged aquatic veg	High Ecological Value	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
SAV	Submerged aquatic veg	High Ecological Value	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
SAV	Submerged aquatic veg	High Ecological Value	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

BIRDS

Map ID	Subelement	Species	Mapping Qualifier	MDVA												Monthly Presence											
				S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt					
107	Wading	Great blue heron	Nesting	-	-	22 Nests	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
150	Wading	Great blue heron	Nesting	-	-	10 Nests	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
232	Watervowl	Canada goose	Nesting	-	-	Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Watervowl	Mallard	Nesting	-	-	Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
798	Raptor	Bald eagle	Nesting	-	-	1 Nest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
841	Watervowl	American black duck	Wintering	-	-	Present	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Watervowl	Bufflehead	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Watervowl	Canada goose	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Watervowl	Canvasback	Wintering	-	-	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Watervowl	Mallard	Wintering	-	-	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Watervowl	Mergansers	Wintering	-	-	100S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Watervowl	Tundra swan	Wintering	-	-	10S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Map ID	Subelement	Species	Mapping Qualifier	MDVA												Monthly Presence											
				S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Eggs	Larvae	Juveniles	Adults					
1638	Diadromous	Striped bass	Migration	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

REPTILES & AMPHIBIANS

Map ID	Subelement	Species	Mapping Qualifier	MDVA												Monthly Presence											
				S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Eggs	Larvae	Juveniles	Adults					
1762	Turtle	Diamondback terrapin	Concentration Area	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

INVERTEBRATES

Map ID	Subelement	Species	Mapping Qualifier	MDVA												Monthly Presence											
				S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Eggs	Larvae	Juveniles	Adults					
1789	Crab	Blue crab	Highly Abundant	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1834	Crab	Blue crab	Highly Abundant-Wint	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

WIDESPREAD IN MAPPED AREA (> 10 SQUARE KILOMETERS)

BIRDS

Subelement	Species	MDVA			Monthly Presence																
		Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mg.(S)	Mg.(F)	Molt
Diving	Common loon	Wintering			10S														-	-	-
	D. crested cormorant	Wintering			100S														Mar-May	Aug-Nov	-
Grebens		Wintering			Present														-	-	-
Red-throated loon		Wintering			10S														-	-	-
Waterfowl	American black duck	Wintering			10S														-	-	-
	Brant	Wintering			10S														-	-	-
Bufflehead		Wintering			1,000S													-	-	-	-
Canada goose		Wintering			10,000S													-	-	-	-
Canvasback		Wintering			1,000S													-	-	-	-
Long-tailed duck		Wintering			10S													-	-	-	-
Mallard		Wintering			100S													-	-	-	-
Mergansers		Wintering			1,000S													-	-	-	-
Ruddy duck		Wintering			1,000S													-	-	-	-
Scaup		Wintering			1,000S													-	-	-	-
Scoters		Wintering			10S													-	-	-	-
Tundra swan		Wintering			100S													-	-	-	-

FISH

Subelement	Species	MDVA			Monthly Presence																	
		Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
Diadromous	Alewife	Migration			-													-	-	Aug-Nov	Feb-Apr	
	American shad	Migration			-													-	-	Aug-Nov	Feb-Jun	
Atlantic sturgeon	General Distribution	-E	E	-														-	-	-	Jan-Dec	
Atlantic sturgeon	Migration	-E	E	-														-	-	-	Jun-Nov	
Blueback herring	Migration	-																-	-	Aug-Nov	Mar-Jun	
Hickory shad	Migration	-																-	-	Jun-Sep	Feb-May	
Shortnose sturgeon	General Distribution	E/E	E	-														-	-	Jan-Dec	Jan-Dec	
Striped bass	Nursery Area																	-	Mar-May	Jan-Dec	-	

REPTILES & AMPHIBIANS

Subelement	Species	MDVA			Monthly Presence																	
		Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Hatch	Internest	Juveniles	Adults
Turtle	Green sea turtle	General Distribution	T/T	T	Rare												-	-	-	May-Dec*	-	
	Green sea turtle	General Distribution	T/T	T	Common												-	-	-	Jul-Oct	-	

INVERTEBRATES

Subelement	Species	MDVA			Monthly Presence																	
		Mapping Qualifier	S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
Crab	Blue crab	Highly Abundant-Summ															-	-	-	Jun-Oct	Jun-Oct	
		Concentration Area															-	-	-	Jun-Oct	Jun-Oct	

MARINE MAMMALS

Subelement	Species	Mapping Qualifier	MD/VA	S F	Concentration	J F M A M J J A S O N D	Mating	Calving	Pupping	Molt
Manatee	West Indian manatee	General Distribution	-IE	E	Rare	-	-	-	-	-

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)

BIRDS

Subelement	Species	Monthly Presence											
		J	F	M	A	M	J	J	A	S	O	N	D
Diving	Brown pelican	-	-	-	-	-	-	-	-	-	-	-	-
	D. crested cormorant	-	-	-	-	-	-	-	-	-	-	-	-
Gull/Tern	Gulls	-	-	-	-	-	-	-	-	-	-	-	-
	Terns	-	-	-	-	-	-	-	-	-	-	-	-
Waterfowl	Wood duck	-	-	-	-	-	-	-	-	-	-	-	-

FISH

Subelement	Species	Monthly Presence											
		J	F	M	A	M	J	J	A	S	O	N	D
Diacromous	Alewife	-	-	-	-	-	-	-	-	-	-	-	-
	American eel	-	-	-	-	-	-	-	-	-	-	-	-
	Blueback herring	-	-	-	-	-	-	-	-	-	-	-	-
	Gizzard shad	-	-	-	-	-	-	-	-	-	-	-	-
	Striped bass	-	-	-	-	-	-	-	-	-	-	-	-
Estuarine Nursery	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-
	Atlantic croaker	-	-	-	-	-	-	-	-	-	-	-	-
	Atlantic herring	-	-	-	-	-	-	-	-	-	-	-	-
	Atlantic menhaden	-	-	-	-	-	-	-	-	-	-	-	-
	Black drum	-	-	-	-	-	-	-	-	-	-	-	-
	Black sea bass	-	-	-	-	-	-	-	-	-	-	-	-
	Bluefish	-	-	-	-	-	-	-	-	-	-	-	-
	Kingfishes	-	-	-	-	-	-	-	-	-	-	-	-
	Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-
	Northern puffer	-	-	-	-	-	-	-	-	-	-	-	-
	Pigfish	-	-	-	-	-	-	-	-	-	-	-	-
	Red drum	-	-	-	-	-	-	-	-	-	-	-	-
	Sheepshead	-	-	-	-	-	-	-	-	-	-	-	-
	Spot	-	-	-	-	-	-	-	-	-	-	-	-
	Spotted seatrout	-	-	-	-	-	-	-	-	-	-	-	-
	Summer flounder	-	-	-	-	-	-	-	-	-	-	-	-
	Weakfish	-	-	-	-	-	-	-	-	-	-	-	-
	White perch	-	-	-	-	-	-	-	-	-	-	-	-
	Winter flounder	-	-	-	-	-	-	-	-	-	-	-	-
Estuarine Resident	Pipefish	-	-	-	-	-	-	-	-	-	-	-	-
	Silversides	-	-	-	-	-	-	-	-	-	-	-	-
	Silversides	-	-	-	-	-	-	-	-	-	-	-	-
Fish	Anchoovies	-	-	-	-	-	-	-	-	-	-	-	-

FISH (continued)

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
	Striped burrfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Blue catfish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Freshwater	Atlantic spadefish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Cleannose skate	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Sep
	Red hake	-	-	-	-	-	-	-	-	-	-	-	-	Sep-May*
	Silver hake	-	-	-	-	-	-	-	-	-	-	-	-	Oct-May
	Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	Jul-Nov
Marine Pelagic	Cobia	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
	Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Lined seahorse	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
	Spanish mackerel	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
		-	-	-	-	-	-	-	-	-	-	-	-	May-Sep

REPTILES & AMPHIBIANS

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Turtle	Diamondback terrapin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Bivalve	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Stout razor clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Cephalopod	Squid	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Brown shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Mantis shrimp	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	White shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct

MARINE MAMMALS

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Dolphin	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES


ESI POLYGON HABITAT TYPES		Habitat Classification		Area (Acres)	Area (Sq. Miles)
ESI Rank					
10A		Salt and Brackish Water Marshes		266.40	0.42
10B		Freshwater Marshes		16.80	0.03
10C		Swamps		333.65	0.52
10D		Scrub and Shrub Wetlands		94.93	0.15
9A		Sheltered Tidal Flats		5.81	0.01
7		Exposed Tidal Flats		8.45	0.01

ESI SHORELINE HABITAT TYPES		Shoreline Habitat Classification		Length (Meters)	Length (Miles)	% of ESI Shoreline
ESI Rank						
10A		Salt and Brackish Water Marshes		4,1710.92	25.92	39%
10C		Swamps		2,802.68	1.74	3%
10D		Scrub and Shrub Wetlands		1,161.89	0.72	1%
9A		Sheltered Tidal Flats		826.37	0.51	1%
9B		Vegetated Low Banks		29,221.80	18.16	28%
8B		Sheltered, Solid Man-Made Structures		6,778.82	4.21	6%
8C		Sheltered Riprap		2,383.30	1.48	2%
7		Exposed Tidal Flats		1,646.02	1.02	2%
6B		Riprap		5,687.32	3.53	5%
4		Coarse Grained Sand Beaches		8,186.40	5.09	8%
3B		Scarps and Steep Slopes (Sand)		1,810.89	1.13	2%
1B		Exposed, Solid Man-Made Structures		3,522.55	2.19	3%
		Total ESI Shoreline:		105,738.97	Total ESI Shoreline:	65.70
		Total Shoreline:		71,754.39	Total Shoreline:	44.59

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%

All underlying GIS data can be obtained from response.restoration.noaa.gov





Map 67D
Central Bay



SEE BACK OF MAP
for details about mapped species and
other species that occur in mapped area.
Data Published: September 2016

0 Not for Navigation 1 Miles
0 1 Kilometers 1:50,000



Map 67D Central Bay

BIOLOGICAL RESOURCES

Note: An asterisk (*) indicates that life stage occurs in this range but not in all months included

DISPLAYED ON MAP

BENTHIC

Subelement	Species	Mapping Qualifier	MD/VA	S	F	Concentration	Monthly Presence
SAV	Submerged aquatic veg	High Ecological Value	J F M A M J J A S O N D	J F M A M J J A S O N D	J F M A M J J A S O N D	Ephemeral	J F M A M J J A S O N D

BIRDS

Map ID	Subelement	Species	Mapping Qualifier	MD/VA	S	F	Concentration	Monthly Presence	
				J F M A M J J A S O N D	J F M A M J J A S O N D	Nest	Mig.(S)	Mig.(F)	Molt
143	Wading	Great blue heron	Nesting		7 Nests				-
148	Wading	Great blue heron	Nesting		32 Nests				-
149	Wading	Great blue heron	Nesting		18 Nests				-
166	Wading	Great blue heron	Nesting		11 Nests				-
232	Waterfowl	Canada goose	Nesting						Jun-Jul
	Waterfowl	Mallard	Nesting						Jul-Aug
679	Waterfowl	Brant	Wintering						-
	Waterfowl	Bufflehead	Wintering						-
	Waterfowl	Canada goose	Wintering						-
	Waterfowl	Canvasback	Wintering						-
	Waterfowl	Mallard	Wintering						-
	Waterfowl	Mergansers	Wintering						-
	Waterfowl	Scaup	Wintering						-
798	Raptor	Bald eagle	Nesting		1 Nest				Dec-Jun
848	Waterfowl	American black duck	Wintering		10S				-
	Waterfowl	Bufflehead	Wintering						-
	Waterfowl	Canada goose	Wintering						-
	Waterfowl	Canvasback	Wintering						-
	Waterfowl	Mallard	Wintering						-
	Waterfowl	Mergansers	Wintering						-
	Waterfowl	Ruddy duck	Wintering						-
	Waterfowl	Scaup	Wintering						-
	Waterfowl	Tundra swan	Wintering						-
849	Waterfowl	Bufflehead	Wintering						-
	Waterfowl	Canada goose	Wintering						-
	Waterfowl	Canvasback	Wintering						-
	Waterfowl	Mallard	Wintering						-
	Waterfowl	Mergansers	Wintering						-
	Waterfowl	Scaup	Wintering						-
	Waterfowl	Common loon	Wintering						-
1023	Diving	D. crested cormorant	Wintering						Mar-May Aug-Nov
	Diving	Diving	Wintering						-

BIRDS (continued)

Map ID	Subelement	Species	MD/VA			Monthly Presence															
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
	Diving	Red-throated loon			Wintering													-	-	-	-
	Watertowl	Long-tailed duck			Wintering													-	-	-	-
	Watertowl	Scoters			Wintering													-	-	-	-
1059	Watertowl	American black duck			Wintering												-	-	-	-	-
	Watertowl	Ruddy duck			Wintering												-	-	-	-	-
	Watertowl	Tundra swan			Wintering												-	-	-	-	-
1173	Diving	Grebes			Wintering												-	-	-	-	-
	Watertowl	American black duck			Wintering												-	-	-	-	-
	Watertowl	Bufflehead			Wintering												-	-	-	-	-
	Watertowl	Canada goose			Wintering												-	-	-	-	-
	Watertowl	Canvasback			Wintering												-	-	-	-	-
	Watertowl	Long-tailed duck			Wintering												-	-	-	-	-
	Watertowl	Mallard			Wintering												-	-	-	-	-
	Watertowl	Mergansers			Wintering												-	-	-	-	-
	Watertowl	Ruddy duck			Wintering												-	-	-	-	-
	Watertowl	Tundra swan			Wintering												-	-	-	-	-

FISH

Map ID	Subelement	Species	MD/VA			Monthly Presence															
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Eggs	Larvae	Juveniles
1556	Diadromous	Atlantic sturgeon			-IE												-	-	-	-	-
	Diadromous	Shortnose sturgeon			General Distribution	E/E	E										-	-	-	-	-
	Diadromous	Striped bass			Nursery Area												-	Mar-May	Jan-Dec	-	-
1575	Diadromous	Atlantic sturgeon			General Distribution	-IE	E										-	-	-	-	-
	Diadromous	Shortnose sturgeon			General Distribution	E/E	E										-	-	-	-	-
1601	Diadromous	Atlantic sturgeon			Migration	-IE	E										-	-	-	-	-
	Diadromous	Hickory shad			Migration												-	Jun-Sep	Feb-May	-	-
1621	Diadromous	Alewife			Migration												-	-	-	-	-
	Diadromous	American shad			Migration												-	Aug-Nov	Feb-Jun	-	-
	Diadromous	Blueback herring			Migration												-	Aug-Nov	Feb-Jun	-	-
1638	Diadromous	Striped bass			Migration												-	-	-	-	-

REPTILES & AMPHIBIANS

Map ID	Subelement	Species	MD/VA			Monthly Presence															
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Eggs	Larvae	Juveniles
		Diamondback terrapin			Concentration Area												May-Aug	-	-	-	-
1762	Turtle	Green sea turtle			General Distribution	T/T	T										-	-	-	-	-
1767	Turtle	Green sea turtle			General Distribution	T/T	T										-	-	-	-	-
	Turtle				Common												-	-	-	-	-
1638					Rare												-	-	-	-	-

INVERTEBRATES

Map ID	Subelement	Species	MD/VA			Monthly Presence															
			S	F	Concentration	J	F	M	A	M	J	J	A	S	O	N	D	Nest	Eggs	Larvae	Juveniles
1788	Crab	Blue crab			Highly Abundant-Summ												-	-	-	-	-
1814	Bivalve	Eastern elliptio			Vulnerable Occurrence												-	May-Dec*	-	-	-

Central Bay: Map 67D

MARINE MAMMALS

Map ID	Subelement	Species	MD/VA												Monthly Presence												
			Mapping Qualifier			S F			Concentration			J F M A M J J A S O N D			Mating			Calving			Pupping			Molt			
		General Distribution	-E	E	Rare																						
1844	Manatee	West Indian manatee																									

ALSO PRESENT IN MAPPED AREA (GENERAL DISTRIBUTION)

BIRDS

Subelement	Species	Monthly Presence															
		J	F	M	A	M	J	J	A	S	O	N	D	Nest	Mig.(S)	Mig.(F)	Molt
Diving	Brown pelican																
	D. crested cormorant																
Gull/Tern	Gulls																
	Terns																
Waterfowl	Wood duck																

FISH

Subelement	Species	Monthly Presence																
		J	F	M	A	M	J	J	A	S	O	N	D	Spawn	Eggs	Larvae	Juveniles	Adults
Diacromous	Alewife													Feb-Apr	Feb-Apr	Feb-Apr	Feb-Oct	Feb-Apr
	American eel																Jan-May	Jan-Dec
	American shad													Feb-Jun	Feb-Jun	Mar-Jul	Mar-Oct	Feb-Jun
	Blueback herring													Mar-Jun	Apr-Jun	May-Oct	Mar-Jun	Mar-Dec
	Gizzard shad													Mar-Jun	-	Jan-Dec	Jan-Dec	Jan-Dec
	Striped bass													-	-	Apr-Dec	Jan-Dec	May-Oct
Estuarine Nursery	Atlantic croaker													-	-	Dec-Apr	Jan-Dec	May-Nov
	Atlantic croaker													-	-	-	-	Jan-Jun
	Atlantic herring													-	-	-	-	Jan-Jun
	Atlantic menhaden													-	-	-	-	Jan-Dec
	Black drum													-	-	-	-	Apr-Dec
	Black sea bass													-	-	-	-	Apr-Nov
	Bluefish													-	-	-	-	May-Oct
	Kingfishes													-	-	-	-	Apr-Nov
	Northern puffer													-	-	-	-	Mar-Nov
	Northern puffer													-	-	-	-	Mar-Oct
	Pigfish													-	-	-	-	Apr-Nov
	Red drum													-	-	-	-	May-Nov
	Sheepshead													-	-	-	-	Apr-Nov
	Spot													-	-	-	-	Nov-Apr
	Spotted seatrout													-	-	-	-	Apr-Nov
	Summer flounder													-	-	-	-	Feb-May
	Weakfish													-	-	-	-	Jun-Dec
	White perch													-	-	-	-	Jan-Dec
Estuarine Resident	Winter flounder													-	-	-	-	Dec-Mar
	Pipefish													-	-	-	-	Jan-Dec
	Silversides													-	-	-	-	Jan-Dec
	Silversides													-	-	-	-	Mar-Nov

FISH (continued)

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Fish	Anchovies	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Striped burrfish	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
Freshwater	Blue catfish	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Marine Benthic	Atlantic spadefish	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
	Cleasnose skate	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Cownose ray	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Sep
Northern searobin	-	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
Red hake	-	-	-	-	-	-	-	-	-	-	-	-	-	Sep-May*
Silver hake	-	-	-	-	-	-	-	-	-	-	-	-	-	Oct-May
Spotted hake	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Butterfish	-	-	-	-	-	-	-	-	-	-	-	-	-	Mar-Nov
Cobia	-	-	-	-	-	-	-	-	-	-	-	-	-	Jun-Sep
Harvestfish	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Nov
Lined seahorse	-	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Spanish mackerel	-	-	-	-	-	-	-	-	-	-	-	-	-	May-Sep

REPTILES & AMPHIBIANS

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Turtle	Diamondback terrapin	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Oct

INVERTEBRATES

Subelement	Species	Monthly Presence												Adults
		J	F	M	A	M	J	J	A	S	O	N	D	
Bivalve	Eastern oyster	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Softshell clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	Stout razor clam	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Cephalopod	Squid	-	-	-	-	-	-	-	-	-	-	-	-	Apr-Dec
Crab	Blue crab	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
Shrimp	Brown shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct
	Mantis shrimp	-	-	-	-	-	-	-	-	-	-	-	-	Jan-Dec
	White shrimp	-	-	-	-	-	-	-	-	-	-	-	-	May-Oct


MARINE MAMMALS

Subelement	Species	Monthly Presence												Molt
		J	F	M	A	M	J	J	A	S	O	N	D	
Dolphin	Bottlenose dolphin	-	-	-	-	-	-	-	-	-	-	-	-	-

For additional information about species locations and extent, reference the underlying GIS data available from response.restoration.noaa.gov

SHORELINE RESOURCES

ESI POLYGON HABITAT TYPES	
	Habitat Classification
10A	Salt and Brackish Water Marshes
10B	Freshwater Marshes
10C	Swamps
10D	Scrub and Shrub Wetlands



	Area (Acres)	Area (Sq. Miles)
89.29	0.14	
279.16	0.44	
1,541.07	2.41	
135.06	0.21	

	Length (Meters)	% of ESI Shoreline
19,587.49	23%	
14,729.51	17%	
13,425.51	16%	
377.31	< 1%	
19,813.72	23%	
9,499.60	11%	
1,702.80	2%	
288.10	< 1%	
3,591.13	4%	
25.60	< 1%	
2,516.28	3%	
748.31	1%	
75.08	< 1%	

Total ESI Shoreline: 86,380.44 Total ESI Shoreline: 53.67
 Total Shoreline: 63,240.50 Total Shoreline: 39.30

Note: A shoreline segment may include multiple shoreline habitats. If any segments include multiple habitats, the combined length of all habitats may exceed the length of the mapped shoreline, and the percent of ESI shoreline values will sum to greater than 100%


All underlying GIS data can be obtained from response.restoration.noaa.gov



Human-Use Features

 AQ	Aquaculture	 Park
 	Airport	 Port
 	Anchorage	 Recreational Fishing
 	Artificial Reef	 Repeated Measurement Site
 	Abandoned Vessel	 State Protected
 	Beach	 Tribal Land
 	Boat Ramp	 Water Intake
 	Commercial Fishing	 Wildlife Refuge
 	Critical Habitat	 US Highway
 	EPA Facility	 State Highway or Route
 	Ferry	 Interstate
 	Heliport	 Road
 	Historical Site	 Shipping Lane
 	Lock and Dam	 Rail Route
 	Marina	 Pipeline
 	Military	 Ferry Route
 	Management Area	 Management Area
 	National Estuarine Research Reserve	 Map ID
 	National Park	
 	Nature Conservancy	

ESI Shoreline and Habitat Ranking



Examples of Double and Triple Shoreline Rankings:



Shorelines often contain varied geomorphology, and therefore may require two or three ESI types to describe. These symbols will look similar to the examples above. The first shoreline type listed is the most landward shore type.

Sensitive Biological Resources

 Birds	 Invertebrates	 Terrestrial Mammals
● Birds	● Invertebrates	 Small Mammal
 Alcid/Pelagic	 Crab/Invertebrate/Shellfish	 Ungulate
 Diving	 Bivalve/Barnacle/Chordate	 Bat
 Gull/Tern/Bird	 Cephalopod	 Habitats
 Passerine	 Insect	 Upland/Wetland/Plant
 Raptor	 Gastropod	 Reptiles &
 Shorebird	 Lobster	 Amphibian
 Wading	 Shrimp	 Turtle
 Waterfowl	 Marine Mammals	 Threatened or Endangered Species
 Fish	 Dolphin	 Map ID
 Fish	 Pinniped	
 Nursery	 Whale	
 Fish	 Manatee	

Guidelines for Interpreting Environmental Sensitivity Index (ESI) Maps

The following guidelines may help map users interpret the ESI maps. Additional information about resources mapped for this atlas can be found in the ESI Intro pages and associated metadata.

Sensitive Biological Resources (1:50,000 scale maps): Species are arranged into eight major categories or "elements" and further subdivided into "subelement" groupings representing species that share similar lifestyle characteristics and risks to oiling.

Biological resources may be mapped as points, lines, and/or polygons. Each element is represented by a unique color and/or hatch pattern, and icons illustrate the subelement(s) found within each feature. If a species is state or federally listed as threatened or endangered, a solid red square is placed behind the icon. When multiple elements occur in the same location, overlapping hatch patterns will be shown.

The associated Map ID "links" to the map report where the species and attributes are listed. To maximize readability, biological polygons covering more than 10 square km are not displayed on the map. Instead, these species are listed in the "Widespread in Mapped Area" section of the report. Occurrences of non-listed species assigned a "General Distribution" mapping qualifier are listed under "Also Present in Mapped Area" regardless of polygon size.

Shoreline Habitat Resources (all maps): The shoreline was mapped at mean-high water, then classified based on vulnerability to spilled oil and ease of clean-up. Shorelines are ranked on a scale from 1 (least vulnerable) to 10 (most vulnerable). Cooler colors represent less vulnerable shoreline types; warm and hot colors indicate increased vulnerability. A shoreline may have more than one habitat type present. When this happens, the most landward shoreline type is mapped on the shoreline, and the more seaward type(s) are mapped adjacent to the water. The areal extent of intertidal and marsh habitats may also be mapped as polygons. Most often these polygons represent tidal flats and marshes, though in particularly rocky shored areas, these may include exposed, wave-cut platforms and sand and gravel beaches.

Human-Use Features (1:100,000 scale maps): Locations of human-use features and jurisdictional boundaries are mapped as points, lines, or polygons. As with biological features, the human-use features appearing on the map include Map ID which "links" to a description of the feature in the map report. Additional features, particularly jurisdictional and management boundaries that cover the majority or entirety of the mapped area, are listed in the report, but not shown on the map.

