

COOK INLET/ KENAI PENINSULA, ALASKA

SHORELINE TYPES

- 1 Exposed rocky shores
- 2 Exposed wave-cut platforms
- 3 Fine-grained sand beaches
- 4 Coarse-grained sand beaches
- 5 Exposed tidal flats
- 6 Mixed sand and gravel beaches
- 7 Gravel beaches
- 8 Sheltered rocky shores
- 9 Sheltered tidal flats
- 10 Marshes

BIOLOGICAL RESOURCES

- MARINE MAMMALS**

 - Seals, sea lions and others
 - Whales and dolphins

MARINE BIRDS


 - Birds of prey
 - Diving birds
 - Seabirds
 - Shorebirds
 - Wading birds
 - Waterfowl

SHELLFISH

 - Bivalve mollusks
 - Crabs
 - Shrimp

FISHES

 - Anadromous fishes
 - Nonanadromous fishes



SOCIOECONOMIC FEATURES

- Parks
- Marinas
- Boat ramps

WATERFOWL
STAGING AREA
SPRING, FALL

Sugarloaf
Mountain

RIGHT ARM

INISKIN
BAY

WATERFOWL NESTING AND STAGING AREAS ARE
ASSOCIATED WITH COASTAL WETLANDS AND ARE
GENERALLY CONFINED WITHIN THESE HABITATS

SEE MAP #33 FOR SPECIES
IN NEARSHORE AREA

WATERFOWL
STAGING AREA
SPRING, FALL

Mapped, edited, and published by the Geological Survey
Control by USGS and USC&GS

rpi inc.
Research Planning Institute, Inc.
925 Gervais Street
Columbia, South Carolina, USA 29201
(803) 256-7322

TRUE NORTH
MAGNETIC NORTH
APPROXIMATE MEAN
DECLINATION, 1958

SCALE 1:63,360

CONTOUR INTERVAL 100 FEET

DATUM IS MEAN SEA LEVEL

DEPTH CURVES AND SOUNDINGS IN FEET - DATUM IS MEAN LOWER LOW WATER

SHORELINE SHOWN REPRESENTS THE APPROXIMATE LINE OF MEAN HIGH WATER

THE MEAN RANGE OF TIDE IS APPROXIMATELY 12 FEET

FOR SALE BY U. S. GEOLOGICAL SURVEY
FAIRBANKS, ALASKA 99701, DENVER, COLORADO 80225, OR WASHINGTON, D. C. 20242
A FOLDER DESCRIBING TOPOGRAPHIC MAPS AND SYMBOLS IS AVAILABLE ON REQUEST

ADJOINS MAP #41,42

**ESI MAP
#32**

ILIAMNA (C-2), ALASKA

N5930-W15322.5/15X22.5

1958

MINOR REVISIONS 1972

ADJOINS MAP #33,26