


COOK INLET/  
KENAI PENINSULA, ALASKA

SHORELINE TYPES

- Sensitivity
- 1 Exposed rocky shores
  - 2 Exposed wave-cut platforms
  - 3 Fine-grained sand beaches
  - 4 Coarse-grained sand beaches
  - 5 Exposed tidal flats
  - 6 Mixed sand and gravel beaches
  - 7 Gravel beaches
  - 8 Sheltered rocky shores
  - 9 Sheltered tidal flats
  - 10 Marshes

BIOLOGICAL RESOURCES


- MARINE MAMMALS
- Seals, sea lions, and otters
  - Whales and dolphins
- MARINE BIRDS
- Birds of prey
  - Diving birds
  - Seabirds
  - Shorebirds
  - Wading birds
  - Waterfowl
- SHELLFISH
- Bivalve molluscs
  - Crabs
  - Shrimp
- FISHES
- Anadromous fishes
  - Nonanadromous fishes



SOCIOECONOMIC FEATURES

- Parks
- Marinas
- Boat ramps

PRESENT IN NEARSHORE AREA



GEORGIEVSK  
REDOUBT  
STATE HISTORICAL  
PARK

WATERFOWL  
STAGING AREA  
SPRING, FALL

WATERFOWL NESTING AND STAGING AREAS ARE  
ASSOCIATED WITH COASTAL WETLANDS AND ARE  
GENERALLY CONFINED WITHIN THESE HABITATS

**rpi inc.**  
Research Planning Institute, Inc.  
925 Cervais Street  
Columbia, South Carolina, USA 29201  
(803) 256-7322

predetermined by the Bureau of Land Management  
Folio S-16, Seward Meridian  
Swamps, as portrayed, indicate only the wetter areas,  
usually of low relief, as interpreted from aerial photographs

CONTOUR INTERVAL 50 FEET  
DOTTED LINES REPRESENT 25 FOOT CONTOURS  
NATIONAL GEODETIC VERTICAL DATUM OF 1929  
DEPTH CURVES IN FEET-DATUM IS MEAN LOWER LOW WATER  
SHORELINE SHOWN REPRESENTS THE APPROXIMATE LINE OF MEAN HIGH WATER  
THE AVERAGE RANGE OF TIDE IS APPROXIMATELY 17 FEET

FOR SALE BY U.S. GEOLOGICAL SURVEY  
FAIRBANKS, ALASKA 99701, DENVER, COLORADO 80225, OR RESTON, VIRGINIA 22092  
A FOLDER DESCRIBING TOPOGRAPHIC MAPS AND SYMBOLS IS AVAILABLE ON REQUEST

ADJOINS MAP # 23

ESI MAP  
#20

KENAI (B-4), ALASKA  
N6015-W15107.5/15X22.5

1951  
LIMITED REVISIONS 1980