



SPECIES PRESENT  
IN NEARSHORE AREAS

- 18,19  
26,38
- 15,20-  
23,27  
28,39-  
42
- 17,25  
43-49
- 4-7  
9,10  
50-64
- 2,3  
55,56
- 57-65
- 66
- 12,14  
67-70
- 11,13
- 29-31
- 3,5,8  
12-14  
16-20  
22,26  
27-42
- 1,2,6  
7,10  
43

**ST. MARYS RIVER  
SHORELINE TYPES**

- 1 Exposed bedrock bluffs
- 2 Exposed unconsolidated sediment bluffs
- 3 Shelving bedrock shores
- 4 Sand shores
- 5 Mixed sand and gravel shores
- 6 Gravel shores
- 7 Riprap and harbor structures
- 8 Sheltered bluffs (not present)
- 9 Low vegetated banks
- 10 A Fringing wetlands
- 10 B Extensive wetlands

**BIOLOGICAL RESOURCES**

- BIRDS**
  - Diving birds
  - Gulls and terns
  - Raptors
  - Shorebirds
  - Wading birds
  - Waterfowl
- FINFISH**
  - Anadromous fish
  - Freshwater fish
- MAMMALS**
  - Furbearers

**KEY TO WILDLIFE MARKERS**

COLOR CODED = TYPES OF ORGANISMS  
POINT LOCALITY  
DOTS = SEASONALITY  
SPRING (S) SUMMER (S) FALL (S) WINTER (S)  
SYMBOL = ECOLOGICAL TYPE  
NUMBER = SPECIES (from regional lists)  
\* = BREEDING SEASON

**SOCIOECONOMIC FEATURES**

- Parks and preserves
- Recreational beaches
- Water intakes
- Marinas
- Boat ramps
- Station locations

Mapped, edited, and published by the Geological Survey  
Control by USGS, USGS/USFS, and U.S. Lake Survey  
Topography by photogrammetric methods from aerial  
photographs taken 1954-55. Field checked 1964

**rpi inc.**  
Research Planning Institute, Inc.  
925 Gervais Street  
Columbia, South Carolina, USA 29201  
(803) 256-7322



**ESI MAP  
#11**

GOETZVILLE, MICH.-ONT.  
N4600-W8400/7.5

1964

AMS 4176 II SE-SERIES V862